

Multiple contaminations of chickens with Campylobacter, Escherichia coli and Salmonella in Yaounde (Cameroon).

Ariane Nzouankeu, Antoinette Ngandjio, Guy Ejenguele, Thomas Njine,
Marguerite Ndayo Wouafo

► **To cite this version:**

Ariane Nzouankeu, Antoinette Ngandjio, Guy Ejenguele, Thomas Njine, Marguerite Ndayo Wouafo. Multiple contaminations of chickens with Campylobacter, Escherichia coli and Salmonella in Yaounde (Cameroon).. Journal of Infection in Developing Countries, Independent, 2010, 4 (9), pp.583-686. 10.3855/jidc.1019 . pasteur-00550846

HAL Id: pasteur-00550846

<https://hal-riip.archives-ouvertes.fr/pasteur-00550846>

Submitted on 31 Dec 2010

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Multiple contaminations of chickens with *Campylobacter*, *Escherichia coli* and *Salmonella* in Yaounde (Cameroon)

Ariane Nzouankeu^{1,2}, Antoinette Ngandjio¹, Guy Ejenguele¹, Thomas Njine², Marguerite Ndayo Wouafo¹

¹Pasteur Centre of Cameroon, Yaounde, Cameroon

²University of Yaounde I, Faculty of Science, Yaounde, Cameroon

Abstract

Introduction: Food-borne diseases associated with *Campylobacter*, *Escherichia coli*, and *Salmonella* are mainly caused by the consumption of raw or undercooked poultry meat. The objective of this study was to evaluate the prevalence of *Campylobacter*, *Escherichia coli*, and *Salmonella* in chickens.

Methodology: One hundred and fifty chickens collected from eight retail markets in Yaounde were examined for the presence of *Campylobacter*, *Escherichia coli*, and *Salmonella* using standard bacteriological procedures.

Results: Of the 150 chickens collected, 135 (90%) were contaminated with *Campylobacter* (68.9% *C. coli* and 31.1% *C. jejuni*). All the chickens were positive for *E. coli*. Among the 150 isolates, 17 (11.3%) were enteropathogenic *E. coli* (EPEC). Additionally, 103 *Salmonella* strains were recovered from 90 chickens. *Salmonella* Enteritidis (45.6%) and *Salmonella* Hadar (28.1%) were the most frequent serotypes.

Multiple contamination was found in 142 chickens (94.6%), of which 83 (55.3%) were concurrently contaminated with *Campylobacter*, *Escherichia coli*, and *Salmonella*.

Conclusion: These results show that chickens in Cameroon are highly contaminated with *Campylobacter*, *Escherichia coli*, and *Salmonella*. The multiple contaminations of chickens is a potential risk of infection for consumers and highlights the necessity of public awareness for food safety.

Key words: *Campylobacter*; *Escherichia coli*; *Salmonella*; multiple contamination

J Infect Dev Ctries 2010; 4(9):583-586.

(Received 09 March 2010 - Accepted 19 April 2010)

Copyright © 2010 Nzouankeu *et al.* This is an open-access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Introduction

Contaminated, raw, or undercooked poultry products, especially chicken meat, have been shown to be a critical link in transmitting food-borne pathogens to humans, resulting in numerous cases of diseases [1]. Most of these food-borne illnesses are caused by three major bacteria: *Campylobacter*, *Salmonella*, and *E. coli* [2-3]. Infection with these bacteria usually results in a self-limiting gastroenteritis; however, young children, older adults, and the immunocompromised may experience invasive disease [4-5]. *Campylobacter* has been identified as the predominant cause of Guillain-Barre's syndrome and reactive arthritis [6]. *E. coli* is particularly responsible for different forms of diarrhoea, such as the bloody diarrhoea and hemolytic uremic syndrome caused by *E. coli* O157:H7 [7]. *Salmonella* is known to be responsible for systemic salmonellosis infections which could be life threatening.

The objective of this study was to evaluate the prevalence of *Campylobacter*, *Escherichia coli*, and *Salmonella* in chickens sold in retail markets in Yaounde, Cameroon to determine the risks associated with the chicken handling and consumption.

Methodology

Sample collection

Chickens were purchased from eight retail markets in Yaounde. These markets were selected to ensure complete coverage of all the subdivisions of the town. Sampling visits were made once a week from February 2006 to January 2007. The entire chicken carcasses were immediately placed in sterile plastic bags and transported in a cool box to the laboratory, where samples were processed within an hour.

Table 1. Co-contamination with *Campylobacter*, *E. coli* and *Salmonella* in chickens

Co-infections	Number of contaminated chickens	Total
<i>C. coli</i> + <i>E. coli</i> *+ Enteritidis (<i>C. coli</i> + EPEC + Enteritidis)	18 2	20
<i>C. coli</i> + <i>E. coli</i> * + Hadar (<i>C. coli</i> + EPEC + Hadar)	12 3	15
<i>C. coli</i> + <i>C. jejuni</i> + <i>E. coli</i> * + Enteritidis (<i>C. coli</i> + <i>C. jejuni</i> + EPEC + Enteritidis)	9 1	10
<i>C. coli</i> + <i>E. coli</i> * + <i>Salmonella.sp</i> (<i>C. coli</i> + EPEC + <i>Salmonella.sp</i>)	7 2	9
<i>C. coli</i> + <i>C. jejuni</i> + <i>E. coli</i> * + <i>Salmonella. Sp</i> (<i>C. coli</i> + <i>C. jejuni</i> + EPEC + <i>Salmonella. Sp</i>)	6 1	7
<i>C. coli</i> + <i>E. coli</i> *+ Enteritidis +Hadar (<i>C. coli</i> + EPEC + Enteritidis +Hadar)	5 1	6
<i>C. jejuni</i> + <i>E. coli</i> * + Hadar (<i>C. jejuni</i> + EPEC + Hadar)	5 1	6
<i>C. coli</i> + <i>C. jejuni</i> + <i>E. coli</i> * + Hadar (<i>C. coli</i> + <i>C. jejuni</i> + EPEC + Hadar)	3 1	4
<i>C. jejuni</i> + <i>E. coli</i> * + <i>Salmonella. sp</i>	4	4
<i>C. jejuni</i> + <i>E. coli</i> * + Enteritidis (<i>C. jejuni</i> + EPEC + Enteritidis)	1 1	2
Total	83	83

*non EPEC

Bacteria isolation and identification

Ten grams of skin from each chicken neck were used for bacterial isolation.

Campylobacter

Campylobacter were isolated according to the ISO 10272-1 [8] under microaerophilic conditions. Presumptive *Campylobacter* colonies were confirmed using API CAMPY strips (Biomerieux, Marcy-l'Etoile, France).

E. coli

E. coli were cultured following the AFNOR V08-053 standard [9] and confirmed with API 20E strips (Biomerieux, Marcy l'étoile, France). Serotyping was performed to detect EPEC (Enteropathogenic *E. coli*) strains with somatic O, flagellar H and surface A, B, L, and K anti-sera (Bio-Rad Laboratories, Marnes La Coquette, France).

Salmonella

Salmonella isolation was conducted based on the methods described in the ISO 6579 [10], confirmed with API 20E Strips (Biomerieux, Marcy l'Etoile, France), and serotyped with the somatic O and flagella H *Salmonella* anti-sera, according to the Kauffman-White scheme.

Results

A total of 150 chickens were collected and examined for the presence of *Campylobacter*, *E. coli*, and *Salmonella*. Among them, 135 (90%) were contaminated with *Campylobacter*, and 164 *Campylobacter* strains were isolated (68.9% were *C. coli* and 31.1% *C. jejuni*). Out of 84 chickens that were contaminated with *C. coli*, 29 were contaminated with *C. coli* and *C. jejuni*, while 22 chickens were contaminated with *C. jejuni* alone.

All the chicken carcasses were contaminated with *E. coli*. Seventeen (11.3%) of these strains agglutinated the antisera O 127 B8 (7), O126 B16 (4), O128 B12 (2), O142 K86 (2), O119 B14 (1), and O111 B4 (1). and were reported as enteropathogenic *E. coli* (EPEC).

A total of 103 *Salmonella* strains were isolated from 90 chickens (60%). Seventy-nine chickens were contaminated with only one serotype, ten with two different serotypes and one with four different serotypes. The serotypes obtained were Enteritidis, Hadar, Tilburg, Mikawasima, Bareilly, Cleveland, Colindale, Duesseldorf, Eko, Gwoza, Harburg, Hato, Hiduddify, Liverpool, Manhattan, Muenster, Reading, Saintpaul, and serotype II. The most frequent isolates were Enteritidis (45.6%) and Hadar (28.1%) [11].

Multiple contaminations with the bacteria genus listed above were obtained in 142 chickens (94.7%). The eight remaining chickens (5.3%) were contaminated only with *E. coli*. Fifty-two chickens (34.7%) had *Campylobacter* and *E. coli*, seven (4.7%) had *E. coli* and *Salmonella*, and 83 (55.3%) had *Campylobacter*, *E. coli* and *Salmonella*. Table 1 shows the different types of co-infections with these three bacteria together.

Discussion

Food-borne diseases represent a major public health problem worldwide. This study reveals a high prevalence of *Campylobacter*, *E. coli*, and *Salmonella* in chickens in Yaounde. These bacteria have already been implicated in outbreaks of food poisoning [4]. The high prevalence of *Campylobacter* (90%) in this study corroborates data from several other countries [12-13]. This may be explained by the fact that *Campylobacter* is commensal in many avian caeca [14]. Cross-transmission of *Campylobacter* is very high during poultry processing because the chickens are contaminated with their fecal materials [15]. In the present study, 68.9% of *Campylobacter* isolates were *C. coli*. Similar results have been obtained in Thailand [16] and South Africa [17]. In most of the cases, *C. jejuni* was the most frequent *Campylobacter* in poultry [18].

All the chickens purchased for this study were contaminated with *E. coli*. This may be due to the fact that *E. coli* is part of the normal enteric flora of chickens [5]. Among the 150 *E. coli* strains identified in this work, only 11.3% were enteropathogenic (EPEC); these strains usually cause diarrhoea in infants (younger than 2 years) through one or more virulence mechanisms [19]. The proportion of EPEC in this study (11.3%) was lower than that obtained by Farooq *et al.* in 2009 [20] in which all the *E. coli* isolated from 25 chickens were EPEC.

The prevalence of *Salmonella* in chickens in this study was 60%. This is nearly similar to what has been observed in some other developing countries: 68.2% in Ethiopia [21] and 72% in Thailand [22]. Enteritidis was the most predominant *Salmonella* serotype in this study as well as in many others [23-24]. Hadar, the second most important serotype (28.1%), has also been frequently isolated in chickens [25-26].

The multiple contaminations of many chickens by *Campylobacter*, *E. coli*, and *Salmonella* observed in this study are disturbing. These bacteria, which are dangerous for the young, the old, and the

immunocompromised, could be rapidly life-threatening. This study suggests the importance of monitoring poultry for enteric pathogens in poultry to avoid cross-contamination to humans. Efforts should be made to educate producers, retailers, and consumers on the proper handling and cooking of chicken meat.

Acknowledgments

This research was supported by the Institut Pasteur International network through ACIP (*Action Concertée Inter-Pasteurienne*) funding. The authors would like to thank Dr Anfumbon Kfutwah, who helped with English correction.

References

1. Wilson IG (2002) *Salmonella* and *Campylobacter* contamination of raw retail chickens from different producers: a six year survey. *Epidemiol Infect* 129: 635-645.
2. Todd EC (1997) Epidemiology of foodborne diseases: a worldwide review. *World Health Stat* 50: 30-50.
3. Meng J, Zhao S, Doyle MP (1998) Virulence genes of Shiga-toxin producing *Escherichia coli* isolated from food, animals and humans. *Int J Food Microbiol* 45: 229-235.
4. Sackey BA, Mensah P, Collison E, Sakyi-Dawson E (2000) *Campylobacter*, *Salmonella*, *Shigella* and *Escherichia coli* in live and dressed poultry from metropolitan Accra. *Int J Food Microbiol*. 71: 21-28.
5. Zhao C, Ge B, De Villena J, Sudler R, Yeh E, Zhao S, White DG, Wagner D, Meng J (2001) Prevalence of *Campylobacter* spp., *Escherichia coli*, and *Salmonella* Serovars in Retail Chicken, Turkey, Pork, and Beef from the Greater Washington, D.C., Area. *Appl Environ Microbiol* 67: 5431-5436.
6. Shane SM (2000) *Campylobacter* infection of commercial poultry. *Rev Sci Tech Off Int Epiz* 19: 376-395.
7. Griffin PM (1995) *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. In Blaser MJ, Smith PD, Ravdin JI, Greenberg HB, and Guerrant RL, editors. *Infections of gastrointestinal tract*. New York: Raven Press. 739-761.
8. ISO 10272-1 (2006) Microbiology of food and animal feeding stuffs. Horizontal method for detection and enumeration of *Campylobacter* spp. Part 1: Detection method. International Organization for Standardization.
9. AFNOR. (2002). NF V08-053 - Microbiology of food and animal feeding stuffs – Horizontal method for the enumeration of beta-glucuronidase positive *Escherichia coli* using 5-bromo-4-chloro-3-indolyl beta-D-glucuronide by colony count technique at 44°C -Routine method.
10. ISO 6579 (2002) Microbiology of food and animal feeding stuffs. Horizontal method for detection of *Salmonella* spp. 4th edition. International Organization for Standardization.
11. Wouafo M, Nzouankeu A, Kinack JA, Fonkoua M-C, Ejenguele G, Njine T, Ngandjio A. Prevalence and Antimicrobial Resistance of *Salmonella* Serotypes in Chickens from Retail Markets in Yaounde (Cameroon). *Microb Drug resist*. In press.
12. Ledergerber U, Regula G, Stephan R, Danuser J, Bissig B, Stärk KDC (2003) Risk factors for antibiotic resistance in *Campylobacter* spp. Isolated from raw poultry meat in Switzerland. *BMC Public Health* 3: 39-48.

13. Denis M, Refrégier-Petton J, Laisney M-J, Ermel G, Salvat G (2001) *Campylobacter* contamination in French chicken production from farm to consumers. Use of a PCR assay for detection and identification of *Campylobacter jejuni* and *Campylobacter coli*. *J Appl Microbiol* 91: 225-267.
14. Keener KM, Bashor MP, Curtis PA, Sheldon BW, Kathariou S (2004) Comprehensive Review of *Campylobacter* and Poultry Processing. *Compr Rev Food Sci Food Saf* 3: 105-116.
15. Pacheco K, Peña Y, Gamboa MM, Hernánd F (1999) Prevalence of *Campylobacter* in chickens in a tropical environment. *Rev Biomed* 10: 63-64.
16. Padungtod P, Kaneene JB (2005) *Campylobacter* in food animals and humans in northern Thailand. *J Food Prot* 68: 2519-2526.
17. van Nierop W, Dusé AG, Marais E, Aithma N, Thothobolo N, KasselM, Stewart R, Potgieter A, Fernandes B, Galpin JS, Bloomfield SF (2005) Contamination of chicken carcasses in Gauteng, South Africa, by *Salmonella*, *Listeria monocytogenes* and *Campylobacter*. *Int J Food Microbiol* 99: 1-6.
18. Suzuki H, Yamamoto S (2009) *Campylobacter* Contamination in Retail Poultry Meats and By-Products in the World: A Literature Survey. *J Vet Med Sci* 71: 255-261.
19. Levine MM, Edelman R (1984) Enteropathogenic *Escherichia coli* of classic serotypes associated with infant diarrhea: epidemiology and pathogenesis. *Epidemiol Rev* 6: 31-51.
20. Farooq S, Hussain I, Mir M.A, Bhat MA, Wani SA (2009) Isolation of atypical enteropathogenic *Escherichia coli* and Shiga toxin 1 and 2 producing *Escherichia coli* from avian species in India. *Lett Appl Microbiol* 48: 692-697.
21. Tibaijuka B, Molla B, Hildebrandt G, Kleer J (2003) Occurrence of *Salmonellae* in retail raw chicken products in Ethiopia. *Berl Munch Tierarztl Wochenschr* 116: 55-58.
22. Boonmar S, Bangtrakulnonth A, Pornrunangwong S, Marnrim N, Kaneko K, Ogawa M (1998) Predominant serovars of *Salmonella* in humans and foods from Thailand. *J Vet Med Sci* 60: 877-880.
23. Goncagül G, Günaydin E, Carli TK (2005) Prevalence of *Salmonella* serogroups in chicken meat. *Turk J Vet Anim Sci* 29: 103-106.
24. Al-Zenki S, Nasser A, Al-Safar A, Alomirah H, Al-Haddad A, Hendriksen RS, Aarestrup FM (2007) Prevalence and antibiotic resistance of *Salmonella* isolated from poultry farm and processing plant in the environment in the state of Kuwait. *Foodborne Pathog Dis* 4: 367-373.
25. Cardinale E, Gros-Claude JDP, Tall F, Cisse EF, Gueye EF, Salvat G (2003) Prevalence of *Salmonella* and *Campylobacter* in retail chicken carcasses in Senegal. *Rev Elev Med Vet Pays Trop* 56: 13-16.
26. Capita R, Alonso-Calleja C, Prieto M (2007) Prevalence of *Salmonella enterica* serovars and genovars from chicken carcasses in slaughterhouses in Spain. *J Appl Microbiol* 103: 1366-1375.

Corresponding author

Ndayo Wouafo Marguerite
 Pasteur Centre of Cameroon
 Laboratory of Hygiene and Environment – Microbiology
 PO Box 1274
 Yaounde, Cameroon
 Tel: +237 2223 1803 / +237 7747 7362
 Fax: +237 2223 1564
 Email: wouafo@pasteur-yaounde.org

Conflict of interests: No conflict of interests is declared.