

Novel astroviruses in insectivorous bats.

D. K. W. Chu, L. L. M. Poon, Y. Guan, J. S. M. Peiris

► **To cite this version:**

D. K. W. Chu, L. L. M. Poon, Y. Guan, J. S. M. Peiris. Novel astroviruses in insectivorous bats.. Journal of Virology, American Society for Microbiology, 2008, 82 (18), pp.9107-14. 10.1128/JVI.00857-08 . pasteur-00588955

HAL Id: pasteur-00588955

<https://hal-riip.archives-ouvertes.fr/pasteur-00588955>

Submitted on 27 Apr 2011

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Novel astroviruses in insectivorous bats**

2

3 DKW Chu¹, LLM Poon¹, Y Guan¹, JSM Peiris^{1,2}

4

5 ¹ State Key Laboratory of Emerging Infectious Diseases, Department of Microbiology,

6 The University of Hong Kong, Hong Kong SAR, China

7 ² HKU-Pasteur Research Centre, Hong Kong SAR, China.

8

9

10

11 Department of Microbiology, The University of Hong Kong

12 State Key Laboratory for Emerging Infectious Diseases

13 HKU-Pasteur Research Centre,

14 Hong Kong SAR

15

16 Key words: astroviruses, bats, zoonosis, coronaviruses, wild-life, evolution.

17 Running title: Novel astroviruses in insectivorous bats

18

19 Address correspondence to:

20 LLM Poon (llmpoon@hkucc.hku.hk), Tel (852)-2819-9943 or

21 JSM Peiris (malik@hkucc.hku.hk), Tel (852)-2855-4888

22 Department of Microbiology, The University of Hong Kong,

23 University Pathology Building, Queen Mary Hospital,

24 Pokfulam, Hong Kong SAR

25

26 **Abstract**

27 Bats are increasingly recognized to harbor a wide range of viruses and in most
28 instances these viruses appear to establish long term persistence in these animals.
29 They are the reservoir of a number of human zoonotic diseases including Nipah,
30 Ebola and SARS. We report the identification of novel groups of astroviruses in
31 apparently healthy insectivorous bats found in Hong Kong, in particular bats
32 belonging to the genera *Miniopterus* and *Myotis*. Astroviruses are important causes of
33 diarrhea in many animal species including humans. Many of the bat astroviruses
34 (BatAstV) form distinct phylogenetic clusters in the genus mammastrovirus within the
35 family *Astroviridae*. Virus detection rates of 36%-100% and 50%-70% were found in
36 *Miniopterus magnater* and *Miniopterus pusillus* bats respectively, captured within a
37 single bat-habitat during four consecutive visits spanning one year. There was high
38 genetic diversity of viruses in bats found within this single habitat. Some BatAstV
39 may be phylogenetically related to human astroviruses and further studies in a wider
40 range of bat species in different geographic locations are warranted. These findings
41 are likely to provide new insights into the ecology and evolution of astroviruses and
42 reinforce the role of bats as a reservoir of viruses with potential to pose a zoonotic
43 threat to human health.

44

45

46

47 **Introduction**

48 The Astroviridae are a family of non-enveloped, positive sense single-stranded RNA
49 viruses of approximately 28-30nm in size with a characteristic star-like surface
50 structure. The genomes of these viruses range in sizes from 6.4 - 7.3 kb and are
51 polyadenylated at their 3' ends (19). The viral RNA is synthesized by its RNA
52 dependent RNA polymerase (RdRp). The genome consists of 3 open reading frames
53 (ORFs) designated as ORF1a, ORF1b and ORF2 (Figure 1). ORF1a encodes
54 nonstructural polyprotein 1a while the entire ORF1 encodes polyprotein 1ab with a
55 ribosomal frame shift at the ORF1a / 1b junction (9). Efficiency of ORF1b translation
56 is estimated as only 25 - 28% of ORF1a (18). ORF2 encodes the viral structural
57 polyprotein which is required for virion formation. Studies with human astroviruses
58 showed that the structural polyprotein is intracellularly cleaved into functional units
59 including a capsid forming unit and a host binding motif unit (7, 12, 20).

60
61 Astroviruses have been identified from a variety of mammals (genus mamastrovirus)
62 and birds (genus avastrovirus) including humans (HAstV), bovine (BAstV), pigs
63 (PAstV), ovine (OAstV), mink (MAstV), dogs, cats, mice, chickens (CAstV) and
64 turkeys (TAstV). In most species, these viruses are associated with gastroenteritis but
65 some avian astroviruses have been associated with both intestinal and extraintestinal
66 manifestations [reviewed in (19)]. Human astroviruses appear to cause milder disease
67 than rotaviruses but are the second or third most common viral agent found in
68 children with diarrhea (5, 8). Astrovirus can also cause significant disease in the
69 elderly (16) and in immunocompromised patients (6).

70

71 Most of the surveillance studies of astroviruses focused on humans and domesticated

72 animals and relatively little is known about the prevalence of astroviruses in wildlife.
73 The role of bats as the reservoirs for zoonotic diseases including rabies, Hendra,
74 Nipah and Ebola has been highlighted in recent years [reviewed in ref. (2)].
75 Insectivorous bats have also been shown to harbor a range of novel coronaviruses
76 including the precursor of SARS coronavirus (15, 25). Some species of bats live in
77 close proximity to human habitation and thus it is important to have a better
78 understanding of the virus ecology found in bats. The range and diversity of
79 coronaviruses found in bats has led to the hypothesis that bat coronaviruses may be
80 the precursors of most other mammalian group 1 and 2 coronaviruses (30). These
81 findings highlight the importance of identifying novel viruses in wildlife in general
82 and bats in particular. We used random primers to detect novel viruses in bat faecal
83 specimens. Here, we report the discovery of novel astroviruses in bats in Hong Kong.
84 The remarkably high prevalence and genetic diversity of astroviruses in various bat
85 species found within a relatively small geographic area highlights the need for study
86 in other species of bats and in other geographic locations.

87

88 **Methods**

89

90 Sample collection

91 The sampling of bats was carried out in two phases and the sampling methods have
92 been described previously (4, 25). Phase 1 was carried out in 2004 and 2005 and
93 during this phase, swab samples were collected from different species of bats captured
94 in the wild in different habitats in Hong Kong. Phase 2 was carried out in 2005 and
95 2006 and swab samples were collected from bats of the genus *Miniopterus* captured
96 on four sampling occasions (June, August and December 2005 and March 2006) in a
97 single habitat, an abandoned mine cave in Lin Ma Hang, Hong Kong, near the border
98 with Mainland China (4). In both phases, species of bats captured for sampling were
99 healthy and identified by a bat taxonomist. Rectal and throat swabs, together with
100 fresh fecal samples if available, were collected. Swabs were placed in viral transport
101 medium (Earle's balanced salt solution, 0.2% sodium bicarbonate, 0.5% bovine serum
102 albumin, 200 µg of vancomycin per liter, 18 µg of amikacin per liter, 160 U of
103 nystatin per liter) in screw cap tubes and transported in a cool box to the laboratory
104 for processing. Bats were released at the site after sampling.

105

106 Viral nucleic acid extraction and RT-PCR

107 RNA from 140 µl of sample in transport medium was extracted by QIAamp virus
108 RNA mini kit (QIAGEN) following the protocol provided by the manufacturer.
109 Purified RNA was eluted in 60 µl of Elution buffer provided in the extraction kit.
110 cDNA was generated from RNA using Superscript III reverse transcriptase
111 (Invitrogen) in a 20 µl reaction containing 150 ng of random hexamers or 0.5 µM of a
112 gene specific reverse primer 5'-TTTGGTCCNCCNCTCCAAA-3' targeting the 3' end

113 of ORF1b, 10 mM of dithiothreitol, 0.5 mM of deoxynucleoside triphosphate mix, 1x
114 First-Strand buffer (Invitrogen), and 200 U of reverse transcriptase. Reaction mixtures
115 were incubated at 25°C for 5 min, followed by 50°C for 60 min and then the enzyme
116 was inactivated by heating at 70°C for 15 min.

117

118 Random hexamer-generated cDNA was screened for the presence of astrovirus using
119 hemi-nested PCR targeting RdRp gene. A 50 µl PCR reaction was set up containing 1
120 U Accuprime *Taq* DNA polymerase in 1x reaction buffer (Invitrogen), 2 µM each of
121 forward and reverse primers and 2 µl of cDNA or 1 µl of first PCR product as a
122 template. First-round PCR was carried out with a mix of two forward primers
123 5'-GARTTYGATTGGRCKCGKTAYGA-3' and
124 5'-GARTTYGATTGGRCKAGGTAYGA-3', and reverse primer

125 5'-GGYTTKACCCACATNCCRAA-3'. After an initial incubation at 94°C for 1 min,
126 30 cycles of amplification were carried out consisting of denaturation at 94°C for 30
127 sec, annealing at 50°C for 30 sec and extension at 68°C for 30 sec. Hemi-nested PCR
128 was carried out with a mix of two forward primers

129 5'-CGKTAYGATGGKACKATHCC-3' and 5'-AGGTAYGATGGKACKATHCC-3'

130 and the same reverse primers used in the first-round PCR using the same

131 thermocycling conditions as for the first-round PCR except that 40 cycles of

132 amplification were performed. Water controls were included in each run of the

133 RT-PCR assay. PCR products were analyzed by standard agarose gel electrophoresis.

134 The expected product size of the 2nd PCR is 422 base pairs. All positive results were

135 verified by direct DNA sequencing of the PCR amplicons.

136

137 Cloning and sequencing of PCR products

138 PCR products were purified by QIAquick PCR purification kit (QIAGEN) following
139 the manufacturer's instruction. Long PCR products (product sizes > 1000 bp) were
140 gel purified with QIAquick gel extraction kit (QIAGEN) and then cloned into
141 pCR2.1-TOPO plasmids (Invitrogen) for DNA sequencing. Multiple clones of a PCR
142 product were picked and sequenced by using BigDye Terminator v3.1 Cycle
143 Sequencing kit (Applied Biosystems). Sequencing products were analyzed by PRISM
144 3700 DNA analyzer (Applied Biosystems).

145
146 One astrovirus positive bat specimen, bat astrovirus AFCD337, was chosen for
147 sequencing of the viral genome directly from the original clinical specimen by using
148 cDNA generated by random hexamers and consensus primer RT-PCR. ORF2 region
149 and the 3' end of the virus sample were amplified using 3' Rapid Amplification of
150 cDNA Ends (RACE) System (Invitrogen) following the protocol provided by the
151 manufacturer (primers and PCR conditions available on request). ORF1a (partial) and
152 ORF1b sequences were assembled from multiple overlapping sequences derived from
153 PCR amplicons. Nine additional ORF1a (partial) / ORF1b astrovirus sequences from
154 other samples were obtained using similar method. The deduced sequence of these
155 samples has at least 3-fold sequence coverage.

156

157 Phylogenetic analysis

158 Sequence editing and sequence identities calculation were done using BioEdit version
159 7.0.4 (<http://www.mbio.ncsu.edu/BioEdit/bioedit.html>). Alignments of nucleotide
160 sequences and amino acid sequences were done using Clustal W (29) with default
161 parameters. Phylogenetic trees were constructed using Clustal X V2.0 (14) and Mega
162 4 (28) by the neighbour joining method with the nucleotide substitution model of

163 maximum composite likelihood and default parameters. Bootstrap values of the
164 phylogenetic tree constructed were generated by doing 1000 replicates.

165

166 Nucleotide sequence accession numbers.

167 The sequence of bat astrovirus AFCD337 reported in this paper were deposited in
168 GenBank under accession numbers xxxxx, and RdRp and the partial ORF1b
169 sequences of other strains of bat astroviruses were deposited under accession numbers
170 xxxxx-xxxxx. For the genetic analysis in this paper, other astrovirus genomes were
171 retrieved from GenBank including human astrovirus type 1 strain Oxford (L23513),
172 human astrovirus type 1 strain Dresden (AY720892), human astrovirus type 4 strain
173 Dresden (AY720891), human astrovirus type 4 isolate Goiania/GO/12/95/Brazil
174 (DQ070852), human astrovirus type 4 isolate Guangzhou (DQ344027), human
175 astrovirus type 5 isolate Goiania/GO/12/94/Brazil (DQ028633), human astrovirus
176 type 8 (AF260508), mink astrovirus (NC_004579), ovine astrovirus (NC_002469)
177 and turkey astrovirus type-1 (NC_002470).

178 **Results**

179

180 Detection of astroviruses in bats

181 A total of 262 bats were captured and rectal and throat swabs sampled in the two
182 phases sample collection in Hong Kong in 2004 to 2006. Bats sampled included nine
183 species, *Cynopterus sphinx*, *Hipposideros armiger*, *Miniopterus magnater*,
184 *Miniopterus pusillus*, *Miniopterus schreibersii*, *Myotis chinensis*, *Myotis ricketti*,
185 *Pipistrellus abramus* and *Rhinolophus rouxi*. A hundred and sixteen positives were
186 detected from 250 available rectal samples tested, representing a positive rate of 46%
187 (Table 1). On the other hand, only 19 (8%) positive throat swabs were found in 246
188 available samples tested. With the exception of two throat swabs collected from
189 *Miniopterus pusillus* and *Rhinolophus rouxi* in 2004, all the throat swabs positive
190 results came from bats with astrovirus detected in rectal samples. All the positive PCR
191 products were sequenced to confirm the identity of the amplicon and since the
192 sequences of the different PCR amplicons were largely non-identical, PCR
193 cross-contamination can be excluded.

194

195 Astroviruses were detected in 7 out of the 9 species of bats screened, i.e. *Miniopterus*
196 *magnater*, *Miniopterus pusillus*, *Miniopterus schreibersii*, *Myotis chinensis*, *Myotis*
197 *ricketti*, *Pipistrellus abramus* and *Rhinolophus rouxi* (Table 1). The detection rates of
198 astroviruses in these species of bats were remarkably high and ranged from 25% to
199 100%. However, astrovirus was not detectable in our samples collected from
200 *Cynopterus sphinx* (n=11) and *Hipposideros armiger* (n=10).

201

202 These same specimens had previously been tested for bat coronaviruses (3, 4, 25).

203 While 6% of bats were co-infected with both a bat astrovirus and a bat coronavirus,
204 such co-infection appears to be randomly distributed and there was no positive or
205 negative statistical association between the presence of these two viruses (Chi square
206 test with Yates correction, $p=0.82$).

207

208 Longitudinal study of astrovirus in bats at a single habitat

209 The bats listed in Table 1 include 157 *Miniopterus magnater* and *Miniopterus pusillus*
210 bats which were captured at four separate visits over a two year period at one habitat,
211 an abandoned mine cave in Hong Kong. *M. magnater* was found throughout the
212 period while *M. pusillus* was mainly found in two visits carried out in Dec 2005 and
213 Mar 2006 and only 1 found in a visit in Aug 2005. Sixty-two (54%) out of 115 rectal
214 swabs and 2 (2%) out of 116 throat swabs collected from *M. magnater* and 18 (55%)
215 out of 33 rectal swabs and 5 (15%) out of 33 throat swabs collected from *M. pusillus*
216 were positive for astrovirus. The overall positive rate in individual bats (either rectal
217 or throat swab or both positive) for *M. magnater* ranged from 36-100% at each of the
218 four visits which spanned the winter, spring and summer seasons and for *M. pusillus*
219 was 50% and 70% in the two instances when adequate bats were sampled.

220

221 Genetic and Phylogenetic analysis of a novel astrovirus from *Miniopterus pusillus*

222 A bat astrovirus (BatAstV) AFCD337 detected in a rectal specimen from a
223 *Miniopterus pusillus* bat collected in March 2006 was chosen as a representative virus
224 for more extensive genome sequencing. Approximately 74% of the genome of this
225 novel astrovirus was obtained by direct RT-PCR amplification from a rectal swab
226 sample. The partial 5067 nucleotide (excluding the poly-A tail) genome was
227 constructed by aligning sequences from multiple overlapping regions. Amino acid

228 sequences deduced from the viral genome include part of the open reading frame 1a
229 (ORF1a) and the complete ORF1b, ORF2 and 3' untranslated region followed by the
230 poly-A tail at the 3' end (Figure 1).

231

232 Amino acid sequences encoded by ORFs of the novel bat astrovirus were compared
233 with that of other astrovirus genomes including human astrovirus (HAstV) types 1, 4,
234 5 and 8 (17, 26) as well as mink astrovirus (MAstV) (22), ovine astrovirus (OAstV)
235 and turkey astrovirus type-1 (TAstV-1) (11) (Table 2). The findings show that the
236 identified bat astrovirus BatAstV AFCD337 is a novel mamastroviruses virus clearly
237 distinct from other known astroviruses. It has <53% and <27% genetic similarity to
238 other known astroviruses in the ORF1b and ORF2 regions, respectively.

239

240 The putative ORF1a (partial) and ORF1b of the bat astrovirus sequenced have sizes of
241 909 nt and 1572 nt respectively. A region at 5' end of ORF1a remains unsequenced so
242 far. ORF1a and ORF1b encodes for non-structural proteins which are essential for
243 virus replication. Characteristic features of BatAstV AFCD337 ORF1a and ORF1b
244 includes the protease motif in ORF1a; an astrovirus "slippery sequence" (AAAAAC)
245 at the junction between ORF1a and ORF1b which is required for inducing a
246 ribosomal shift event; a RdRp motif in ORF1b; and a conserved sequence at the end
247 of ORF1b of astroviruses (11). A conserved stem-loop structure that is predicted at the
248 3' end of the genomic RNA of human, ovine, porcine and turkey astroviruses type-1
249 was not found in BatAstV AFCD337 (10).

250

251 The putative ORF2 of the virus has a size of 2553 nt, which is the largest astrovirus
252 capsid gene known. The N-terminal half of the ORF2 protein, which was previously

253 shown to be more conserved among astroviruses and proposed to be the core
254 assembly domain of the viral capsid (12), was also found to be relatively conserved in
255 this bat astrovirus. The amino acid sequence similarities of this N-terminal half of the
256 bat astrovirus capsid protein to HAstV-1 Oxford, OAstV and MAstV are 36.3%,
257 45.0% and 39.5% respectively; compared with <27% similarity for the ORF2 region
258 overall (Table 2). Thus the C-terminal half of this bat astrovirus protein was highly
259 divergent when compared with other astroviruses. This observation supports the
260 speculation that the C-terminal half of the protein is located on the surface of the viral
261 particle and constitutes a region of the capsid that contributes to the species-specific
262 tropism of the virus (12).

264 Phylogenetic analysis on astrovirus in bats.

265 Phylogenetic analyses on ORF1a (partial) and ORF1b region and the ORF2 confirms
266 that BatAstV AFCD337 is a novel distinct astrovirus (Figure 2). To better define the
267 genetic diversity within the bat astroviruses by species, time and geographic location,
268 77 PCR RdRp amplicons (422 nucleotides) obtained from the rectal swab samples in
269 the screening PCR assay were selected for genetic sequence analysis. Bat astrovirus
270 gene sequences were aligned with that of other astroviruses including MAstV, OAstV,
271 HAstV-1 Oxford, HAstV-1 Dresden, HAstV-2, HAstV-4 Dresden, HAstV-4 Goiania,
272 HAstV-4 Guangzhou, HAstV-5 Goiania and HAstV-8. An avian astrovirus, Turkey
273 astrovirus was included as an out-group. A phylogenetic tree was constructed from the
274 sequence alignment (Figure 3). The 72 astroviruses detected in bats cluster together to
275 form a novel group of viruses within the cluster of mamastroviruses (Figure 3).
276 Within this group are found two subgroups of viruses, one which include the majority
277 of astroviruses detected from *Miniopterus magnater*, *Miniopterus pusillus* and

278 *Miniopterus schreibersii* (including BatAstV AFCD337) and another subgroup which
279 includes most of the viruses detected in *Myotis chinensis* and *Myotis ricketti*. Other
280 than these two major groups of bat astroviruses, a few astroviruses detected in
281 *Miniopterus magnater* and *Miniopterus pusillus* and a virus detected in a *Pipistrellus*
282 *abramus* appear to have an outgroup relationship to the others, albeit with weak levels
283 of statistical confidence in this phylogenetic topology (Figure 3). To further
284 investigate this, a 750 nt region of the RdRp and ORF1b (3' end) was sequenced of
285 these and other representative bat astroviruses. A phylogenetic tree based on the
286 aligned protein encoding sequences was shown (Figure 4). The phylogeny of the
287 major group of viruses related to BatAstV AFCD337 is confirmed. Interestingly
288 however, AFCD11 from *Pipistrellus abramus* and AFCD57 from *Miniopterus*
289 *magnater* appear phylogenetically related to the human astroviruses although with
290 modest bootstrap support. Their close relationship has been also confirmed by
291 phylogenetic analysis using an alternative method, MrBayes (data not shown).
292
293 Occasionally the same virus strain can be found in multiple bats sampled at the same
294 habitat in a single sampling trip (i.e. strains AFCD74 and AFCD79; strains AFCD175
295 and AFCD228). However, most viruses detected even at the same sampling occasion
296 at a single habitat are genetically diverse and no dominant strain could be discerned,
297 even though the detection rates of the virus was remarkably high.

298 **Discussion**

299 We report the discovery of novel astroviruses in 7 out of 9 species of apparently
300 healthy bats captured in Hong Kong. These astrovirus positive species include
301 *Miniopterus magnater*, *Miniopterus pusillus*, *Miniopterus schreibersii*, *Myotis*
302 *chinensis*, *Myotis ricketti*, *Pipistrellus abramus* and *Rhinolophus rouxi*. Attempts at
303 viral culture were so far unsuccessful (unpublished data). Phylogenetic analysis
304 revealed that 72 out of 77 BatAstV RdRp genes analyzed clustered together to form a
305 novel group of astroviruses. This virus group can be divided into two sub-groups,
306 subgroup A detected from *Miniopterus* species and subgroup B detected from *Myotis*
307 species. In the longitudinal study carried out in the abandoned mine cave habitat, the
308 subgroup A viruses appears to be circulating between *Miniopterus magnater* and
309 *Miniopterus pusillus* bats without any evidence of species restriction. This is in
310 marked contrast to the bat coronaviruses within the same habitat where Bat CoV 1A
311 and 1B appear to have a marked host restriction to *Miniopterus magnater* and
312 *Miniopterus pusillus*, respectively (4). Multiple clones of partial ORF1 sequence
313 (approximately 1000 nt) for 10 representative samples were analyzed and no evidence
314 of multiple infection was found although more systematic studies on this aspect are
315 needed.

316

317 The diversity of astroviruses in bats is remarkable. There is no significant
318 phylogenetic clustering of viruses found within a single sampling occasion. The
319 values of astrovirus RdRp amino acid pair-wise similarity found within a single bat
320 species, i.e. *Miniopterus magnater* captured in a single habitat (mine cave) ranged
321 between 51.1% to 100%, with 97.9% of these values lower than 90%. In contrast, the
322 pair-wise amino acid sequence similarities between the same gene region of human

323 astroviruses from geographically diverse regions were estimated to range between
324 92.6% to 99.1%. It has been previously reported that the amino acid sequence
325 identities of RdRp gene (covering 80% of the RdRp gene regions analysed in this
326 report) between 4 groups of avastrovirus, i.e. turkey astrovirus type 1-like viruses,
327 turkey astroviruses type 2-like viruses, avian nephritis virus-like viruses and
328 chicken-origin astroviruses detected in different regions were also highly diverse
329 ranging from 50.1% and 73.8% (23). The high virus detection rates of BatAstV in our
330 surveillance taken together with the marked genetic diversity of viruses from bats
331 within the same habitat are reminiscent of our observations with bat coronaviruses (4).
332 These findings may suggest that bats are persistently infected with astroviruses
333 although mark-recapture studies are needed to confirm this contention. Other
334 mammalian astrovirus infections tend to be short-lived in immunocompetent humans
335 or other animals. However, type 3 HAsV has been associated with persistent
336 gastroenteritis in immunocompetent children although the same virus serotype was
337 not repeatedly demonstrated over the full period of clinical diarrhea (1).
338
339 Five BatAstV sequences from *Miniopterus magnater*, *Miniopterus pusillus* and
340 *Pipistrellus abramus* failed to cluster with the subgroup A and B referred to above and
341 some of these (AFCD11 from *Pipistrellus abramus* and AFCD57 from *Miniopterus*
342 *magnater*) cluster rather with the human astroviruses although with weak statistical
343 support (Figure 4). Whether these bat viruses are related to the precursor of HAsV is
344 yet to be further investigated. Further sequence data from these strains may help
345 elucidate this phylogenetic association.
346
347 Evidence of recombination between astroviruses and also that between coronaviruses

348 is well documented (13, 24, 27). It has been reported that a stem-loop motif in 3'
349 untranslated region (UTR) was found conserved in mamastroviruses, TAstV-1, and in
350 avian infectious bronchitis virus which is a group 3 coronavirus (10). However, this
351 stem loop motif was not recognized in the BatAstV AFCD337. This 3' UTR
352 stem-loop motif is also absent in TAstV-2. However a phylogenetic analysis of 3'
353 UTRs did not indicate a close phylogenetic relationship between the two sequences of
354 BatAstV AFCD337 (81 nt) and TAstV-2 (accession no.: NC_005790) (196 nt) which
355 lack the 3'UTR stem-loop structure (data not shown). Recently a report on the
356 identification of a novel coronavirus from liver tissue of a whale with pulmonary
357 disease and terminal acute liver failure showed that the ORF6 of the novel
358 coronavirus possessed significant amino acid similarity to human astrovirus capsid
359 proteins (21). The high rates of infection of bats in the same mine cave habitat with
360 coronaviruses and astroviruses implies frequent co-infection with both viruses.
361 Therefore we searched for sequence of BatAstV AFCD337 similar to bat
362 coronaviruses using BLAST program with algorithms allowing a word-size down to
363 seven bases. However no sequences with significant similarity was detected between
364 bat astroviruses and coronaviruses co-circulating within the same species within the
365 same habitat.

366

367 The discovery of novel diverse astroviruses in bats and the genetic analysis of such
368 viruses is likely to provide new insights into the ecology and evolution of astroviruses
369 and reinforces the role of bats as a reservoir of viruses which sometimes pose a
370 zoonotic threat to human health. More extensive surveillance for astroviruses in bats
371 of different species and in different geographic areas is needed to further address these
372 questions.

373 **Acknowledgements**

374 This project is supported by National Institutes of Health (NIAID contract
375 HHSN266200700005C) and a Research Excellence Award to JSMP from The University of
376 Hong Kong. The study was approved and supported by the Department of Agriculture,
377 Fisheries and Conservation, Hong Kong, Special Administrative Region, People's Republic of
378 China. We thank K. S. Cheung, C. T. Shek and C. S. M. Chan of the Department of
379 Agriculture, Fisheries and Conservation, Hong Kong for facilitating this study.
380

ACCEPTED

381 **References**

- 382 1. **Caballero, S., S. Guix, W. M. El-Senousy, I. Calico, R. M. Pinto, and A.**
383 **Bosch.** 2003. Persistent gastroenteritis in children infected with astrovirus:
384 association with serotype-3 strains. *J Med Virol* **71**:245-50.
- 385 2. **Calisher, C. H., J. E. Childs, H. E. Field, K. V. Holmes, and T. Schountz.**
386 2006. Bats: important reservoir hosts of emerging viruses. *Clin Microbiol Rev*
387 **19**:531-45.
- 388 3. **Chu, D. K., J. S. Peiris, H. Chen, Y. Guan, and L. L. Poon.** 2008. Genomic
389 characterizations of bat coronaviruses (1A, 1B and HKU8) and evidence for
390 co-infections in *Miniopterus* bats. *J Gen Virol* **89**:1282-7.
- 391 4. **Chu, D. K., L. L. Poon, K. H. Chan, H. Chen, Y. Guan, K. Y. Yuen, and J.**
392 **S. Peiris.** 2006. Coronaviruses in bent-winged bats (*Miniopterus* spp.). *J Gen*
393 *Virol* **87**:2461-6.
- 394 5. **Dalton, R. M., E. R. Roman, A. A. Negredo, I. D. Wilhelmi, R. I. Glass,**
395 **and A. Sanchez-Fauquier.** 2002. Astrovirus acute gastroenteritis among
396 children in Madrid, Spain. *Pediatr Infect Dis J* **21**:1038-41.
- 397 6. **Gallimore, C. I., C. Taylor, A. R. Gennery, A. J. Cant, A. Galloway, D.**
398 **Lewis, and J. J. Gray.** 2005. Use of a heminested reverse transcriptase PCR
399 assay for detection of astrovirus in environmental swabs from an outbreak of
400 gastroenteritis in a pediatric primary immunodeficiency unit. *J Clin Microbiol*
401 **43**:3890-4.
- 402 7. **Geigenmuller, U., N. H. Ginzton, and S. M. Matsui.** 2002. Studies on
403 intracellular processing of the capsid protein of human astrovirus serotype 1 in
404 infected cells. *J Gen Virol* **83**:1691-5.
- 405 8. **Herrmann, J. E., D. N. Taylor, P. Echeverria, and N. R. Blacklow.** 1991.
406 Astroviruses as a cause of gastroenteritis in children. *The New England*
407 *Journal of Medicine* **324**:1757-1760.
- 408 9. **Jiang, B., S. S. Monroe, E. V. Koonin, S. E. Stine, and R. I. Glass.** 1993.
409 RNA sequence of astrovirus: distinctive genomic organization and a putative
410 retrovirus-like ribosomal frameshifting signal that directs the viral replicase
411 synthesis. *Proc Natl Acad Sci U S A* **90**:10539-43.
- 412 10. **Jonassen, C. M., T. O. Jonassen, and B. Grinde.** 1998. A common RNA
413 motif in the 3' end of the genomes of astroviruses, avian infectious bronchitis
414 virus and an equine rhinovirus. *J Gen Virol* **79 (Pt 4)**:715-8.
- 415 11. **Jonassen, C. M., T. T. Jonassen, T. M. Sveen, and B. Grinde.** 2003.
416 Complete genomic sequences of astroviruses from sheep and turkey:
417 comparison with related viruses. *Virus Res* **91**:195-201.
- 418 12. **Krishna, N. K.** 2005. Identification of structural domains involved in

- 419 astrovirus capsid biology. *Viral Immunol* **18**:17-26.
- 420 13. **Lai, M. M., and D. Cavanagh.** 1997. The molecular biology of coronaviruses.
421 *Adv Virus Res* **48**:1-100.
- 422 14. **Larkin, M. A., G. Blackshields, N. P. Brown, R. Chenna, P. A. McGettigan,**
423 **H. McWilliam, F. Valentin, I. M. Wallace, A. Wilm, R. Lopez, J. D.**
424 **Thompson, T. J. Gibson, and D. G. Higgins.** 2007. Clustal W and Clustal X
425 version 2.0. *Bioinformatics* **23**:2947-8.
- 426 15. **Lau, S. K., P. C. Woo, K. S. Li, Y. Huang, H. W. Tsoi, B. H. Wong, S. S.**
427 **Wong, S. Y. Leung, K. H. Chan, and K. Y. Yuen.** 2005. Severe acute
428 respiratory syndrome coronavirus-like virus in Chinese horseshoe bats. *Proc*
429 *Natl Acad Sci U S A* **102**:14040-5.
- 430 16. **Lewis, D. C., N. F. Lightfoot, W. D. Cubitt, and S. A. Wilson.** 1989.
431 Outbreaks of astrovirus type 1 and rotavirus gastroenteritis in a geriatric
432 in-patient population. *J Hosp Infect* **14**:9-14.
- 433 17. **Lewis, T. L., H. B. Greenberg, J. E. Herrmann, L. S. Smith, and S. M.**
434 **Matsui.** 1994. Analysis of astrovirus serotype 1 RNA, identification of the
435 viral RNA-dependent RNA polymerase motif, and expression of a viral
436 structural protein. *J Virol* **68**:77-83.
- 437 18. **Lewis, T. L., and S. M. Matsui.** 1996. Astrovirus ribosomal frameshifting in
438 an infection-transfection transient expression system. *J Virol* **70**:2869-75.
- 439 19. **Mendez, E., and C. F. Arias.** 2007. Astroviruses, p.981-1000. D. M. Knipe, P.
440 M. Howley, D. E. Griffin, et al. *Fields Virology*, 5th ed., vol. 1. Lippincott
441 Williams & Wilkins, Philadelphia, Pa.
- 442 20. **Mendez, E., T. Fernandez-Luna, S. Lopez, M. Mendez-Toss, and C. F.**
443 **Arias.** 2002. Proteolytic processing of a serotype 8 human astrovirus ORF2
444 polyprotein. *J Virol* **76**:7996-8002.
- 445 21. **Mihindukulasuriya, K. A., G. Wu, J. St Leger, R. W. Nordhausen, and D.**
446 **Wang.** 2008. Identification of a Novel Coronavirus from a Beluga Whale
447 Using a Pan-viral Microarray. *J Virol*.
- 448 22. **Mittelholzer, C., K. O. Hedlund, L. Englund, H. H. Dietz, and L. Svensson.**
449 2003. Molecular characterization of a novel astrovirus associated with disease
450 in mink. *J Gen Virol* **84**:3087-94.
- 451 23. **Pantin-Jackwood, M. J., E. Spackman, and P. R. Woolcock.** 2006.
452 Molecular characterization and typing of chicken and turkey astroviruses
453 circulating in the United States: implications for diagnostics. *Avian Dis*
454 **50**:397-404.
- 455 24. **Pantin-Jackwood, M. J., E. Spackman, and P. R. Woolcock.** 2006.
456 Phylogenetic analysis of Turkey astroviruses reveals evidence of

- 457 recombination. *Virus Genes* **32**:187-92.
- 458 25. **Poon, L. L., D. K. Chu, K. H. Chan, O. K. Wong, T. M. Ellis, Y. H. Leung,**
459 **S. K. Lau, P. C. Woo, K. Y. Suen, K. Y. Yuen, Y. Guan, and J. S. Peiris.**
460 2005. Identification of a novel coronavirus in bats. *J Virol* **79**:2001-9.
- 461 26. **Silva, P. A., D. D. Cardoso, and E. Schreier.** 2006. Molecular
462 characterization of human astroviruses isolated in Brazil, including the
463 complete sequences of astrovirus genotypes 4 and 5. *Arch Virol* **151**:1405-17.
- 464 27. **Strain, E., L. A. Kelley, S. Schultz-Cherry, S. V. Muse, and M. D. Koci.**
465 2008. Genomic Analysis of Closely Related Astroviruses. *J Virol*.
- 466 28. **Tamura, K., J. Dudley, M. Nei, and S. Kumar.** 2007. MEGA4: Molecular
467 Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*
468 **24**:1596-9.
- 469 29. **Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G.**
470 **Higgins.** 1997. The CLUSTAL_X windows interface: flexible strategies for
471 multiple sequence alignment aided by quality analysis tools. *Nucleic Acids*
472 *Res* **25**:4876-82.
- 473 30. **Vijaykrishna, D., G. J. Smith, J. X. Zhang, J. S. Peiris, H. Chen, and Y.**
474 **Guan.** 2007. Evolutionary insights into the ecology of coronaviruses. *J Virol*
475 **81**:4012-20.
- 476
- 477

478 **Figure Legends**

479

480 Figure 1. Schematic diagrams of Mink astrovirus genome and BatAstV AFCD337
481 partial genome. ORFs, protease motif (Pro) and RdRp motif (RdRp) are shown in the
482 diagram. The unsequenced putative 5' end genome region of BatAstV AFCD337 is
483 represented by a dotted line. The 2.5 kb ORF1a (partial), ORF1b region, and ORF2
484 regions used for phylogenetic analysis in Figure 2 are indicated by arrows.

485

486 Figure 2. Phylogenetic analysis of partial ORF1a (~800 nt), ORF1b and ORF2
487 nucleotide sequences comparing BatAstV AFCD337 with astroviruses of other
488 species. Nine other BatAstV sequences are included in the ORF1a (partial) and
489 ORF1b phylogenetic trees. Alignment was based on the encoded amino acid
490 sequences. Abbreviations used for different astroviruses are bat (BatAstV), mink
491 (MAstV), ovine (OAstV), human (HAstV) and turkey type-1 (TAstV-1).

492

493 Figure 3. Phylogenetic tree constructed with RdRp gene sequences (422 nt) amplified
494 by the RT-PCR screening assay. Sequences of 77 bat astroviruses, and other respective
495 sequences of different astroviruses isolated from human (HAstV), mink (MAstV),
496 ovine (OAstV) and turkey type-1 (TAstV-1) were included and aligned based on the
497 nucleotide sequences. For the bat specimens collected during the phase 2 longitudinal
498 study, the sampling dates are indicated in the tree as follows: (1) - June 2005; (2) -
499 August 2005; (3) - December 2005; (4) - March 2006.

500

501 Figure 4. Phylogenetic tree constructed with 750 nt sequences of RdRp gene and
502 ORF1b (3' end) of representative astroviruses isolated from bat (BatAstV) human

503 (HAsV), mink (MAstV), ovine (OAsV) and turkey type-1 (TAsV-1). These
504 sequences were aligned based on the encoded amino acid sequences and reverse
505 translated back to nucleotides for the phylogenetic analysis.

ACCEPTED

506 Table 1. Detection of astrovirus in bats by RT-PCR

	Rectal samples			Throat samples			Bats		
	No. tested	No. positive	Positive %	No. tested	No. positive	Positive %	No. tested	No. positive	Positive %
<u>Insectivorous bats</u>									
<i>Hipposideros armiger</i>	10	0	0%	10	0	0%	10	0	0%
<i>Miniopterus magnater</i>	122	67	55%	123	4	3%	132	67	51%
<i>Miniopterus pusillus</i>	73	31	42%	71	6	8%	74	32	43%
<i>Miniopterus schreibersii</i>	3	3	100%	2	1	50%	3	3	100%
<i>Myotis chinensis</i>	9	3	33%	9	1	11%	9	3	33%
<i>Myotis ricketti</i>	12	10	83%	12	5	42%	12	10	83%
<i>Pipistrellus abramus</i>	2	1	50%	1	0	0%	3	1	33%
<i>Rhinolophus rouxi</i>	8	1	13%	8	2	25%	8	2	25%
<u>Fruit bats</u>									
<i>Cynopterus sphinx</i>	11	0	0%	10	0	0%	11	0	0%
<u>Total</u>	250	116	46%	246	19	8%	262	118	45%

507 Table 2. Amino acid sequence similarities between prototype BatAstV AFCD337, subgroup A (BatAstV AFCD68), subgroup B (BatAstV
 508 WCF140) and other astroviruses.

Strains	BatAstV AFCD337		
	ORF1a (partial)	ORF1b	ORF2
BatAstV AFCD68	79.9%	74.2%	-
BatAstV WCF140	54.0%	68.5%	-
HAstV-1 Dresden	17.2%	48.4%	22.3%
HAstV-1 Oxford	17.9%	49.2%	22.1%
HAstV-4 Guangzhou	17.5%	49.0%	22.3%
HAstV-4 Goiania/95/Brazil	17.9%	49.2%	22.2%
HAstV-4 Dresden	17.9%	48.8%	21.8%
HAstV-5 Goiania/94/Brazil	17.5%	49.2%	20.9%
HAstV-8	17.2%	48.8%	20.5%
MAstV	31.2%	52.7%	24.1%
OAstV	28.9%	52.6%	26.2%
TAstV-1	13.9%	40.2%	14.1%

509

Figure 1




Figure 2







Figure 3



Subgroup A

Subgroup B

0.05

Figure 4

