



## **Molecular epidemiology of Crimean- Congo hemorrhagic fever virus genome isolated from ticks of Hamadan province of Iran.**

F. Tahmasebi, S.M. Ghiasi, E. Mostafavi, M. Moradi, N. Piazak, A. Mozafari, A. Haeri, A.R. Fooks, Sadegh Chinikar

### **► To cite this version:**

F. Tahmasebi, S.M. Ghiasi, E. Mostafavi, M. Moradi, N. Piazak, et al.. Molecular epidemiology of Crimean- Congo hemorrhagic fever virus genome isolated from ticks of Hamadan province of Iran.. Journal of Vector Borne Diseases, 2010, 47 (4), pp.211-6. pasteur-00748082

**HAL Id: pasteur-00748082**

**<https://riip.hal.science/pasteur-00748082>**

Submitted on 4 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Molecular epidemiology of Crimean-Congo hemorrhagic fever virus genome isolated from ticks of Hamadan province of Iran

F. Tahmasebi<sup>a</sup>, S.M. Ghiasi<sup>b</sup>, E. Mostafavi<sup>b</sup>, M. Moradi<sup>b</sup>, N. Piazak<sup>c</sup>, A. Mozafari<sup>d</sup>, A. Haeri<sup>e</sup>, A.R. Fooks<sup>f</sup> & S. Chinikar<sup>b</sup>

<sup>a</sup>Islamic Azad University, Research and Science Branch, Tehran; <sup>b</sup>Laboratory of Arboviruses and Viral Haemorrhagic Fevers (National Reference Laboratory), Pasteur Institute of Iran, <sup>c</sup>Entomology Laboratory, Parasitology Department, Pasteur Institute of Iran, Tehran; <sup>d</sup>Iran University of Medical Science, Tehran; <sup>e</sup>Medical School, Shahid Beheshti Medical University, Tehran, Iran; <sup>f</sup>Veterinary Laboratory Agency, Weybridge, New Haw, Addlestone, U.K.

### Abstract

**Background & objectives:** Crimean-Congo hemorrhagic fever (CCHF) virus is a tick-borne member of the genus *Nairovirus*, family *Bunyaviridae*. CCHFV has been isolated from at least 31 different tick species. The virus is transmitted through the bite of an infected tick, or by direct contact with CCHFV-infected patients or the products of infected livestock. This study was undertaken to study the genetic relationship and distribution of CCHFV in the tick population of Hamadan province of Iran.

**Method:** In this study, RT-PCR has been used for detection of the CCHFV genome.

**Results:** This genome was detected in 19.2% of the ticks collected from livestock of different regions of the Hamadan province in western Iran. The infected species belonged to *Hyalomma detritum*, *H. anatolicum*, *Rhipicephalus sanguineus* and *Argas reflexus*. With one exception, genetic analysis of the virus genome isolates showed high sequence identity to each other. Even though they clustered in the same group with the strain circulating in Iran, they had a closer relationship to the Matin strain.

**Interpretation & conclusion:** Vector control programs should be applied for reducing population density of potential tick vectors in this province. Further surveys are indicated in this region to provide a better view of the distribution and epidemiology of the virus.

**Key words** *Argas reflexus*; CCHF; RT-PCR

### Introduction

Crimean-Congo hemorrhagic fever virus (CCHFV) is a tick-borne virus with a negative-sense, single-stranded (ss) RNA genome containing S (small), M (medium), and L (large) segments. It belongs to the genus *Nairovirus* and the family *Bunyaviridae*. The virus causes a severe hemorrhagic syndrome in humans, with a fatality rate of up to 50%, while in animals, the virus causes subclinical infection<sup>1–6</sup>. The CCHFV is transmitted to humans through the bite of

*Ixodid* ticks (mostly *Hyalomma* genus) or by contact with blood or tissues from infected livestock. Moreover, it has been demonstrated that most statistics about transmission routes in Iran relate to people in close contact with tissues and blood from CCHF-affected livestock<sup>7–15</sup>. In addition to zoonotic transmission, CCHFV can spread from person to person and is one of the rare hemorrhagic fever viruses able to cause nosocomial outbreak in hospitals<sup>16–17</sup>. The virus is distributed across Africa, Eastern Europe, the Middle East, and Asia. The dis-

tribution of CCHFV coincides precisely with the distribution of *Hyalomma* ticks<sup>3,5,16, 18,19</sup>.

Although sporadic surveys of CCHF in livestock and humans have been undertaken since 1970, it wasn't until the 1999 outbreak that CCHF was recognized as one of the country's major public health concerns. Since 2000, CCHF infection has been demonstrated in 23 out of the 30 provinces of Iran; with Sistan-v-Baluchistan, Isfahan, Fars, and Khuzestan being the most heavily infected provinces. The first confirmed case of human CCHF in the Hamadan province in western Iran, was reported in Hamadan city in 2004. Recent surveys have shown circulation of the virus in livestock existing in the province<sup>16,20</sup>, but no further data on the genetic variability of the CCHFV strain from the Hamadan province have been available. However, several partial and complete S-genome segment sequences have become available through GenBank (Accession Nos. AY366373–AY366379 for partial sequences and DQ446212–DQ446215 for complete sequences). This study describes the genetic relationship and distribution of CCHFV in the tick populations from Hamadan province.

## Material & Methods

**Study area:** Hamadan province is one of the highest and coldest regions in Iran. This district is located between 45° 32' and 48° E and 34° 47' and 35° 1' N and consists of eight townships. Hamadan is among the major sheep and goat raising provinces in Iran. The economy is mainly based on agriculture and livestock husbandry, and so a large proportion of the population comes into close contact with livestock. This provides a high risk of exposure to CCHFV. In this study the province was divided into three regions, with two townships selected randomly from each region. Therefore, six townships were studied—Razan and Kabotar-Ahang in the north; Hamadan and Bahar in the centre; and Malayer and Toyserkan in the south of the province (Fig. 1).

**Sample collection and preparation:** In this study, carried out during 2007, 70 villages were randomly selected as the study area. In each village, 20–30 sheep were checked for tick collection. The entire body of each sheep, but mainly: the ears, nape of the neck, perineum, scrotum and the tail base, were inspected for the presence of ticks. Collected ticks were




Fig. 1: Locations of collecting the ticks to characterize the situation of CCHFV infection in Hamadan province, the sampled townships are shown with stars

kept alive in separate labeled vials and sent to the Pasteur Institute of Iran, for species identification and molecular detection of CCHFV.

**RT-PCR:** Ticks were individually washed twice by PBS (PBS, pH 7.4) and crushed with a mortar and pestle in 200–300  $\mu$ l of PBS. Total RNA was extracted using an RNeasy mini kit (QIAGEN, Cat No. 2215716) according to the manufacturer's instructions. The extracted total RNA was stored at  $-70^{\circ}\text{C}$  until use. For the RT-PCR, a master mix was prepared as follows: 28  $\mu$ l of RNase free water, 10  $\mu$ l of buffer ( $5 \times \text{conc.}$ ), 2  $\mu$ l of dNTP mixture, 2  $\mu$ l of enzyme mixture containing reverse transcriptase and Taq DNA polymerase enzymes, 1  $\mu$ l of primer F2 ( $5' \mu\text{-TGGACACCTTCACAACTC-3}\mu$ ), 1  $\mu$ l of Primer R3 ( $5' \mu\text{-GACAATTCCTACACC-3}\mu$ ) and 1  $\mu$ l of RNase inhibitor. The F2 and R3 primers amplify a 536 bp fragment inside the S-segment of the CCHFV genome. For qualitative analysis in real-time RT-PCR, 1  $\mu$ l of SYBR Green dye with 1/10000 dilution was also used in the master mix. The thermal cycling program for the RT-PCR, for both techniques, included 30 min at  $50^{\circ}\text{C}$  for reverse transcription reaction (cDNA synthesis); followed by 35 cycles of  $95^{\circ}\text{C}$  for 30 sec,  $50^{\circ}\text{C}$  for 30 sec,  $72^{\circ}\text{C}$  for 45 sec, and a final extension at  $72^{\circ}\text{C}/5$  min. For gel-based RT-PCR product analysis, 5  $\mu$ l of the PCR products was mixed with 1  $\mu$ l loading buffer ( $6 \times \text{conc.}$ ) and then analyzed by gel electrophoresis. In real-time analysis, both amplification and dissociation curves were evaluated with respect to negative and positive controls, as well as in the gel-based analysis<sup>1,8,16,21</sup>.

**Sequencing and genetic analysis:** PCR products were purified using quick-spin PCR purification kit (Intron) and directly underwent sequencing by specific primers. All sequenced amplicons were placed in the GenBank with Accession Nos. GU456723 to GU456728 related to CT9, CT10, CT12, CT13, CT14 and CT15, respectively. For phylogenetic analysis, a 499 bp long fragment of the S-segment was used. The sequences of CCHF viruses were analyzed by a neighbor-joining method with Kimura two




Fig. 2: The proportion of tick species infected with CCHFV

parameter distances by using Mega 4 software. Bootstrap confidence limits were based on 500 replicates. Evolutionary divergence, distance matrix and subsequently sets of phylogenetic trees were calculated by the software<sup>22</sup>.

## Results

A total of 328 adult ticks (199 males and 129 females) were collected from the north, center and south of Hamadan province. These ticks were grouped into five species, including: *H. detritum* (89.6%) (Found at all locations), *Hyalomma anatolicum* (3.4%), and *Rhipicephalus sanguineus* (6.1%) [both species found in central and southern regions, *Argas reflexus* (0.4%) collected from northern parts], and *Ornithodoros lahorensis* (0.3%). The two latter species, *A. reflexus* and *O. lahorensis*, showed narrow distributions (Fig. 2).

The presence of CCHFV was confirmed by RT-PCR technique in which a 536 bp amplicon is amplified inside the S segment of the CCHFV genome isolates from 63 collected ticks. Among positive ones, 42.9 and 57.1% were males and females, respectively. The infected tick species belonged to *H. detritum* (16.32%), *H. anatolicum* (18.18%), *R. sanguineus* (55%) and *A. reflexus* (100%). *Ornithodoros lahorensis* was RT-PCR negative.

The genome isolates belonging to different parts of the study area were named as: CT9 (isolates of *R.*




Fig. 3: Phylogenetic tree (neighbour-joining method) calculated for CCHFV S-segment sequences. New sequences in this paper (CT9, CT10, CT12, CT13, CT14 and CT15) and previously described sequences from Iran (AY 366373–8, DQ446212–14) with exceeding of 70% are shown.

*sanguineus*); CT10, CT12 and CT15 (all isolates of *H. detritum*); CT13 (isolated from *A. reflexus*); and CT14 (isolated from *H. anatolicum*). By using various methods for reconstructing phylogeny, a CCHFV phylogenetic tree was obtained based on individual sequences of the 536 bp fragment of the nucleoprotein gene. The genetic analysis revealed some nucleotide substitutions. Notably, the isolates (with the exception of CT9) were all clustered with Matin strain (Pakistani strain) and showed a closer relationship to this strain as the previously published Iranian isolates of CCHFV genome had been clustered in one group close to the Matin strain. The CT9 isolate showed lower identity (89.7%) and more divergence (6.8) compared to the other isolates (Fig. 3).

## Discussion

The data obtained provide evidence for the presence of CCHFV in different regions of Hamadan province. CCHFV infection is shown in *H. detritum*, *H. anatolicum*, *R. sanguineus* and *A. reflexus*. These findings are in concordance with previous studies in other parts of the country. *Hyalomma* ticks are the primary vectors for the widespread transmission of CCHFV throughout Europe, Asia, the Middle East, and Africa<sup>3,5,19</sup>. Although in the epidemiology of CCHF virus, *Hyalomma* ticks are considered to be the most important vector and reservoir, the virus has also been reported from other genera of ticks. In 1979, CCHFV was first isolated in Iran from *O. lahorensis*<sup>23</sup>. In a study done in Ardabil province, north-west of Hamadan, 33.3% of the ticks, including *Hyalomma* sp, *Rhipicephalus* sp and *Ornithodoros* sp, were infected with CCHFV. In another study in Bahar (central part of Hamadan Province) 11.3% of ticks, including: *H. marginatum*, *H. dromedarii*, *H. anatolicum*, *R. bursa*, *R. sanguineus* and *H. punctata*, were CCHFV positive<sup>24</sup>. Current research shows CCHF infection in 19.3% of ticks collected from Hamadan province. Therefore, people in this province, involved with livestock, should protect themselves from direct contact with CCHFV contaminated tissue and blood.

In the present study, CCHFV contamination was confirmed in *A. reflexus*; an argasid (soft) tick which mainly infests pigeons, but occasionally infests mammals. This suggests that more consideration should be given to the detection of the CCHF virus genome in soft ticks. Soft ticks have a long life-span and can preserve the CCHF virus for long periods, potentially providing a persistent virus reservoir, for infection of the corresponding livestock and hard ticks in a region, and consequently, possible outbreaks of CCHF.

This report presents data on the CCHF situation in ticks, and the genetic variability of the CCHFV isolates obtained from different parts of Hamadan province. The relatively high divergence of the CT9 iso-



late, compared to the other isolates, may correlate with high geographical distance. CT9 was isolated from the southern part of the province, whereas the other isolates were mostly found in the central and northern parts. The remarkable genetic variability of this isolate could also be attributed to amplification of the virus and recombination events which may have occurred in the species of host tick from which it was isolated: *R. sanguineus*. This uncertainty shows that further molecular epidemiology surveys are required in this region to identify the exact phylogenetic relationship between the circulating strains<sup>6,25</sup>. Comparison of the data produced from three regions in the province suggested that ticks in the northern parts are infected more than the central and southern regions. Persons with high risk occupations such as shepherds, farmers, veterinarians, and other people who are in close contact with ticks should take precautions to avoid exposure to virus-infected ticks or virus-contaminated animal blood or tissues. Applying commercially available insect repellents, such as diethyl toluimidean DEET, to exposed skin, and the use of clothing impregnated with permethrin, can give some protection against tick bites. This study indicates that CCHF should be regarded as a serious health problem, worthy of consideration by the health centers in this province, and neighbouring regions, in order to develop and implement strategies to decrease the tick population, and alert high risk professionals. Further surveys on human and animal populations in these regions should be recommended to provide a better view of the distribution and epidemiology of the virus in this province.

### Acknowledgement

We would like to thank the other members of the National Reference Laboratory for Arboviruses and Viral Haemorrhagic Fevers at Pasteur Institute of Iran for their technical support. This study was funded by the budget of the National Reference Laboratory for Arboviruses and Viral Haemorrhagic Fevers at Pasteur Institute of Iran. Authors declare no conflict of interest.

### References

1. Chinikar S, Persson SM, Johansson M, Bladh L, Goya M, Houshmand B, *et al.* Genetic analysis of Crimean-Congo hemorrhagic fever virus in Iran. *J Med Virol* 2004; 73(3): 404–11.
2. Dickson DL, Turell MJ. Replication and tissue tropisms of Crimean-Congo hemorrhagic fever virus in experimentally infected adult *Hyalomma truncatum* (Acari: Ixodidae). *J Med Entomol* 1992; 29(5): 767–73.
3. Ergonul O. Crimean-Congo haemorrhagic fever. *Lancet Infect Dis* 2006; 6(4): 203–14.
4. Honig JE, Osborne JC, Nichol ST. The high genetic variation of viruses of the genus *Nairovirus* reflects the diversity of their predominant tick hosts. *Virology* 2004; 318(1): 10–6.
5. Whitehouse CA. Crimean-Congo hemorrhagic fever. *Antiviral Res* 2004; 64(3): 145–60.
6. Hewson R, Chamberlain J, Mioulet V, Lloyd G, Jamil B, Hasan R, *et al.* Crimean-Congo haemorrhagic fever virus: sequence analysis of the small RNA segments from a collection of viruses worldwide. *Virus Res* 2004; 102(2): 185–9.
7. Dohm DJ, Logan TM, Linthicum KJ, Rossi CA, Turell MJ. Transmission of Crimean-Congo hemorrhagic fever virus by *Hyalomma impeltatum* (Acari: Ixodidae) after experimental infection. *J Med Entomol* 1996; 33(5): 848–51.
8. Durden LA, Logan TM, Wilson ML, Linthicum KJ. Experimental vector incompetence of a soft tick, *Ornithodoros sonrai* (Acari: Argasidae), for Crimean-Congo hemorrhagic fever virus. *J Med Entomol* 1993; 30(2): 493–6.
9. Gonzalez JP, Camicas JL, Cornet JP, Faye O, Wilson ML. Sexual and transovarian transmission of Crimean-Congo haemorrhagic fever virus in *Hyalomma truncatum* ticks. *Res Virol* 1992; 143(1): 23–8.
10. Gordon SW, Linthicum KJ, Moulton JR. Transmission of Crimean-Congo hemorrhagic fever virus in two species of *Hyalomma* ticks from infected adults to cofeeding immature forms. *Am J Trop Med Hyg* 1993; 48(4): 576–80.
11. Hassanein KM, El-Azazy OM. Isolation of Crimean-Congo hemorrhagic fever virus from ticks on imported Sudanese sheep in Saudi Arabia. *Ann Saudi Med* 2000; 20(2): 153–4.
12. Logan TM, Linthicum KJ, Bailey CL, Watts DM, Dohm DJ, Moulton JR. Replication of Crimean-Congo hemorrhagic fever virus in four species of ixodid ticks (Acari)

- infected experimentally. *J Med Entomol* 1990; 27(4): 537–42.
13. Logan TM, Linthicum KJ, Bailey CL, Watts DM, Moulton JR. Experimental transmission of Crimean-Congo hemorrhagic fever virus by *Hyalomma truncatum* Koch. *Am J Trop Med Hyg* 1989; 40(2): 207–12.
  14. Shepherd AJ, Swanepoel R, Cornel AJ, Mathee O. Experimental studies on the replication and transmission of Crimean-Congo hemorrhagic fever virus in some African tick species. *Am J Trop Med Hyg* 1989; 40(3): 326–31.
  15. Zeller HG, Cornet JP, Camicas JL. Experimental transmission of Crimean-Congo hemorrhagic fever virus by west African wild ground-feeding birds to *Hyalomma marginatum* rufipes ticks. *Am J Trop Med Hyg* 1994; 50(6):676–81.
  16. Chinikar S, Ghiasi SM, Hewson R, Moradi M, Haeri A. Crimean-Congo hemorrhagic fever in Iran and neighbouring countries. *J Clin Virol* 2010; 47: 110–4.
  17. Joubert JR, King JB, Rossouw DJ, Cooper R. A nosocomial outbreak of Crimean-Congo haemorrhagic fever at Tygerberg Hospital. Pt III. Clinical pathology and pathogenesis. *S Afr Med J* 1985; 68(10): 722–8.
  18. Papa A, Velo E, Papadimitriou E, Cahani G, Kota M, Bino S. Ecology of the Crimean-Congo hemorrhagic fever endemic area in Albania. *Vector Borne Zoonotic Dis* 2009; 9(6): 713–6.
  19. Swanepoel R, Shepherd AJ, Leman PA, Shepherd SP, McGillivray GM, Erasmus MJ, *et al.* Epidemiologic and clinical features of Crimean-Congo hemorrhagic fever in southern Africa. *Am J Trop Med Hyg* 1987; 36(1): 120–32.
  20. Chinikar S, Goya MM, Shirzadi MR, Ghiasi SM, Mirahmadi R, Haeri A, *et al.* Surveillance and laboratory detection system of Crimean-Congo haemorrhagic fever in Iran. *Transbound Emerg Dis* 2008; 55(5–6): 200–4.
  21. Duh D, Nichol ST, Khristova ML, Saksida A, Hafner-Bratkovic I, Petrovec M, *et al.* The complete genome sequence of a Crimean-Congo hemorrhagic fever virus isolated from an endemic region in Kosovo. *Virol J* 2008; 5: 7.
  22. Tamura KDJ, Nei M, Kumar S. MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0 *Mol Biol Evol* 2007;1093/molbev/msm092.
  23. Sureau PKJ, Casals J, Digoutte JOP, Salaun JJ, Piazak N, Calvo MA. Isolation of Thogoto, Wad medani, Wanowrie, and Crimean-Congo hemorrhagic fever viruses from ticks of domestic animals in Iran. *Ann Virol (Inst Pasteur)* 1980; 131E: 185–200.
  24. Moradi AR CS, Oshaghi MA, Vatandoost H, Houlakoui Naeini K, Zahirnia AH, Telmadarraiy Z. Molecular detection of Crimean-Congo hemorrhagic fever (CCHF) virus in Ticks (Ixodidae, Argasidae) of Hamadan province, Iran. *Biochem Cell Arch* 2008; 8(1): 119–23.
  25. Hewson R, Gmyl A, Gmyl L, Smirnova SE, Karganova G, Jamil B, *et al.* Evidence of segment reassortment in Crimean-Congo haemorrhagic fever virus. *J Gen Virol* 2004; 85(Pt 10): 3059–70.

*Corresponding author:* Dr Sadegh Chinikar, Head of Arboviruses and Viral Hemorrhagic Fevers Laboratory (National Reference Laboratory), Pasteur Institute of Iran, No. 69, Pasteur Ave, 1316943551, Tehran, Iran.  
E-mail: chinikar@pasteur.ac.ir; sadeghchinikar@yahoo.com

*Received:* 5 April 2010

*Accepted in revised form:* 8 November 2010