

# Bacterial interference with host epithelial junctional complexes: Probiotic bacteria vs. A/E lesion-forming *Escherichia coli*

Topouzova-Hristova Tania, Stephanova Elena, Stoitsova Stoyanka

► **To cite this version:**

Topouzova-Hristova Tania, Stephanova Elena, Stoitsova Stoyanka. Bacterial interference with host epithelial junctional complexes: Probiotic bacteria vs. A/E lesion-forming *Escherichia coli*. *Journal of BioScience and Biotechnology*, 2012, 1 (2), pp.83-89. pasteur-00748280

**HAL Id: pasteur-00748280**

**<https://hal-riip.archives-ouvertes.fr/pasteur-00748280>**

Submitted on 5 Nov 2012

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## MINI-REVIEW

Tania Topouzova-Hristova<sup>1</sup>  
Elena Stephanova<sup>1</sup>  
Stoyanka Stoitsova<sup>2</sup>

## Bacterial interference with host epithelial junctional complexes: Probiotic bacteria vs. A/E lesion-forming *Escherichia coli*

### Authors' addresses:

<sup>1</sup>Faculty of Biology, Sofia University, Sofia-1000, Bulgaria.

<sup>2</sup>Institute of Microbiology "Stephan Angeloff", Bulgarian Academy of Sciences, Sofia-1113, Bulgaria.

### Correspondence:

Stoyanka Stoitsova  
Institute of Microbiology "Stephan Angeloff", Bulgarian Academy of Sciences, Acad. G. Bonchev St., Bl. 26 Sofia-1113, Bulgaria  
Tel.: +359-2 9793159  
e-mail: stoitsova\_microbiobas@yahoo.com

### Article info:

Received: 29 July 2012

Accepted: 30 August 2012

### ABSTRACT

During colonization, enteropathogenic (EPEC) and enterohaemorrhagic (EHEC) *Escherichia coli* are capable to manipulate host cytoskeleton and colonize gut epithelia by a specific mode of attachment known as the attaching and effacing lesion (A/E lesion). While actin rearrangements during A/E lesion formation have been extensively investigated, the possible alterations of other cytoskeletal elements like those comprising the intercellular junctional complexes (JC) of polarized cells during infection have only lately attracted attention. The present mini-review addresses the opposite effects of two groups of bacteria, A/E lesion-forming pathogenic *E. coli* and probiotic bacterial strains, on JC. JC are important in maintaining gut barrier functions. EPEC and EHEC can disrupt JC which as a consequence leads to reduction in the transepithelial electrical resistance (TER) and an increase of the permeability to macromolecules. Probiotic bacteria on the other hand stabilize JC thus increasing TER and reducing permeability to macromolecular markers. Probiotic strains can protect JC integrity of polarized cells from the damage caused by EPEC or EHEC. Together with the promise of these results, of concern is the fact that the outcome of the studies can differ dependent on experimental protocols. Studies with living bacteria and different strain combinations have also put forward strain specific effects. Therefore, an important practical item for future studies is the identification of the molecules synthesized by probiotic bacteria that may be active on JC stability.

**Key words:** EPEC, EHEC, probiotic bacteria, epithelial junctional complexes

### Introduction

Enteropathogenic and enterohaemorrhagic *Escherichia coli* (resp., EPEC and EHEC) are adapted to colonize gut epithelia by a specific mode of attachment known as attaching and effacing lesion (A/E lesion). This is produced due to the capacity of A/E pathogens to manipulate host cytoskeleton. Using their Type III secretion machinery, the bacteria inject into the eukaryotic cell a translocated intimin receptor (Tir). Tir is then integrated in the host cell membrane and interacts with intimin molecules available at the bacterial surface thus enforcing the pathogen's adherence. Epithelial microvilli in such loci lose their typical structure and form pedestals where actin accumulates under the elevated cell membrane (Frankel & Philips, 2008).

This type of lesion has been known for decades, and the molecular mechanisms by which EPEC and EHEC manipulate the host-cell actin cytoskeleton have been extensively examined. Meanwhile the putative alterations of other cytoskeletal components have been more or less neglected.

One of the basic disease symptoms that accompany EPEC and EHEC infections is diarrhea. An important pathogenic phenomenon related with diarrhea is the damage of the barrier functions of the intestine leading to uncontrolled influx of water and solutes. These barrier functions in the healthy intestine are maintained due to the integrity of the epithelial junctional complexes (JC). Nevertheless, it was only lately that the effects of EPEC and EHEC on JC have drawn attention.

**MINI-REVIEW**

As an opposite, recent studies have shown the beneficial effects of probiotic bacteria on JC integrity. The present review will focus on the mode of JC damage by A/E lesion-forming *E. coli* and the promise of the probiotic application against this.

**Junctional complexes and epithelial barrier damage by A/E pathogens**

Interaction of A/E bacteria with the polarized (epithelial) host cells is accompanied by a complex set of events related to cytoskeleton rearrangement and impairment of the epithelial barrier function. The epithelial barrier function is maintained by two large membrane domains (apical and basolateral), the junctional complex between adjacent cells, composed of the tight junction (TJ), adherens junction (AJ), and desmosomes (D). Bundles of actin filaments support the subapical membrane and some specialized membrane structures such as microvilli and junctional complexes (TJ and AJ).

The main cellular structure that maintains cell polarity and does not allow lateral diffusion of apical proteins as well as pericellular permeability of large soluble molecules is the **tight junction**. Tight junctions are composed of a branching network of transmembrane proteins (claudins and occludins) embedded in both contact membranes, with extracellular domains joining to one another. The cytoplasmic actin directly interacts with claudins and with the zonula occludens (ZO) family of cytoplasmic proteins (Delorme-Axford & Coyne, 2011). ZO-1 binds numerous transmembrane and cytoplasmic proteins and is required for the assembly of both adherens and tight junctions. Depletion of ZO-1 leads to reversible loss of junctional integrity and barrier function and alterations in transepithelial permeability (Van Itallie *et al.*, 2009; Yu *et al.*, 2010).

Another junction, important for epithelial integrity sustaining, localized near the TJ, is the **adherens junction** (AJ). The adhesive properties of AJ are determined by specialized integral membrane proteins comprising the protein family of cadherins. E-cadherin is a protein specific for epithelial cells, and is functionally linked to the generation of a polarized epithelial phenotype. The extracellular region of this protein binds to cadherins of adjacent cells and the intracellular regions of e-cadherins interact with cytoplasmic actin by catenins and other regulatory proteins (Tian *et al.*, 2011). *In vitro* studies have shown that E-cadherin is not only necessary for adherens junction formation but is also essential for the assembly of

other junctional complexes such as desmosomes, gap junctions and tight junctions (Tunggal *et al.*, 2005).

During certain disease states, including viral or A/E pathogens infection, the ability of the intestinal epithelium to regulate absorption and secretion is severely disturbed, resulting in diarrhea (Shifflett *et al.*, 2005; Delorme-Axford & Coyne, 2011). Despite the advances made in understanding of EPEC and EHEC pathogenesis at the molecular level, how the infection triggers diarrhea is still unclear. Both EPEC and EHEC alter intestinal epithelial barrier function by TJ disruption and loss of interaction of the transmembrane TJ proteins occludin and claudin-1 with the cytosolic plaque protein ZO-1, which links them to cortical F-actin (Hanajima-Ozawa *et al.*, 2007). EHEC lower transepithelial electric resistance (TER), increase the paracellular flux of mannitol, and alter ZO-1 distribution by the signaling pathways including myosin light chain kinase and conventional protein kinase C (PKC) (Philpott *et al.*, 1998; Berkes *et al.*, 2003). In contrast, conventional PKCs do not appear to participate in the EPEC-associated disruption of TJs (Berkes *et al.*, 2003). *In vitro* studies have demonstrated that the EPEC effector protein EspF plays a central role in decreasing TER and altering the intestinal epithelial TJ structure. Specifically, EPEC induce disruption of TJ architecture as evidenced by a loss of TJ protein-protein interactions, redistribution of TJ proteins, and the appearance of aberrant TJ strands in the lateral membrane (Shifflett *et al.*, 2005; Hanajima-Ozawa *et al.*, 2007). Surprisingly, recruitment of ZO-1 only does not lead to the loss of TER hence other proteins may be involved in these pathological conditions. The knockout of ZO-1 in mouse epithelial cells causes a pronounced delay in junction assembly, but has no effect on already mature junctions so apparently ZO-1 is not essential for tight junction formation (McNeil *et al.*, 2006).

In a recent study on *in vivo* effect of EHEC infection on murine intestinal barrier function, no alterations of ZO-1 distribution was detected, but reduction of TER, redistribution of occludin and claudin-3 and increased expression of claudin-2 was observed in the colon of infected mice 5-10 days after infection (Roxas *et al.*, 2010). These controversial results need more comprehensive studies on the molecular mechanisms of infection-induced JC alterations.

An important factor known to regulate epithelial monolayer permeability and TJ stability is the adherens junction, composed mainly of the transmembrane protein E-cadherin and cytosolic catenins. It has been shown that EPEC infection induces significant changes in the adherens

## MINI-REVIEW

junctions of Caco-2 monolayers, accompanied by significant phosphorylation of PKC- $\alpha$  associated with cadherins, leading to the dissociation of the cadherin/beta-catenin complex (Malladi et al., 2004). Some pathogenic bacteria, such as enteropathogenic *Escherichia coli*, *Shigella flexneri*, and *Campylobacter jejuni*, cleaved E-cadherin of host cells to disrupt the epithelial barrier, but it remained unknown whether this may be a general virulence mechanism (Hoy et al., 2012).

### **Probiotic bacteria can counteract A/E lesion-forming pathogenic bacteria by stabilizing junctional complexes and epithelial barrier function**

According to the FAO/WHO definition, probiotics are live non-pathogenic microorganisms that have beneficial effects on human health:

[http://www.fao.org/es/esn/food/foodandfood\\_probio\\_en.stm](http://www.fao.org/es/esn/food/foodandfood_probio_en.stm)

The mechanisms by which different probiotics achieve this are variable, numerous and are beyond the scope of this review.

Lately, the capacity of live probiotic bacteria to promote gut barrier function has been the scope of an increased number of studies and the focus of several reviews (Liévin-Le Moal & Servin, 2006; Walter, 2008; Guttman & Finlay, 2009; Lutgendorff et al., 2009.; Kim et al., 2010; Ohland & MacNaughton, 2010; Ahrne et al., 2011; Gill et al., 2011; Ulluwishewa et al., 2011). Using *in vitro* models of cell culture monolayers, probiotic strains have been shown to increase TER, reduce permeability to macromolecules and stimulate the expression of JC proteins. This was monitored in various experiments by TER measurements, Western blotting, qRT-PCR and microarray techniques. On the cellular level, the probiotic bacteria when applied alone have been shown to promote the synthesis of JC components by the cultured cells. This kind of stabilization, in particular that of the TJ, could be a most probable reason for maintaining TER and permeability values even after prolonged co-cultivation with cell cultures in the absence of antibiotics. Table 1. provides several examples on the mechanisms of the action of probiotic bacteria: *Lactobacillus plantarum*, *L. rhamnosus*, *L. casei* and *E. coli* Nissle 1917, on cultured-cell monolayers.

These effects further proved beneficial in the presence of diarrheagenic *E. coli* from several pathotypes (Table 1).

Importantly, in the strain and dose combinations used in the cited studies, the probiotic bacteria had no antibacterial activity against the pathogens. Hence the effects on JC stability can be considered as pivotal for the observed outputs. While very promising, these data are nevertheless not easy to compare with one another due to the variety of applied experimental protocols (Table 1). Bacteria in the test systems were used in different amounts and proportions. Some data imply that pre-treatment with the probiotic strain was decisive for the beneficial effect (Sherman et al., 2005; Johnson-Henry et al., 2008). In other cases, co-cultivation of the strains was performed (Qin et al., 2009; Liu et al., 2010, Zyrek et al., 2007). Only a few data have shown that the probiotic, if applied after the EPEC, can result in some restoration of TER values and reduction of the JC damage (Parassol et al., 2005). And finally, strain-specific effects have to be underlined as well, and this concerns both the *E. coli* and the probiotic strains (Stöber et al., 2010).

*In vivo* evidence for effects of probiotic bacteria on uninfected or infected animal models are even sparser. In rats, *L. acidophilus* has been shown to increase occludin expression and maintain gut epithelial tight junctions (Qin et al., 2005). In germfree mice, *L. reuteri* ATCC PTA 6475 protected the animals from disease manifestation caused by EHEC (Eaton et al., 2011).

### **Metabolic products and molecular mechanisms related with the effects of probiotic bacteria and A/E lesion-forming pathogens on epithelial JC: questions and perspectives**

How do these *in vitro* results rely to situations of heavy infectious diarrheas?

Together with strain-specific differences, when the application of live probiotic bacteria is considered usually time is needed to settle the correct type of ecosystem that will ensure the probiotic' benefits. Unfortunately, in infections with A/E lesion-forming pathogens there is rarely a lot of time available. A/E infections are acute, and especially in cases of Shiga toxin-producing strains there is a high risk. Antibiotics are undesirable since massive death of bacteria will result in release of high amounts of toxins thus increasing greatly the danger for the patient. Prospects were sought in the application of bacteriophages, bacteriocins, and low molecular-weight inhibitors, but for the time being in trials none of the interventions proved superior to supportive

## MINI-REVIEW

Table 1. Protective mechanisms of live probiotic bacteria on epithelial barrier integrity in the absence or presence of pathogenic *E. coli*.

Probiotic strain	In interaction with:		References
	Eukaryote model	<i>E. coli</i> strain	
<i>L. plantarum</i> MB452	Caco-2	-	Anderson et al., 2010
<i>L. plantarum</i> CGMCC 1258	Caco-2; NCM460	<i>E. coli</i> O124:NM ATCC 43893 (EIEC); <i>E. coli</i> O111:NM ATCC 43887 (EPEC)	Qin et al., 2009 ; Liu et al., 2010
<i>L. rhamnosus</i> OLL2838	Isolated mouse intestinal epithelial cells	-	Miyauchi et al., 2009
<i>L. rhamnosus</i> R0011	Hep-2, T84	<i>E. coli</i> O157:H7 (EHEC); <i>E. coli</i> O127:H6 (EPEC)	Sherman et al., 2005
<i>L. rhamnosus</i> GG	MDCK-1 and T84	<i>E. coli</i> O157:H7 CL56	Johnson-Henry et al., 2008
<i>L. casei</i> DN-114 001	T84	<i>E. coli</i> E2348/69 (EPEC)	Parassol et al., 2005
<i>L. acidophilus</i> , <i>L. fermentum</i> , <i>L. gasseri</i> , <i>L. rhamnosus</i>	T84	-	Hummel et al., 2012
<i>E. coli</i> Nissle 1917	T84	-	Veltman et al., 2012
<i>E. coli</i> Nissle 1917	T84	<i>E. coli</i> E2348/69 (EPEC)	Zyrek et al., 2007

Legend: TER - transepithelial electric resistance; TJ - tight junction; EHEC - enteroinvasive *E. coli*; FAS - Fluorescent actin staining test for actin accumulation. Cells shaded green: probiotic strains alone; white cells - protective effects of probiotic strains against *E. coli*-induced damage.

## MINI-REVIEW

therapy alone (Muniesa *et al.*, 2012).

Provided the beneficial effects of probiotic strains on JC stability and epithelial barrier function, one reasonable task would be the identification of the molecules responsible for these effects. Such molecules may prove useful in therapy by themselves, without the need to apply the live probiotics and wait for the ecosystem to settle.

To check for JS-stabilizing activity of released products, cell-free supernatants of probiotic bacteria were applied to cultured cells. It was shown that cell-free supernatant of *Bifidobacterium lactis* and *B. infantis* reduced TER and prevented the EHEC-induced damage of tight junctions (Ewaschuk *et al.*, 2008; Putaala *et al.*, 2008). Apart from the stabilization of epithelia, released products (total supernatants and selected fractions from them) from *L. acidophilus* La-5 and *L. reuteri* ATCC 55730 were shown to influence EHEC virulence by reducing extracellular concentrations of autoinducer-2 and suppressing quorum sensing-regulated expression of virulence-related genes (Medellin-Peña *et al.*, 2007; Jelčić *et al.*, 2008; Medellin-Peña & Griffiths, 2009). Still of concern, when working with spent culture media from probiotic strains there remains the possibility of strain-specific responses of the A/E-forming pathogens.

The promise in these types of studies is the expectation for a further more precise identification of the molecules responsible for JC stabilization.

One group of such molecules that lately attracted attention is the surface proteins of lactobacilli. Thus, the surface layer adhesive protein (SLAP) from *L. plantarum* CGMCC 1258, while inhibiting EPEC adhesion to Caco-2 cells, stimulated the expression of TJ- associated proteins and promoted the maintenance of TER and TJ structure, the levels of extracellular signal-regulated kinase (EPK) and EPK phosphorylation (Liu *et al.*, 2011b).

Another surface protein in focus is the micro integral membrane protein (MIMP) from *L. plantarum* CGMCC 1258. NCM460 cells transiently infected with MIMP are characterized by increased synthesis of the TJ protein claudin-1 by NCM460 cells. Together with this the protein reduced the attachment of EPEC to the cultured cells (Liu *et al.*, 2011a).

### Conclusion

While still quite new, the studies on the interactions between A/E lesion-forming bacteria, probiotic bacteria and epithelial junctional complexes, due to the promise for

practical applications, put in focus the need of extensive studies, including more strain combinations, culture cell lines and *in vivo* models. For the time being, we have already some clarity on the cellular targets (e.g., the opposite action of probiotic bacteria and A/E lesion forming bacteria on JC and their components). Yet, by measuring the effects on TER, or JC molecules amounts or expression, we still address the output and not the mechanisms. And these seem to be the correct address for future work.

### Acknowledgement

The support of National Science Fund, Republic of Bulgaria, DO02-301, is thankfully acknowledged.

### References

- Ahrne S, Johansson Hagslatt M-L. 2011. Effect of lactobacilli on paracellular permeability in the gut. *Nutrients*, 3(1): 104-117.
- Anderson RC, Cookson AL, McNabb WC, Park Z, McCann MJ, Kelly WJ, Roy NC. 2010. *Lactobacillus plantarum* MB452 enhances the function of the intestinal barrier by increasing the expression levels of genes involved in tight junction formation. *BMC Microbiology*, 10: 316.
- Berkes J, Viswanathan V, Savkovic S, Hecht G. 2003. Intestinal epithelial responses to enteric pathogens: effects on the tight junction barrier, ion transport, and inflammation. *Gut*, 52(3): 439-451.
- Delorme-Axford E, Coyne CB. 2011. The actin cytoskeleton as a barrier to virus infection of polarized epithelial cells. *Viruses*, 3(12): 2462-2477.
- Eaton KA, Honkala A, Auchtung TA, Britton RA. 2011. Probiotic *Lactobacillus reuteri* ameliorates disease due to enterohemorrhagic *Escherichia coli* in germfree mice. *Infect. Immun.*, 79(1): 185-191.
- Ewaschuk JB, Diaz H, Meddings L, Diederichs B, Dmytrash A, Backer J, Loojer-van Langen M, Madsen KL. 2008. Secreted bioactive factors from *Bifidobacterium infantis* enhance epithelial cell barrier function. *Am. J. Physiol. Gastrointest. Liver. Physiol.*, 295(5): G1025- G1034.
- FAO/WHO. 2001. [http://www.fao.org/es/esn/food/foodandfood\\_](http://www.fao.org/es/esn/food/foodandfood_)
- Frankel G., Phillips AD. 2008. Attaching effacing *Escherichia coli* and paradigms of Tir-triggered actin polymerization: getting off the pedestal. *Cellular Microbiol.*, 10(3): 549-556.
- Gill N, Wlodarska M, Finlay BB. 2011. Roadblocks in the gut: barriers to enteric infection. *Cellular Microbiol.*, 13(5): 660-669.
- Guttman JA, Finlay BB. 2009. Tight junctions as targets of infectious agents. *Biochim. Biophys. Acta*, 1788(4): 832-841.
- Hanajima-Ozawa M, Matsuzawa T, Fukui A, Kamitani Sh, Ohnishi H, Abe A, Horiguchi Y, Miyake M. 2007. Enteropathogenic *Escherichia coli*, *Shigella flexneri*, and *Listeria monocytogenes* recruit a junctional protein, Zonula occludens-1, to actin tails and pedestals. *Infect. Immun.*, 75(2): 565-573.
- Hoy B, Geppert T, Boehm M, Reisen F, Plattner P, Gadermaier G,

## MINI-REVIEW

- Sewald N, Ferreira F, Briza P, Schneider G, Backert S, Wessler S. 2012. Distinct roles of secreted HtrA proteases from gram-negative pathogens in cleaving the junctional protein and tumor suppressor E-cadherin. *J. Biol. Chem.*, 287(13): 10115-10120.
- Hummel S, Veltman K, Cichon C, Sonnenborn U, Schmidt MA. 2012. Differential targeting of the E-cadherin/ $\beta$ -catenin complex by Gram-positive probiotic lactobacilli improves epithelial barrier function. *Appl. Environ. Microbiol.* 78(4): 1140-1147.
- Jelčić I, Hüfner E, Schmidt H, Hertel C. 2008. Repression of the locus of enterocyte effacement-encoded regulator of gene transcription of *Escherichia coli* O157:H7 by *Lactobacillus reuteri* culture supernatant is LuxS and strain dependent. *Appl. Environ. Microbiol.*, 74(10): 3310-3314.
- Johnson-Henry KC, Donato KA, Shen-Tu G, Gordanpour M, Sherman PM, 2008. *Lactobacillus rhamnosus* strain GG prevents enterohemorrhagic *Escherichia coli* O157:H7-induced changes in epithelial barrier function. *Infect. Immun.*, 76(4): 1340-1348.
- Kim M, Ashida H, Ogawa M, Yoshokawa Y, Mimuro H, Sasakawa C. 2010. Bacterial interactions with the host epithelium. *Cell Host Microbe*, 8(1): 20-35
- Liévin-Le Moal V, Servin AL. 2006. The front line of enteric host defense against unwelcome intrusion of harmful microorganisms: mucins, antimicrobial peptides, and microbiota. *Clin. Microbiol. Rev.*, 19(2): 315-337.
- Liu Z, Shen T, Zhang P, Ma Y, Moyer MP, Qin H. 2010. Protective effects of *Lactobacillus plantarum* against epithelial barrier dysfunction of human colon cell line NCM460. *World J Gastroenterol.*, 16(45): 5759-5765.
- Liu Z, Ma Y, Yang J, Zhang P, Moyer MP, Qin H. 2011a. Expression of the *Lactobacillus plantarum* surface layer MIMP protected the NCM460 epithelial cells from enteroinvasive *Escherichia coli* infection. *Cellular Physiol. Biochem.*, 27(1): 99-108.
- Liu Z, Shen T, Zhang P, Ma Y, Qin H. 2011b. *Lactobacillus plantarum* surface layer adhesive protein protects intestinal epithelial cells against tight junction injury induced by enteropathogenic *Escherichia coli*. *Mol. Biol. Rep.*, 38(5): 3471-3480.
- Lutgendorff F, Nijmeijer R, Sandström, Trulsson L, Magnusson K-E, Timmerman H, van Minnen L, Rijkers G, Gooszen H, Akkermans L, Söderholm J. 2009. Probiotics prevent intestinal barrier dysfunction in acute pancreatitis in rats via induction of ileal mucosal glutathione biosynthesis. *PLoS ONE*, 4(2): e4512.
- Malladi V, Shankar B, Williams P, Balakrishnan A. 2004. Enteropathogenic *Escherichia coli* outer membrane proteins induce changes in cadherin junctions of Caco-2 cells through activation of PKC $\alpha$ . *Microbes Infect.* 6(1): 38-50.
- McNeil E, Capaldo CT, Macara IG. 2006. Zonula occludens-1 function in the assembly of tight junctions in Madin-Darby Canine kidney epithelial cells. *Mol. Biol. Cell*, 17(4): 1922-1932.
- Medellin-Peña MJ, Griffiths MW. 2009. Effect of molecules secreted by *Lactobacillus acidophilus* strain La-5 on *Escherichia coli* O157:H7 colonization. *Appl. Environ. Microbiol.*, 75(4): 1165-1172.
- Medellin-Peña MJ, Wang H, Johnson R, Anand S, Griffiths MW. 2007. Probiotics affect virulence-related gene expression in *Escherichia coli* O157:H7. *Appl. Environ. Microbiol.*, 73(13): 4259-4267.
- Miyauchi E, Morita H, Tanabe S. 2009. *Lactobacillus rhamnosus* alleviates intestinal barrier dysfunction in part by increasing the expression of Zonula occludens-1 and myosin light-chain kinase in vivo. *J. Dairy Sci.*, 92(6): 2400-2408.
- Muniesa M, Hammerl J, Hertwig S, Appel B, Brüssow H. 2012. Shiga toxin-producing *Escherichia coli* O104:H4: a new challenge for microbiology. *Appl. Environ. Microbiol.*, 78(12): 4065-4073.
- Ohland CL, MacNaughton WK. 2010. Probiotic bacteria and intestinal epithelial barrier function. *Am. J. Physiol. Gastrointest. Liver Physiol.* 298(6): G807-G819.
- Parassol N, Freitas M, Thoreux K, Dalmasso G, Bourdet-Sicard R, Rampal P. 2005. *Lactobacillus casei* DN-114 001 inhibits the increase in paracellular permeability of enteropathogenic *Escherichia coli*-infected T84 cells. *Res. Microbiol.*, 156(2): 256-262.
- Philpott DJ, McKay DM, Mak W, Perdue MH, Sherman PM. 1998. Signal transduction pathways involved in enterohemorrhagic *Escherichia coli*-induced alterations in T84 epithelial permeability. *Infect. Immun.*, 66(4): 1680-1687. [probio\\_en.stm](http://probio_en.stm)
- Putala H, Salusjärvi T, Nordström M, Saarinen M, Ouweland A, Bech Hansen E, Rautonen N. 2008. Effect of four probiotic strains and *Escherichia coli* O157:H7 on tight junction integrity and cyclo-oxygenase expression. *Res. Microbiol.*, 159(9-10): 692-698.
- Qin H, Shen T, Gao Z, Fan X, Hang X, Jiang Y, Zhang H. 2005. Effects of lactobacillus on gut microflora and barrier function of the rats with abdominal infection. *World J. Gastroenterol.*, 11(17): 2591-2596.
- Qin H, Zhang Z, Hang X, Jiang Y. 2009. *L. plantarum* prevents enteroinvasive *Escherichia coli*-induced tight junction proteins changes in intestinal epithelial cells. *BMC Microbiology*, 9: 63.
- Roxas J, Koutsouris A, Bellmeyer A, Tesfay S, Royan S, Falzari K, Harris A, Cheng H, Rhee K J, Hecht G. 2010. Enterohemorrhagic *E. coli* alters murine intestinal epithelial tight junction protein expression and barrier function in a Shiga toxin independent manner. *Laboratory Investigation*, 90(8): 1152-1168.
- Sherman PM, Johnson-Henry KC, Yeung HP, Ngo PS, Goulet J, Tompkins TA. 2005. Probiotics reduce enterohemorrhagic *Escherichia coli* O127:H6-induced changes in polarized T84 epithelial cell monolayers by reducing bacterial adhesion and cytoskeletal rearrangements. *Infect. Immun.*, 73(8): 5183-5188.
- Shifflett DE, Clayburgh DR, Koutsouris A, Turner JR, Hecht GA. 2005. Enteropathogenic *E. coli* disrupts tight junction barrier function and structure *in vivo*. *Laboratory Investigation*, 85(10): 1308-1324.
- Stöber H, Maier E, Schmidt H. 2010. Protective effects of Lactobacilli, Bifidobacteria and Staphylococci on the infection of cultured HT29 cells with different enterohemorrhagic *Escherichia coli* serotypes are strain-specific. *Int. J. Food. Microbiol.*, 144(1): 133-140.
- Tian X, Liu Zh, Niu B, Zhang J, Tan Th, Lee S, Zhao Y, Harris D, Zheng G. 2011. E-cadherin/ $\beta$ -catenin complex and the epithelial barrier. *J. Biomed. Biotechnol.*, 2011, Article ID 567305, doi:10.1155/2011/567305.
- Tunggal JA, Helfrich I, Schmitz A, Schwarz H, Günzel D, Fromm M, Kemler R, Krieg T, Niessen CM. 2005. E-cadherin is essential for in vivo epidermal barrier function by regulating tight junctions. *EMBO J.*, 24(6): 1146-1156.
- Ulluwishewa D, Anderson RC, McNabb W, Moughan PJ, Wells JM,

**MINI-REVIEW**

- Roy NC. 2011. Regulation of tight junction permeability by intestinal bacteria and dietary components. *J. Nutr.*, 141(5): 769-776.
- Van Itallie CM, Fanning AS, Bridges A, Anderson JM. 2009. ZO-1 stabilizes the tight junction solute barrier through coupling to the perijunctional cytoskeleton. *Mol. Biol. Cell*, 20(17): 3930–3940.
- Veltman K, Hummel S, Cichon C, Sonnenborn U, Schmidt M. 2012. Identification of specific miRNAs targeting proteins of the apical junctional complex that stimulate the probiotic effect of *E. coli* Nissle 1917 on T84 epithelial cells. *Int. J. Biochem. Cell Biol.*, 44(2): 341-349.
- Walter J. 2008. Ecological role of lactobacilli in the gastrointestinal tract: implications for fundamental and biomedical research. *Appl. Environ. Microbiol.*, 74(16): 4985-4996.
- Yu D, Marchiando AM, Weber CR, Raleigh DR, Wang Y, Shen L, Turner JR. 2010. MLCK-dependent exchange and actin binding region-dependent anchoring of ZO-1 regulate tight junction barrier function. *PNAS*, 107(18): 8237–8241.
- Zyrek AA, Cichon C, Helms S, Enders C, Sonnenborn U, Schmidt MA. 2007. Molecular mechanisms underlying the probiotic effects of *Escherichia coli* Nissle 1917 involve ZO-2 and PKC $\zeta$  redistribution in tight junction and epithelial barrier repair. *Cellular Microbiol.*, 9(3): 804-816.