



HAL
open science

Rapid identification of mycobacteria to species level by PCR-restriction fragment length polymorphism analysis of the hsp65 gene and proposition of an algorithm to differentiate 34 mycobacterial species.

A. Devallois, K. S. Goh, N. Rastogi

► To cite this version:

A. Devallois, K. S. Goh, N. Rastogi. Rapid identification of mycobacteria to species level by PCR-restriction fragment length polymorphism analysis of the hsp65 gene and proposition of an algorithm to differentiate 34 mycobacterial species.. *Journal of Clinical Microbiology*, 1997, 35 (11), pp.2969-73. pasteur-00931152

HAL Id: pasteur-00931152

<https://riip.hal.science/pasteur-00931152>

Submitted on 17 Feb 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Rapid identification of mycobacteria to species level by PCR-restriction fragment length polymorphism analysis of the hsp65 gene and proposition of an algorithm to differentiate 34 mycobacterial species.

A Devallois, K S Goh and N Rastogi
J. Clin. Microbiol. 1997, 35(11):2969.

Updated information and services can be found at:
<http://jcm.asm.org/content/35/11/2969>

CONTENT ALERTS

These include:

Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), [more»](#)

Information about commercial reprint orders: <http://journals.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Rapid Identification of Mycobacteria to Species Level by PCR-Restriction Fragment Length Polymorphism Analysis of the *hsp65* Gene and Proposition of an Algorithm To Differentiate 34 Mycobacterial Species

ANNE DEVALLOIS, KHYE SENG GOH, AND NALIN RASTOGI*

Unité de la Tuberculose et des Mycobactéries, Institut Pasteur, Morne Jolivière B.P. 484,
97165 Pointe à Pitre Cedex, Guadeloupe, French West Indies

Received 23 May 1997/Returned for modification 10 July 1997/Accepted 19 August 1997

PCR-restriction fragment length polymorphism analysis (PRA) of the *hsp65* gene (A. Telenti, F. Marchesi, M. Balz, F. Bally, E. C. Böttger, and T. Bodmer, *J. Clin. Microbiol.* 31:175–178, 1993) was applied to 108 mycobacterial isolates representing 34 species to evaluate its potential as a rapid reference method. A total of 49 distinct patterns were obtained; 25 species were characterized by a single PRA pattern, while 9 species gave more than one specific pattern. An algorithm describing these 34 species (which includes five additional species and new subgroups of *Mycobacterium kansasii*, *M. abscessus*, and *M. peregrinum*) is proposed. A relatively simple and inexpensive method, PRA may be particularly helpful in routine clinical microbiology laboratories.

Classical identification results for mycobacteria based on cultural and biochemical tests may take several weeks after reception of specimens, and the tests sometimes fail to produce a precise identification. Additional techniques such as thin-layer chromatography (4, 16), gas-liquid chromatography (14), high-performance liquid chromatography (5), and gene sequencing (3, 11, 12, 22, 23) are powerful tools but unfortunately remain limited to reference laboratories. Recent molecular techniques based on probe hybridization (7, 15) or on amplification of a specific genomic region, such as the AmpliCor MTB test (6), provide results limited to a single species per experiment. In contrast to the above-mentioned techniques needing specialized equipment, PCR-restriction fragment length polymorphism analysis (PRA) (26) of the *hsp65* gene present in all mycobacteria (9, 20, 21) offers an easy, rapid, and inexpensive procedure to identify several mycobacterial species in a single experiment. *hsp65* PRA was also used to identify some slowly growing mycobacteria (18), as well as clinical isolates cultured in Bactec 460-TB vials (25). Other PCR-restriction fragment length polymorphism methods based on the analysis of digestion products of specific genes, such as 16S rRNA (2, 10, 29) or *dnaJ* genes (24), using three to five different restriction enzymes have also been reported but remain cumbersome compared to *hsp65* PRA, which uses only two restriction enzymes.

Our primary aim was to evaluate the PRA method and to extend it to other mycobacterial species that were not studied previously, with particular emphasis on the differentiation of *Mycobacterium chelonae*, *M. abscessus*, *M. fortuitum*, and *M. peregrinum*. These rapidly growing nonchromogenic mycobacteria are often found in drinking water and are of significant clinical importance because of their multiple drug resistance (8). These four species were previously referred as the *M. fortuitum* complex (composed of *M. fortuitum* var. *fortuitum*, *M. fortuitum* var. *peregrinum*, *M. chelonae* subsp. *chelonae*, and *M. chelonae* subsp. *abscessus*) and could be differentiated from other rapidly growing nonchromogenic mycobacteria by an arylsulfatase-positive test result (4). Differentiation between

M. chelonae and *M. abscessus* and between *M. fortuitum* and *M. peregrinum* has been achieved by cultural and biochemical tests, thin-layer chromatography of mycolic acids, and mycobactin analysis (4, 14). In the present investigation, we have applied PRA followed by computer analysis to both reference and clinical isolates and propose a more complete algorithm by adding five more species, as well as new subgroups of *M. kansasii*, *M. abscessus*, and *M. peregrinum*.

A total of 108 mycobacterial strains analyzed by PRA comprised 43 reference strains (Table 1) and 65 clinical isolates, which were identified by classical biochemical identification tests and the Accuprobe tests and DT1-DT6 PCR (7) for the *M. avium* complex isolates. *M. chelonae*, *M. abscessus*, *M. fortuitum*, and *M. peregrinum* strains from a previous study (4) were identified on the basis of cultural and biochemical tests, mycobactin patterns, and mycolid acid content. Clinical isolates of *M. kansasii* were from a previous study (28) and were obtained from E. Tortoli, Florence, Italy. Bacterial DNA was prepared as follows: one loopful of bacteria was suspended in 300 μ l of TE (10 mM Tris, 1 mM EDTA) and 100 μ l of acid-washed glass beads (diameter, <106 μ m; Sigma, St. Louis, Mo.), heated at 94°C for 15 min, and sonicated at 35 kHz for 15 min (water bath sonicator from Gen-Probe Inc., San Diego, Calif.). Five microliters of the supernatant containing the crude DNA extract was used for PCR. PRA based on the amplification of a 439-bp fragment of the *hsp65* gene was performed with primers Tb11 (5'-ACCAACGATGGTGTGTCCAT) and Tb12 (5'-CTTGTCTGAACCGCATACCCT) by the method of Telenti et al. (26). Twenty-two microliters of the PCR product was digested by *Bst*EII (Promega, Madison, Wis.) or *Hae*III (BioLabs, Inc., Beverly, Mass.), and 12 μ l of the restriction digest was loaded on a 4% (wt/vol) NuSieve 3:1 agarose gel (FMC Bioproducts, Rockland, Maine). A 100-bp ladder (Pharmacia Biotech, Uppsala, Sweden) served as an external molecular size marker and was added after every six lanes of migration to reduce migration-related errors. Restriction patterns were videocopied by using the Gel-Analyst software (Bioprobe Systems, Montreuil, France). The Taxotron package software (P. A. D. Grimont, Institut Pasteur, Paris,

* Corresponding author. Phone: 590-893-881. Fax: 590-893-880.

TABLE 1. Reference strains used in the present study

Species or isolate(s)	Strain ^a
<i>M. abscessus</i>	ATCC 19977
<i>M. africanum</i>	ATCC 25420
<i>M. asiaticum</i>	ATCC 25276
<i>M. avium</i>	ATCC 25291
<i>M. bovis</i>	ATCC 19210
<i>M. bovis BCG</i>	CIPT 140040001
Crohn's disease isolates	2569 (CDI), Lyon (EVL)
<i>M. chelonae</i>	NCTC 946
" <i>M. engbackii</i> "	ATCC 27353
<i>M. flavescens</i>	ATCC 14474
<i>M. fortuitum</i>	ATCC 6841
<i>M. gastri</i>	ATCC 15754
<i>M. gordonae</i>	ATCC 14470
<i>M. haemophilum</i>	ATCC 29548, CIPT 141420006
<i>M. intracellulare</i>	ATCC 13950
<i>M. kansasii</i>	ATCC 12478
<i>M. malmoense</i>	ATCC 29571
<i>M. marinum</i>	ATCC 927
<i>M. paratuberculosis</i>	ATCC 19698, 1077(CVM), 7912 (LCRV)
<i>M. peregrinum</i>	ATCC 19420
<i>M. phlei</i>	ATCC 11758
<i>M. porcinum</i>	ATCC 33775
<i>M. scrofulaceum</i>	ATCC 19981
<i>M. senegalense</i>	NCTC 10956
<i>M. shimoidei</i>	ATCC 27962
<i>M. simiae</i>	ATCC 25275
<i>M. smegmatis</i>	ATCC 19420
<i>M. szulgai</i>	NCTC 10831
<i>M. terrae</i>	ATCC 15755
<i>M. tuberculosis</i>	ATCC 27294, ATCC 25177, CIPT 140010059
<i>M. triviale</i>	ATCC 23292
<i>M. ulcerans</i>	ATCC 33728
Wood pigeon	6861 (LCRV), 6409 (LCRV)
<i>M. xenopi</i>	ATCC 19276, ATCC 19970

^a ATCC, American Type Culture Collection; NCTC, National Collection of Type Cultures; CIPT, Collection Institut Pasteur de Paris-Tuberculose; CDI, Centraal Diergeneeskundig Instituut, Lelystad, The Netherlands; EVL, Ecole Vétérinaire de Lyon, Lyon, France; LCRV, Laboratoire Central de Recherches Vétérinaires, Maisons-Alfort, France; CVM, College of Veterinary Medicine, Ames, Iowa.

France) was used to convert the migration file into a molecular weight data file by the Schaffer and Sederoff method.

The results obtained in this investigation are summarized in Tables 2 and 3 and Fig. 1. As noted previously (26), restriction fragments shorter than 60 bp were not taken in account as they

are suspected of being primer or primer dimer bands. Restriction fragments were visible for all the tested strains under normal conditions except for some *M. kansasii* strains which needed larger amounts of the PCR products for restriction analysis. All the 108 isolates were easily differentiated by PRA, and there was a perfect correlation between the PRA identification and the conventional identification. As summarized in Fig. 1, the restriction profiles obtained were in agreement with the algorithm of Taylor et al. (25), who have recently modified the initial algorithm of Telenti et al. (26). In addition, the algorithm proposed in Fig. 1 now incorporates newly found profiles that have not been described before (three *M. kansasii* subgroups, two *M. peregrinum* subgroups, and one *M. abscessus* subgroup) and also previously unanalyzed species such as *M. ulcerans*, "*M. engbackii*," *M. porcinum*, *M. phlei*, and *M. senegalense*.

All the members of the *M. tuberculosis* complex (three reference strains of *M. tuberculosis* and one each of *M. africanum*, *M. bovis*, and *M. bovis BCG* [5, 4, 2, and 2 clinical isolates, respectively]), gave identical PRA profiles (Fig. 1). As illustrated in Fig. 1, within the *M. avium* complex *M. avium* and *M. intracellulare* species were easily differentiated; the three strains of *M. paratuberculosis* and the two isolates of wood pigeon and Crohn's disease mycobacteria all gave the *M. avium* profile, confirming their genetic relatedness to this species (27, 30).

Only one pattern each for *M. fortuitum* subsp. *fortuitum*, *M. fortuitum* subsp. *peregrinum*, *M. fortuitum* 3rd variant, *M. chelonae* subsp. *chelonae*, and *M. chelonae* subsp. *abscessus* was initially described by Telenti et al. (26); this was followed by the description of an additional pattern for a single presumptive *M. chelonae* isolate by Taylor et al. (25). In this investigation a total of 39 isolates of this group (35 clinical isolates and 4 type strains) were studied, and in addition to the previously described patterns, one new pattern for *M. abscessus* and two for *M. peregrinum* were reported (Fig. 1). As all the 35 clinical isolates included in the present investigation have been extensively characterized (4, 14), we were able to add the new profiles to the algorithm proposed in Fig. 1. It should be further underlined that some miscellaneous strains which were not differentiated into either *M. chelonae*-*M. abscessus* or *M. fortuitum*-*M. peregrinum* in previous studies (4, 14) were easily discriminated by PRA (Table 2). In our opinion, it is important to properly classify the organisms before attributing a species-specific PRA pattern to them; e.g., an additional pattern reported for a "presumptive" *M. chelonae* isolate by Taylor et al. (25) was not reproduced for any of the 39 isolates studied here.

TABLE 2. Summary of PRA results for *M. chelonae*, *M. abscessus*, *M. fortuitum*, and *M. peregrinum*

PRA pattern	Bands with <i>Bsr</i> EII and <i>Hae</i> III (bp)	Reference strain	No. of isolates per pattern (this study)	References for pattern
<i>M. chelonae</i> I	325/140 and 210	NCTC 946	10 <i>M. chelonae</i> , 3 miscellaneous ^a	25, 26, and this study
<i>M. chelonae</i> II ^b	325/120 and 140/65/60		None	25
<i>M. abscessus</i> I	245/220 and 160/60	ATCC 19977	2 <i>M. abscessus</i> , 1 miscellaneous ^a	25, 26, and this study
<i>M. abscessus</i> II	245/220 and 210/60		2 <i>M. abscessus</i>	This study
<i>M. fortuitum</i> I	245/120/80 and 155/135	ATCC 6841	4 <i>M. fortuitum</i>	25, 26, and this study
<i>M. fortuitum</i> II ^c	245/120/80 and 150/135		1 miscellaneous ^d	25, 26, and this study
<i>M. peregrinum</i> I	245/220 and 155/150/100	ATCC 19420	4 <i>M. peregrinum</i> , 1 miscellaneous ^d	25, 26, and this study
<i>M. peregrinum</i> II	245/220 and 150/135/100		2 <i>M. peregrinum</i> , 4 miscellaneous ^d	This study
<i>M. peregrinum</i> III	245/140/80 and 155/150/100		1 <i>M. peregrinum</i>	This study

^a These isolates were previously classified as miscellaneous as differentiation between *M. chelonae* and *M. abscessus* was not achieved (4, 14).

^b Pattern reported for a single presumptive *M. chelonae* isolate (25).

^c Pattern previously reported for third-variant isolates ATCC 49403 and ATCC 49404 (26).

^d These isolates were previously classified as miscellaneous as differentiation between *M. fortuitum* and *M. peregrinum* was not achieved (4, 14).

TABLE 3. Summary of PRA results for the five subgroups of *M. kansasii*

PRA pattern	Bands with <i>Bst</i> EII and <i>Hae</i> III	Reference strain	No. of isolates per pattern (this study)	References for pattern
<i>M. kansasii</i> I	245/220 and 140/105/80	ATCC12478	7	17, 25, 26, and this study
<i>M. kansasii</i> II	245/140/80 and 140/105 ^a		8	17 and this study
<i>M. kansasii</i> III	245/140/80 and 140/105/70 ^a		None	17, 26
<i>M. kansasii</i> IV	245/120/80 and 140/115/70 ^a		None	17
<i>M. kansasii</i> V	325/120 and 140/100/80		None	17

^a A minor (5-bp) difference upon *Bst*EII digestion was found by Picardeau et al. (17).

We are consequently unable to reconfirm the inclusion of this pattern for *M. chelonae* for the time being. A summary of previous results and those found in this study is provided in Table 2.

All the 15 clinical isolates of *M. kansasii* studied here came from a previous study describing *M. kansasii* biotypes associated with AIDS in Italy (28). Indeed, on the basis of phenotypic traits based on Tween 80 hydrolysis and the α-fucosidase test and genotypic criteria such as positive or negative hybridization with the AccuProbe test (13), two significant biotypes within the *M. kansasii* species were shown to exist (28). It is interesting to note that in our study, all of the pattern I isolates

belonged to biotype 1 (i.e., AccuProbe test-positive isolates) (28), whereas pattern II isolates belonged to biotype 2 (i.e., AccuProbe test-negative isolates) (28). A majority of *M. kansasii* isolates from France also presented PRA patterns I and II (17) and had genotypes corresponding to those reported by Ross et al. (19). Apart from these two major subgroups, three other minor subgroups have also been reported (17, 26), confirming the heterogeneous nature of *M. kansasii*. These results are summarized in Table 3.

All the results obtained in this investigation, along with an update from previously published observations, have been integrated in the algorithm illustrated in Fig. 1. However, some




FIG. 1. Algorithm of PRA patterns for 34 mycobacterial species. Shaded species illustrate newer PRA patterns that were not reported in the previous algorithms of Telenti et al. (26) and Taylor et al. (25).

patterns may be close enough even upon *Hae*III digestion (isolates with a 5-bp difference for some of the bands or those varying in a single band) and it may be difficult to differentiate between such isolates without appropriate software. Examples may include differentiation between *M. scrofulaceum* and *M. peregrinum* II and between *M. fortuitum* I and *M. fortuitum* II.

Mycobacterial identification to the species level is critical step in patient management, as the results obtained influence both the choice of proper treatment and the eventual need for patient isolation. It is therefore important to develop methods that provide rapid results and that may be used in a wide variety of laboratories around the world. The latter criterion invariably requires that such methods be both simple and cost-effective.

The PRA method was selected in this investigation because of its ease and rapidity and because it may help speciate numerous species of mycobacteria within a single experiment. This broader spectrum of the PRA method, as compared to other molecular methods, is based on the amplification of a conserved *hsp65* gene present in all mycobacteria and in some other bacteria such as *Nocardia* spp. (9). The restriction analysis of a 439-bp fragment within this gene after *Bst*EII and *Hae*III digestions is highly effective for differentiating mycobacteria at the species level (26). The results obtained clearly showed distinct profiles for all the 34 species. For some species, such as *M. kansasii*, *M. chelonae*, *M. abscessus*, *M. fortuitum*, *M. peregrinum*, *M. gordonae*, *M. nonchromogenicum*, *M. simiae*, and *M. flavescens*, the discrimination was even obtained at the subgroup level. For *M. kansasii*, this subdivision was clearly linked to bacteriological and clinical specificities (28) and was also reflected by 16S rRNA sequencing (13) and other molecular techniques (1, 17, 19). It is therefore possible that the discrimination at a subgroup level for other species, although not investigated in detail for the time being, could be similarly linked to bacteriological and clinical specificities. On the other hand, the four members of the *M. tuberculosis* complex were characterized by similar PRA profiles, confirming that they do belong to a similar group genetically.

A substantial number of mycobacterial species and clinical isolates have now been run in parallel by using PRA and other reference methods (17, 25, 26; this study), and the available data have been integrated in the algorithm presented in Fig. 1. It is apparent that some species harbor unique patterns while others present several distinct profiles, which may complicate the interpretation of the results obtained. For example, a lot of PRA profiles were generated in this investigation, and had it not been for the computer-assisted analysis and interpretation of the gels, which significantly reduced the gel-to-gel variations, we may have faced problems in resolving all the patterns obtained in a short time. Furthermore, the computer-assisted analysis was particularly useful for analyzing fragments of between 100 and 200 bp.

Despite the fact that we recommend analysis of PRA profiles using appropriate software, it is important to underline that visual analysis of PRA profiles remains appropriate if related type strains are run in parallel. In our opinion, PRA is particularly useful for identifying clinical isolates giving mixed biochemical results, a situation which may require choosing between two or three closely related species. Last but not least, PRA is a rapid method which, like multiplex PCR or 16S rRNA sequencing, permits the identification of several species in a single experiment.

As far as smaller technical details are concerned, the NuSieve gel is well adapted to the migration of small DNA fragments and less cumbersome than polyacrylamide gels. As 5-bp differences in the sizes of restriction fragments may arise

between different users, we recommend that each laboratory willing to use this method first validate PRA on reference strains. Last but not least, the usual recommendations to prevent PCR-linked contamination may be useful to avoid false identification. We personally use separate manipulation rooms for reagent preparation, specimen preparation, and amplification and detection. Even specimen centrifugations were performed in a separate aerosol-free area (distinct from the above three areas).

In conclusion, PRA is a rapid, cost-effective, and efficient method for the identification of mycobacteria in a clinical microbiology laboratory. Recently, PRA was used on Bactec cultures (25), and it should now be evaluated directly for clinical specimens. If successful, PRA will be the method of choice, as opposed to nucleic acid probes, which require large number of bacilli.

We are grateful to the "Délégation Générale au Réseau International des Instituts Pasteur et Instituts Associés," Institut Pasteur, Paris, France, and the Fondation Française Raoul Follereau, Paris, France, for financial support.

We thank E. Tortoli, Careggi Hospital, Florence, Italy, for kindly providing *M. kansasii* clinical isolates.

REFERENCES

1. Abed, Y., C. Bollet, and P. de Micco. 1995. Demonstration of *Mycobacterium kansasii* species heterogeneity by the amplification of the 16S-23S spacer region. *J. Med. Microbiol.* **43**:156-158.
2. Avannis-Aghajani, E., K. Jones, A. Holtzman, T. Aronson, N. Glover, M. Boian, S. Froman, and C. F. Brunk. 1996. Molecular technique for rapid identification of mycobacteria. *J. Clin. Microbiol.* **34**:98-102.
3. Böttger, B., T. Rogall, T. Flohr, H. Blöcker, and E. C. Böttger. 1990. Detection and identification of mycobacteria by amplification of rRNA. *J. Clin. Microbiol.* **28**:1751-1759.
4. Bosne, S., and V. Levy-Frebault. 1992. Mycobactin analysis as an aid for the identification of *Mycobacterium fortuitum* and *Mycobacterium chelonae* subspecies. *J. Clin. Microbiol.* **30**:1225-1231.
5. Butler, W. R., K. C. Jost, Jr., and J. O. Kilburn. 1991. Identification of mycobacteria by high-performance liquid chromatography. *J. Clin. Microbiol.* **29**:2468-2472.
6. Devallois, A., E. Legrand, and N. Rastogi. 1996. Evaluation of Amplicor MTB test as adjunct to smears and culture for direct detection of *Mycobacterium tuberculosis* in the French Caribbean. *J. Clin. Microbiol.* **34**:1065-1068.
7. Devallois, A., M. Picardeau, K. S. Goh, C. Sola, V. Vincent, and N. Rastogi. 1996. Comparative evaluation of PCR and commercial DNA probes for detection and identification to species level of *Mycobacterium avium* and *Mycobacterium intracellulare*. *J. Clin. Microbiol.* **34**:2756-2759.
8. Falkinham, J. O. I. 1996. Epidemiology of infection by nontuberculous mycobacteria. *Clin. Microbiol. Rev.* **9**:177-215.
9. Hance, A. J., B. Grandchamp, V. Levy-Frebault, D. Lecossier, J. Rauzier, D. Bocart, and B. Gicquel. 1989. Detection and identification of mycobacteria by amplification of mycobacterial DNA. *Mol. Microbiol.* **3**:843-849.
10. Hughes, M. S., R. A. Skuce, L.-A. Beck, and S. D. Neill. 1993. Identification of mycobacteria from animals by restriction enzyme analysis and direct DNA cycle sequencing of polymerase chain reaction-amplified 16S rRNA gene sequences. *J. Clin. Microbiol.* **31**:3216-3222.
11. Kapur, V., L.-L. Li, M. R. Hamrick, B. B. Plikaytis, T. M. Shinnick, A. Telenti, W. R. Jacobs, A. Banerjee, S. Cole, K. Y. Yuen, J. E. Clarridge, B. N. Kreiswirth, and J. M. Musser. 1995. Rapid *Mycobacterium* species assignment and unambiguous identification of mutations associated with antimicrobial resistance in *Mycobacterium tuberculosis* by automated DNA sequencing. *Arch. Pathol. Lab. Med.* **119**:131-138.
12. Kirschner, P., B. Springer, U. Vogel, A. Meier, A. Wrede, M. Kiekenbeck, F.-C. Bange, and E. C. Böttger. 1993. Genotypic identification of mycobacteria by nucleic acid sequence determination: report of a 2-year experience in a clinical laboratory. *J. Clin. Microbiol.* **31**:2882-2889.
13. Lebrun, L., F. Espinasse, J. Poveda, and V. Vincent-Levy-Frebault. 1992. Evaluation of nonradioactive DNA probes for identification of mycobacteria. *J. Clin. Microbiol.* **30**:2476-2478.
14. Lévy-Frebault, V., M. Daffé, K. S. Goh, M.-A. Lanéelle, C. Asselineau, and H. L. David. 1983. Identification of *Mycobacterium fortuitum* and *Mycobacterium chelonae*. *J. Clin. Microbiol.* **17**:744-752.
15. Mabilat, C., S. Desvarenne, G. Panteix, N. Machabert, M.-H. Bernillon, G. Guardiola, and P. Cros. 1994. Routine identification of *Mycobacterium tuberculosis* complex isolates by automated hybridization. *J. Clin. Microbiol.* **32**:2702-2705.

16. Marks, J., and T. Szulga. 1965. Thin-layer chromatography of mycobacterial lipids as an aid to classification; technical procedures; *Mycobacterium fortuitum*. *Tubercle* **46**:400–411.
17. Picardeau, M., G. Prod'homme, L. Raskine, M. P. LePennec, and V. Vincent. 1997. Genotypic characterization of five subspecies of *Mycobacterium kansasii*. *J. Clin. Microbiol.* **35**:25–32.
18. Plikaytis, B. B., B. D. Plikaytis, M. A. Yakrus, W. R. Butler, C. L. Woodley, V. A. Silcox, and T. M. Shinnick. 1992. Differentiation of slowly growing *Mycobacterium* species, including *Mycobacterium tuberculosis*, by gene amplification and restriction fragment length polymorphism analysis. *J. Clin. Microbiol.* **30**:1815–1822.
19. Ross, B. C., K. Jackson, M. Yang, A. Sievers, and B. Dwyer. 1992. Identification of a genetically distinct subspecies of *Mycobacterium kansasii*. *J. Clin. Microbiol.* **30**:2930–2933.
20. Shinnick, T. M. 1987. The 65-kilodalton antigen of *Mycobacterium tuberculosis*. *J. Bacteriol.* **169**:1080–1088.
21. Shinnick, T. M., M. H. Vodkin, and J. C. Williams. 1988. The *Mycobacterium tuberculosis* 65-kilodalton antigen is a heat shock protein which corresponds to common antigen and to the *Escherichia coli* GroEL protein. *Infect. Immun.* **56**:446–451.
22. Soini, H., E. C. Böttger, and M. K. Viljanen. 1994. Identification of mycobacteria by PCR-based sequence determination of the 32-kilodalton protein gene. *J. Clin. Microbiol.* **32**:2944–2947.
23. Springer, B., L. Stockman, K. Teschner, G. D. Roberts, and E. C. Böttger. 1996. Two-laboratory collaborative study on identification of mycobacteria: molecular versus phenotypic methods. *J. Clin. Microbiol.* **34**:296–303.
24. Takewaki, S.-I., K. Okuzumi, I. Manabe, M. Tanimura, K. Miyamura, K.-I. Nakahara, Y. Yazaki, A. Ohkubo, and R. Nagai. 1994. Nucleotide sequence comparison of the mycobacterial *dnaJ* gene and PCR-restriction fragment length polymorphism analysis for identification of mycobacterial species. *Int. J. Syst. Bacteriol.* **44**:159–166.
25. Taylor, T. B., C. Patterson, Y. Hale, and W. W. Safranek. 1997. Routine use of PCR-restriction fragment length polymorphism analysis for identification of mycobacteria growing in liquid media. *J. Clin. Microbiol.* **35**:79–85.
26. Telenti, A., F. Marchesi, M. Balz, F. Bally, E. C. Böttger, and T. Bodmer. 1993. Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis. *J. Clin. Microbiol.* **31**:175–178.
27. Thorel, M. F., M.-C. Blom-Potar, and N. Rastogi. 1990. Characterization of *Mycobacterium paratuberculosis* and “wood pigeon” mycobacteria by isoenzyme profile and selective staining of immunoprecipitates. *Res. Microbiol.* **141**:551–561.
28. Tortoli, E., M. T. Simonetti, C. Lacchini, V. Penati, and P. Urbano. 1994. Tentative evidence of AIDS-associated biotype of *Mycobacterium kansasii*. *J. Clin. Microbiol.* **32**:1779–1782.
29. Vaneechoutte, M., H. D. Beenhouwer, G. Claeys, G. Verschraegen, A. D. Rouk, N. Paepe, A. Elaichouni, and F. Portaels. 1993. Identification of *Mycobacterium* species by using amplified ribosomal DNA restriction analysis. *J. Clin. Microbiol.* **31**:2061–2065.
30. Vincent Lévy-Frèbault, V., M. F. Thorel, A. Varnerot, and B. Gicquel. 1989. DNA polymorphism in *Mycobacterium paratuberculosis*, “wood pigeon mycobacteria,” and related mycobacteria analyzed by field inversion gel electrophoresis. *J. Clin. Microbiol.* **27**:2823–2826.