



HAL
open science

S100A8 and S100A9 induce cytokine expression and regulate the NLRP3 inflammasome via ROS-dependent activation of NF- κ B(1.).

Jean-Christophe Simard, Annabelle Cesaro, Julie Chapeton-Montes, Mélanie Tardif, Francis Antoine, Denis Girard, Philippe A Tessier

► To cite this version:

Jean-Christophe Simard, Annabelle Cesaro, Julie Chapeton-Montes, Mélanie Tardif, Francis Antoine, et al.. S100A8 and S100A9 induce cytokine expression and regulate the NLRP3 inflammasome via ROS-dependent activation of NF- κ B(1.). PLoS ONE, 2013, 8 (8), pp.e72138. 10.1371/journal.pone.0072138 . pasteur-01132221

HAL Id: pasteur-01132221

<https://riip.hal.science/pasteur-01132221>

Submitted on 16 Mar 2015

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License

S100A8 and S100A9 Induce Cytokine Expression and Regulate the NLRP3 Inflammasome via ROS-Dependent Activation of NF- κ B¹

Jean-Christophe Simard¹, Annabelle Cesaro², Julie Chapeton-Montes², Mélanie Tardif², Francis Antoine¹, Denis Girard¹, Philippe A. Tessier^{2*}

1 Laboratoire de Recherche en Inflammation des Granulocytes, Université du Québec Institut National de la Recherche Scientifique, Institut Armand-Frappier, Laval, Québec, Canada, **2** Axe Maladies Infectieuses et Immunitaires, Centre de Recherche du CHU de Québec and Faculté de Médecine, Université Laval, Québec, Canada

Abstract

S100A8 and S100A9 are cytoplasmic proteins expressed by phagocytes. High concentrations of these proteins have been correlated with various inflammatory conditions, including autoimmune diseases such as rheumatoid arthritis and Crohn's disease, as well as autoinflammatory diseases. In the present study, we examined the effects of S100A8 and S100A9 on the secretion of cytokines and chemokines from PBMCs. S100A8 and S100A9 induced the secretion of cytokines such as IL-6, IL-8, and IL-1 β . This secretion was associated with the activation and translocation of the transcription factor NF- κ B. Inhibition studies using antisense RNA and the pharmacological agent BAY-117082 confirmed the involvement of NF- κ B in IL-6, IL-8, and IL-1 β secretion. S100A8- and S100A9-mediated activation of NF- κ B, the NLR family, pyrin domain-containing 3 (NLRP3) protein, and pro-IL-1 β expression was dependent on the generation of reactive oxygen species. This effect was synergistically enhanced by ATP, a known inflammasome activator. These results suggest that S100A8 and S100A9 enhance the inflammatory response by inducing cytokine secretion of PBMCs.

Citation: Simard J-C, Cesaro A, Chapeton-Montes J, Tardif M, Antoine F, et al. (2013) S100A8 and S100A9 Induce Cytokine Expression and Regulate the NLRP3 Inflammasome via ROS-Dependent Activation of NF- κ B¹. *PLoS ONE* 8(8): e72138. doi:10.1371/journal.pone.0072138

Editor: Emma H. Wilson, University of California, Riverside, United States of America

Received: April 4, 2013; **Accepted:** July 6, 2013; **Published:** August 19, 2013

Copyright: © 2013 Simard et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported by a studentship to JCS from the Fonds de la Recherche en Santé du Québec (FRSQ) and a grant from the Canadian Institutes of Health Research to PAT and DG. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have read the journal's policy and have the following conflicts. This manuscript is related to the following patents: 1. P.A. Tessier, K. Vandal, P. Rouleau C. Ryckman, CHEMOTACTIC FACTOR INHIBITOR FOR MODULATING INFLAMMATORY REACTIONS, European Patent Application EP1519749, filing date 20/06/2003, publication date 06/04/2005, granted 06/03/2009, Licensed to InflammatoRx inc. 2. P.A. Tessier, K. Vandal, P. Rouleau C. Ryckman, ANTIBODIES AGAINST S100A8 AND S100A9 PROTEINS FOR MODULATING INFLAMMATORY REACTIONS, US7,553,488, filing date 15/07/2005, publication date 29/12/2005, granted 30/06/2009, Licensed to InflammatoRx inc. This does not alter the authors' adherence to all the PLOS ONE policies of sharing data and materials.

* E-mail: Philippe.Tessier@crchul.ulaval.ca

Introduction

Endogenous signals originating from stressed, injured, and necrotic cells can activate the innate and adaptive immune systems [1]. These signals have been termed as danger-associated molecular patterns (DAMPs) or alarmins [2]. Like pathogen-associated molecular patterns, DAMPs are recognized by a variety of pathogen-recognition receptors (PRRs) including the toll-like receptors, the formyl-methionine receptors, and the receptor for advanced glycation end-products (RAGE) [1]. Hyaluronan, galectins, high-mobility group box protein, IL-1, and S00A8/A9 are examples of DAMPs [2]. Most DAMPs combine intra- and extra-cellular activities. S100A8 and S100A9 are not exceptions and act as regulators of NADPH oxidase inside the cells and as pro-inflammatory factors once secreted [3,4]. S100A8 and S100A9 are arranged as noncovalently bonded homodimers. In addition, in the presence of calcium, S100A8 and S100A9 form a noncovalent heterodimer called S100A8/A9 or calprotectin, which is presumed to be involved in the cellular control of calcium concentrations. Once released extracellularly, they participate actively in the inflammation process by promoting phagocyte migration [5–7]. S100A8 and S100A9 are presumed to

bind to the PRRs, TLR4, and RAGE, thereby leading to transcription of pro-inflammatory genes [8,9]. S100A9 is also a potent activator of various neutrophil functions, such as degranulation and phagocytosis [10,11], supporting its role as a DAMP.

High extracellular concentrations of S100A8 and S100A9 are found in the serum and at inflammatory sites in autoimmune diseases including arthritis [12–14], lupus, and Crohn's disease [15,16]. In addition, recent studies have demonstrated a role for S100A9 in animal models of lupus, arthritis, and Alzheimer's disease [17–20].

IL-1 β is a well-known cytokine, and its effect on inflammation and in autoinflammatory disorders has been extensively studied [21–23]. In monocytes/macrophages, IL-1 β secretion is predominantly controlled by the NLR family, pyrin domain-containing 3 (NLRP3) inflammasome (a multi-protein complex formed by the protein NLRP3), the adaptor protein called apoptotic speck protein (ASC), and the effector protein caspase-1 [24]. Upon activation of the inflammasome, pro-caspase-1 is recruited into the inflammasome complex and is auto-cleaved into its caspase-recruitment and activation domain, a p10 and a p20 fragment, respectively. Two molecules of the p10 fragment then associate with two p20 fragments to form the active heterotetramer caspase-

1 enzyme that cleaves pro-IL-1 β into IL-1 β and allows its secretion [24].

The inflammasome is formed in cells exposed to microbial, environmental, or endogenous danger stimuli. Two signals are traditionally required for NLRP3 inflammasome activation. The first signal, for example, transmitted by lipopolysaccharide (LPS), induces pro-IL-1 β transcription [25,26]. The second required signal is provided by reduction of intracellular K⁺ [27]. Stimuli such as ATP or the microbial pore-forming toxin nigericin induce strong K⁺ efflux, thus promoting rapid activation of caspase-1 and secretion of mature IL-1 β [28].

We recently reported that S100A9 induces the secretion of cytokines/chemokines in human monocytes [17]. We herein report that S100A8 and S100A9 induce the secretion of several pro-inflammatory cytokines, including IL-6, TNF α , and IL-1 β , by stimulating production of reactive oxygen species (ROS). These, in turn, activate the transcription factor NF- κ B, leading to cytokine secretion and expression and activation of the NLRP3 inflammasome.

Materials and Methods

Ethics Statement

These studies were approved by the INRS-Institut Armand-Frappier's ethical committee, and all participants gave written informed consent.

Reagents

Human recombinant S100A8 and S100A9 were produced as previously described [29] and found were to contain less than 1 pg endotoxin/mg of protein. LPS (from *E. coli*), cycloheximide, diphenyleneiodonium chloride (DPI), N-acetylcysteine (NAC), and anti- β -actin were purchased from Sigma-Aldrich (St. Louis, MO). BAY-117082 (BAY) was purchased from Calbiochem (San Diego, CA). Anti-phospho-inhibitory kappa B kinase (IKK)- α / β (Ser176/177), anti-phospho-IKK- γ (Ser376), and anti-caspase-1 were purchased from Cell Signaling Technology (Danvers, MA). Antibodies against IkB- α (H-4), GAPDH (FL), NF- κ B p50 (4D1), NF- κ B p65 (C-20), ASC and pro-IL-1 β (H-153) were obtained from Santa Cruz Biotechnology (Santa Cruz, CA). The NLRP3/NALP-3 antibody was purchased from Enzo Life Sciences (Farmingdale, NY). RPMI 1640, HEPES, penicillin/streptomycin (pen/strep), heat-inactivated FBS, opti-MEM media, and HBSS were purchased from Life Technologies (Grand Island, NY). Ficoll-Paque was obtained from GE Healthcare Bio-Science AB (Uppsala, Sweden). All secondary antibodies were purchased from Jackson ImmunoResearch Laboratories (West Grove, PA). The Proteome Profiler™ Array (Human Cytokine Array Panel A) was purchased from R&D Systems (Minneapolis, MN). THP-Blue™, QuantiBlue™, and standard LPS 0111:B4 were purchased from InvivoGen (San Diego, CA). Protease Inhibitor Cocktail was purchased from Thermo Fisher Canada (Nepean, Ontario). PureProteome Protein A Magnetic Beads were obtained from Millipore (Billerica, MA).

PBMC Isolation

Blood samples were obtained from informed, consenting individuals according to institutionally approved procedures. PBMCs were isolated from venous blood of healthy volunteers using dextran sedimentation, followed by centrifugation over a Ficoll-Paque cushion. PBMCs were harvested and washed after centrifugation. Cell viability was monitored by trypan blue exclusion and consistently found to be greater than 99%. PBMCs

were resuspended at 1×10^6 cells/ml in RPMI 1640 containing 10% autologous decompartmented serum.

Western Blot Analyses

PBMCs were stimulated with S100A8 (10 μ g/ml), S100A9 (10 μ g/ml), LPS (1 μ g/ml), or the equivalent volume of diluent (HBSS, 1 \times) at 37°C for various periods of time, as specified. For detection of IkB- α , cells were pre-treated with 5 μ M cycloheximide to limit rapid turnover of the protein. At the end of the incubation period, cells were lysed in Laemmli's sample buffer (0.25 M Tris-HCl [pH 6.8], 8% SDS, 40% glycerol, and 20% 2-ME), and aliquots of extracts (corresponding to 8×10^5 cells) were subjected to 10% SDS-PAGE and transferred to nitrocellulose membranes for detection of specific proteins. Membranes were blocked for 1 h at room temperature in 5% milk or in B10k-PO Reagent from Millipore for phosphorylated proteins. After washing, primary antibodies were added at a final dilution of 1:1000 in 0.15% TBS-Tween. The membranes were incubated overnight at 4°C, washed with TBS-Tween, and incubated for 1 h at room temperature with a goat anti-rabbit IgG HRP secondary Ab (Jackson ImmunoResearch Laboratories) diluted 1:25,000 in TBS-Tween or a goat anti-mouse IgG HRP secondary Ab (Jackson ImmunoResearch Laboratories) diluted at 1:25,000 in TBS-Tween. After several washes, protein expression was evaluated using the Luminata Forte Western HRP Substrate (Millipore). Membranes were stripped with ReBlot Plus Strong (Millipore) and reprobed to confirm equal loading of proteins.

Immunoprecipitation

PBMCs (6×10^6 cells) were stimulated for 4 hours with the indicated agonists followed by 30 min with 1 mM ATP. Cells were then centrifuged and lysed in ice-cold lysis buffer 1X (50 mM Tris-HCl pH 8, 100 mM NaCl, 1% Triton X-100, 5 mM EDTA and Protease Inhibitor Cocktail 1X) for 1 hour on ice and sonicated 3 times for 20 s. The lysates were first incubated 1 h on ice with magnetic beads (5 μ l per 10^6 cells) for 1 h with agitation to remove non-specific binding. The beads were then removed and the samples were incubated overnight with 5 μ g of rabbit anti-ASC at 4°C with agitation. Protein A magnetic beads were then added (5 μ l per 1×10^6 cells) for 30 min at R.T. with agitation. The complex (antibody, bead, protein) was collected and 50 μ L of Laemmli's sample buffer 1X was added. The samples were then boiled at 100°C for 10 min, and the immunoprecipitates were then loaded on a SDS-page gel for western blot analysis.

EMSA

Nuclear extracts from cells stimulated for 4 h were prepared using the NucBuster protein extraction kit from EMD Millipore. The EMSA reaction was performed using the Gel Shift Assay System from Promega (Madison, WI). Briefly, 7 μ g of nuclear protein was mixed with 1.75 pmol of unlabeled NF- κ B consensus oligonucleotides (competitive assay) or unlabeled AP-1 consensus oligonucleotides (non-competitive assay) for 15 min at room temperature. Next, 0.03 pmol of (³²P)-labeled NF- κ B consensus oligonucleotides were added for 30 min at room temperature. The DNA-protein complex was resolved on 4% non-denaturing polyacrylamide gels. Gels were exposed for 24 h at -80°C, and the bands were visualized using X-OMAT films.

NF- κ B p50 TransFactor ELISA Assay

Assays were performed using the Colorimetric TransFactor ELISA kit (Clontech Laboratories, Mountain View, CA) according to the manufacturer's protocol. Briefly, nuclear extracts were

prepared as described above. Nuclear proteins (25 μg) were incubated with or without competitor oligonucleotides for 15 min on ice. Competitor oligonucleotides correspond to the same DNA sequence as that of the oligo-coated wells, showing the binding specificity between DNA and the transcription factor. Samples were then added to wells for 60 min at room temperature. After four washes, primary antibody was added to each well and incubated for 60 min at room temperature. After four washes, a goat-anti-rabbit IgG-HRP secondary antibody was added to each well and incubated for 30 min at room temperature. After four washes, the tetramethylbenzidine (TMB) substrate was added to each well for 10 min at room temperature. Finally, 1 M H_2SO_4 was added to each well, and absorbance was read at 450 nm using a SpectraMax M5 spectrophotometer (Molecular Devices, Sunnyvale, CA).

Antisense Experiments

PBMCs ($1 \times 10^6/\text{ml}$) were incubated with p50 (5'-tggatctctgc-cattct-3'), p65 (5'-ggggaacagtgctccatggc-3'), or scrambled (5'-ttaccgcgctagacgggca-3') antisense oligonucleotides (AS) with a phosphorothioate backbone at a final concentration of 10 μM or 40 μM at 37°C in serum-free Opti-MEM media (Invitrogen) for 4 h to increase oligonucleotide uptake. Subsequently, cells were stimulated with agonists for 20 h in Opti-MEM media supplemented with 5% FBS. The transfection efficiency was greater than 90%, as determined using the same oligonucleotide sequences with a Cy3 fluorophore coupled to the 5' end and visualized by fluorescence microscopy. p50 and p65 protein expression was monitored by western blot analysis.

Immunofluorescence Microscopy

As described previously [30], following the incubation of Cy5-AS, cells were washed twice in ice-cold PBS and cytocentrifuged on glass coverslips (Fisher Scientific, Ottawa, Canada). Coverslips were mounted using ProLong Gold Antifade Media containing DAPI (Invitrogen). Fluorescent-labeled cells were captured from high-power field (3400) and observed using a Leica microscope equipped with an ebq 100 dc epifluorescent condenser.

Proteome Profiler™ Array

The Human Cytokine Array Panel A (Proteome Profiler™ Array) was purchased from R&D Systems. All steps were performed within 3 weeks of harvesting the cells. Cells from 10 different PBMC donors were stimulated for 24 h with the corresponding agonists. Supernatants were pooled together and used to probe the membranes. The chemiluminescent signals from the bound cytokines present in the supernatants were detected on X-OMAT films. Values obtained from densitometry analyses were considered when above 9000 units, which corresponds to the background value.

Caspase-1 Assay

Caspase-1 activity was measured using the Caspase-1 Colorimetric Assay kit (R&D Systems) according to the recommended protocol. Briefly, 2×10^6 cells were stimulated with agonists for the indicated time period. Cells were lysed in ice-cold lysis buffer and incubated on ice for 10 min. Cells were centrifuged at 10,000 g for 1 min. Supernatants were then used for enzymatic reactions. Caspase-1 colorimetric substrate (WEHD-pNA) from the kit was added to each reaction mix with the corresponding volume of cell lysate, reaction buffer, and DTT. Plates were incubated for 2 h at 37°C. After incubation, plates were read on a microplate reader at

a wavelength of 405 nm. Results are expressed as fold-increase in caspase-1 activity.

IL-6, IL-8, and IL-1 β Production

IL-6, IL-8, and IL-1 β levels were measured using commercially available ELISA kits (Invitrogen). Freshly isolated cells were incubated with S100A8 (10 $\mu\text{g}/\text{ml}$), S100A9 (10 $\mu\text{g}/\text{ml}$), LPS (1 $\mu\text{g}/\text{ml}$), or the diluent (HBSS) for the indicated time period in 96-well plates containing RPMI-1640 supplemented with 10% autologous serum. Supernatants were harvested after centrifugation and stored at -80°C before ELISA.

Measurement of NF- κB Activity Using THP1-XBlue™ Cells

THP1-XBlue™ cells (1×10^5) were pretreated with DPI (10 μM) or vehicle (DMSO) for 1 h and then stimulated with S100A8 (10 $\mu\text{g}/\text{ml}$), S100A9 (10 $\mu\text{g}/\text{ml}$), or LPS (1 $\mu\text{g}/\text{ml}$) for 24 h. Supernatants were harvested and incubated with QuantiBlue™, which turns purple in the presence of secreted alkaline phosphatase (SEAP). SEAP levels were determined spectrophotometrically at 650 nm.

siRNA Transfection

PBMCs were transfected with Silencer Select Pre-designed siRNA from Ambion (Austin, TX) according to the manufacturer's protocol. Briefly, cells were suspended in 500 μL of opti-MEM reduced serum medium containing 3 μL of Lipofectamine RNAiMAX Reagent (Life Technologies) and various concentration of siRNA. siRNA targeting ASC (sense UGAU-CUUUUUAUACACAAUTT, antisense AUUGU-GUAUAAAAAGAUCAGA), NLRP-3 (sense GGAGAGACCUUUAUGAGAATT, antisense UUCU-CAUAAGGUCUCUCCTG) or Control non-silencing siRNA (sense UUCUCCGAACGUGUCACGUTT, antisense ACGU-GACACGUUCGGAGGAGAATT) were used. After 24 hours, cells were washed and stimulated with the indicated agonists.

Statistical Analyses

Experimental data are expressed as means \pm SEM. One-way ANOVA (Dunnett's multiple-comparison test) and two-way ANOVA (Bonferroni's post-test) were performed using Graph-Pad Prism (version 5.01). Differences were considered statistically significant as follows: * $p \leq 0.05$, ** $p \leq 0.01$, and *** $p \leq 0.005$ versus buffer or the appropriate diluent. Densitometric analyses were performed using AlphaEaseFC (FluorChem HD2).

Results

S100A8 and S100A9 Induce Inflammatory Cytokine Secretion

The effect of S100A8 and S100A9 on the production of cytokines/chemokines was first screened using an antibody array. As illustrated in Figure 1A and 1B, S100A8 and S100A9 induced the secretion of several pro-inflammatory cytokines in PBMCs (Fig. 1A and 1B). Among these, the cytokines IL-6, IL-1 β , TNF- α and the chemokines IL-8, growth-regulated oncogene alpha (Gro- α), MIP-1 α , and MIP-1 β were the most secreted. The secretion of IL-6 and IL-8 in PBMCs was also confirmed by ELISA (Fig. 1C). The concentrations of these cytokines were comparable to those observed in the supernatants of cells stimulated with LPS. Isolation of monocytes, CD4+ T cells, CD8+ T cells, and B cells revealed that monocytes were the only cell type responsible for cytokine production (data not shown).




Figure 1. S100A8 and S100A9 induce the secretion of several cytokines in human PBMCs. Cytokine production was evaluated using proteome profiler arrays (A and B) or ELISA (C), as described in Materials and Methods. Cells were stimulated for 24 hours, and supernatants were harvested and used for experiments. (B) Densitometry analyses of antibody arrays in PBMCs. Data represent supernatants pooled from 10 different donors (A and B). Black bars, upregulated cytokines; white bars, unchanged cytokines. Data are the mean \pm SEM of at least five experiments performed on cells from different donors (C). *** $p \leq 0.005$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g001

S100A8 and S100A9 Increase NF- κ B Translocation and Activation

The production of pro-inflammatory cytokines and chemokines largely depends on transcription factors such as NF- κ B and AP-1. Activation of NF- κ B in S100A8- and S100A9-treated PBMCs was therefore examined using EMSA. As shown in Figure 2, after stimulation for 4 h, S100A8 and S100A9 increased the DNA-binding activity of NF- κ B in PBMCs compared with that in untreated cells. As expected, LPS, which was used as a positive control, also increased this activity. These results were confirmed using an NF- κ B p50 TransFactor ELISA kit (Clontech Laboratories), which allows direct measurement of the p50 subunit levels in the nucleus (Fig. 2B). NF- κ B translocates into the nucleus after degradation of its inhibitor I κ B- α , which occurs upon phosphorylation of the p50 subunit by the IKK complex [31]. Phosphorylation of the IKK complex and degradation of I κ B- α in S100A8- and S100A9-stimulated cells was therefore investigated. Both proteins induced phosphorylation of IKK α / β / γ and degradation of I κ B- α in PBMCs (Fig. 2C), which was comparable to the effect induced by LPS. These results were consistent with the translocation of NF- κ B induced by S100A8 and S100A9 in PBMCs.

S100A8- and S100A9-mediated IL-6 and IL-8 Secretion Partially Depends on NF- κ B Activation

Next, we examined the role of NF- κ B activation in S100A8- and S100A9-induced cytokine secretion. First, an antisense strategy was used to down-regulate p50 or p65 (Rel A) expression. As shown in Figure 3, p50 and p65 expression remained similar to that in control cells when scrambled antisense was added to the

culture media. However, p50 antisense markedly decreased the expression of p50 ($\sim 30\%$), whereas p65 antisense inhibited p65 less potently ($\sim 15\%$). These results confirm that the p50 and p65 oligonucleotides were specific. Moreover, all antisense oligonucleotides were able to enter the cells at the same efficiency, as assessed by fluorescence microscopy. Decreased p50 and p65 expression resulted in reduced IL-8 secretion, confirming the importance of NF- κ B for the activity of S100A8 and S100A9 (Fig. 4). Interestingly, inhibition of p50 and p65 did not significantly diminish IL-6 production. The I κ B- α specific inhibitor BAY-117082 was also used to confirm these results. Addition of BAY-117082 almost completely inhibited IL-8 secretion, but IL-6 production was less affected.

Translocation of NF- κ B in Response to S100A8 and S100A9 Depends on ROS Production

ROS actively participate as the second messenger in the induction of several genes in numerous physiological and pathological conditions [reviewed in [32–34]], and function, at least in part, through the activation of NF- κ B [35,36]. Because S100A8 and S100A9 trigger ROS production in other myeloid cells, including human neutrophils [11], the role of ROS in S100A8 and S100A9-induced NF- κ B activation was investigated. First, the production of ROS in monocytes was measured using the dichlorodihydrofluorescein diacetate (H₂DCFDA) probe. S100A8 and S100A9 induced a slow accumulation of ROS over time (Fig. 5A), with ROS production becoming detectable after 90 min of stimulation with S100A9 and after 120 min of stimulation with S100A8. Next, we used the flavoprotein inhibitor DPI, which inhibits ROS generation (Fig. 5B) from NADPH




Figure 2. S100A8 and S100A9 increase translocation of NF- κ B in human PBMCs. NF- κ B translocation and binding were assessed by EMSA (A) and by a NF- κ B p50 TransFactor ELISA kit (B). Nuclear proteins were isolated from cells stimulated for 4 h. Lanes 1, 3, 5, and 7 (c, competitive assay). Lanes 2, 4, 6, and 8 (non-competitive assay). Phosphorylation of the IKK complex and degradation of I κ B- α were assessed by immunoblotting (C). Results shown are of one experiment that is representative of at least four other experiments (A and C). Data are the mean \pm SEM of three experiments performed on cells from different donors (B). * $p \leq 0.05$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g002

oxidases and mitochondria, and monitored the production of IL-8 and IL-6, as well as the activation of NF- κ B. DPI was found to block the production of IL-8 (Fig. 5C) and IL-6 (Fig. 5D) in a dose-dependent manner. Translocation of NF- κ B was evaluated by measuring the p50 subunit levels in the nucleus in the presence of 10 μ M DPI. DPI significantly reduced the p50 subunit levels in

the nucleus of cells stimulated with S100A9 or LPS (Fig. 5E). A similar trend was observed when cells were exposed to S100A8. To confirm the negative role of DPI in S100A8- and S100A9-mediated NF- κ B activation, the THP1-XBlueTM cell line reporting NF- κ B activity was used. As expected, DPI strongly inhibited the activation of NF- κ B in S100A8-, S100A9-, and LPS-treated




Figure 3. Inhibition of p50 and p65 using an antisense (AS) strategy in human PBMCs. Cells were treated for 4 h with p50 AS (10 or 40 μ M), p65 AS (10 or 40 μ M), the scrambled sequence AS (Scr 40 μ M), or the equivalent volume of diluent (No). Cells were then fixed on coverslips for fluorescence microscopy (A) or lysed for immunoblotting of p50 (B) and p65 (C). Results are from one experiment out of 4 total experiments are shown. * $p \leq 0.05$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g003




Figure 4. S100A8 and S100A9 induce IL-6 and IL-8 secretion via activation of NF- κ B in human PBMCs. Secretion of IL-6 (A and B) and IL-8 (C and D) was evaluated using ELISA. Cells were pre-treated with 1–10 μ M of BAY-117082 (or the equivalent amount of diluent) for 30 min or with antisense (Scr, p50, p65) for 4 h, followed by incubation with the indicated agonist for a total of 24 h. Supernatants were harvested and used for ELISA. Data are the mean \pm SEM of at least four experiments performed on cells from different donors. * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.005$, Dunnett's multiple comparison test.

doi:10.1371/journal.pone.0072138.g004

cells (Fig. 5F). Again, S100A9 was found to be a more potent activator than S100A8.

S100A8 and S100A9 Regulate the NLRP3 Inflammasome

Caspase-1 processing is the ultimate step in the activation of the NLRP3 inflammasome and leads to the secretion of IL-1 β , one of the cytokines up-regulated by S100A8 and S100A9 in human PBMCs (Fig. 1). The NLRP3 inflammasome is tightly controlled by a priming step that is dependent on NF- κ B [37]. Because S100A8 and S100A9 induced ROS production that resulted in the activation of NF- κ B, the roles of these proteins in NLRP3 inflammasome regulation and activation were investigated. The expression of pro-IL-1 β and NLRP3 was first analyzed by western blot after 4-h stimulation with S100A8, S100A9, or LPS. Both S100A8 and S100A9 increased pro-IL-1 β levels, albeit more modestly than LPS (Fig. 6A). However, NLRP3 expression was greatly increased following stimulation with S100A8 and S100A9 to levels comparable to those achieved by LPS stimulation (Fig. 6B). Pretreatment of cells with S100A8 or S100A9 followed by stimulation with ATP, a known inflammasome activator, increased the association between ASC and NLRP-3 compared to control cells without affecting the expression of the adaptor protein ASC (Fig. 6C and D). Next, the processing of caspase-1 into the active 10-kDa and 20-kDa subunits was monitored in the presence or absence of a second signal mediated by ATP. Cell priming with S100A8 or S100A9 alone was not sufficient to induce a strong cleavage of caspase-1 (Fig. 6E). However, caspase-1 cleavage was synergistically enhanced by the addition of ATP. Caspase-1 assays

were performed to further confirm the ability of S100A8 and S100A9 to modulate the enzymatic activity of caspase-1. As expected, the catalytic activity of caspase-1 was enhanced by ATP and potentiated by the presence of S100A8, and to a greater extent, by S100A9 (Fig. 6F). Next, the ability of S100A8 and S100A9 to induce IL-1 β secretion in the presence or absence of ATP was examined using ELISA. Interestingly, the levels of extracellular IL-1 β released by monocytes exposed to ATP+S100A9 were similar to those in cells exposed to ATP+LPS; however, the priming with S100A8 was again less potent (Fig. 6G).

NLRP-3 and ASC are Essentials for IL-1 β Secretion in S100A8 and S100A9-treated Cells

Secretion of IL-1 β is known to be regulated, at least in part, through the NLRP-3 inflammasome and its adaptor protein ASC [24,25]. Therefore, the roles of NLRP-3 and ASC in the production of IL-1 β in S100A8 and S100A9-primed cells were investigated. Addition of ASC-specific siRNA led to a 90% depletion of ASC (Fig. 7A) while siRNA against NLRP-3 reduced its expression by about 60% (Fig. 7B). Nevertheless, both silencing markedly reduced the secretion of IL-1 β in S100A8 and S100A9-primed cells.

ROS-mediated NF- κ B Activation is Involved in NLRP3 Regulation by S100A8 and S100A9

NLRP3 activation is intimately linked to ROS generation [32]. Given that stimulation with S100A8 and S100A9 generated




Figure 5. NF- κ B translocation and IL-8 production in S100A8- or S100A9-treated cells are dependent on ROS production. Cells were stimulated for different time periods and ROS production was assessed by oxidation of H₂DCFDA (A and B). In some experiments, cells were pre-treated with DPI for 30 min (B-F). ROS production was measured after 180 min (B). IL-6 and IL-8 secretion was assessed using ELISA (C and D). Translocation of NF- κ B was assessed using the TransFactor p50 kit (E). NF- κ B activity was followed with THP1-XBlue cells (F). Data are the mean \pm SEM of three experiments performed on cells from four different donors (A-E). Data are the mean \pm SEM of three independent experiments (F). * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.005$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g005

intracellular ROS in monocytes and promoted NF- κ B activity, we investigated the possibility that oxidative stress is involved in S100A8- and S100A9-mediated NLRP3 inflammasome regulation and activation. DPI and the anti-oxidant NAC, a well-known ROS scavenger, were used to abrogate the effects of the oxidative stress induced by S100A8 and S100A9. DPI slightly reduced the expression of NLRP3 and pro-IL-1 β in S100A8, S100A9-, and LPS-primed cells. However, NAC efficiently inhibited the expression of both NLRP3 and pro-IL-1 β (Fig. 8A). The inhibition was to a similar extent when cells were treated with the NF- κ B inhibitor BAY-117082, suggesting that oxidative stress is important in the observed phenomenon (ASC expression levels remained constant, Fig. 8C). DPI and NAC strongly inhibited caspase-1 processing, whereas BAY-117082 had little impact on cells stimulated with S100A8, S100A9, or LPS in the presence of ATP (Fig. 8B). These results were corroborated by analyzing IL-1 β release into the supernatant from the same samples. As illustrated in Figure 8D, NAC strongly impaired the secretion of IL-1 β induced by S100A9 and LPS, and to a lesser extent, by S100A8. Again, NAC was a more potent inhibitor of IL-1 β production than either DPI or BAY-117082.

Discussion

In the last few years, studies have classified S100A8 and S100A9 as DAMPs. S100A8 and S100A9 were shown to be actively involved in the inflammatory response by regulating neutrophil functions [5–7,10,11,38]. In this study, we further explored the roles of these two proteins by investigating cytokine production in human PBMCs. Both S100A8 and S100A9 induced the secretion

of various pro-inflammatory cytokines in monocytes, including IL-8, IL-6, and IL-1 β , through mechanisms dependent on ROS production and the activation of ROS-sensitive transcription factors, such as NF- κ B. This study further demonstrated that S100A9, and to a lesser extent, S100A8, regulate the NLRP3 inflammasome by acting as priming agents. IL-1 β secretion mediated by S100A8 or S100A9 alone was modest, but was greatly increased following caspase-1 activation via ATP.

The ability of S100A8 and S100A9 to promote expression of pro-inflammatory mediators including TNF- α , IL-1 β , IL-6, and IL-8 through NF- κ B activation has been previously reported in the cell line THP-1 and human macrophages [39,40]. As in the present study, S100A9 more potently induced cytokine production compared with S100A8. This might reflect the use of different receptors, or a different affinity for their putative receptors TLR4 and/or RAGE [9,41,42]. However, the mechanisms by which these proteins increased NF- κ B-mediated expression of pro-inflammatory cytokines remained unclear. Here, we report up-regulation of several other cytokines and chemokines, such as MIP-1 α , MIP-1 β , Gro- α , IL-23, G-CSF, and GM-CSF, in addition to TNF- α , IL-1 β , IL-6, and IL-8. The expression of most of these cytokines depends on NF- κ B activation, but other transcription factors could also be involved. This is highlighted by the fact that IL-6 secretion in response to S100A8 and S100A9 is less affected by NF- κ B inhibition compared to IL-8 and IL-1 β secretion, suggesting the involvement of other transcription factors in the secretion of IL-6. Indeed, multiple transcription factor-binding sites were identified in the human IL-6 promoter, including AP-1 [43–45]. AP-1 is intimately linked with the activation of mitogen-activated protein kinases (MAPKs) such as




Figure 6. S100A8 and S100A9 regulate the NLRP3 inflammasome. Cells were incubated for 4 h with S100A8 (10 μ g/ml), S100A9 (10 μ g/ml), LPS (1 μ g/ml), or the equivalent volume of buffer (Ctrl). For some experiments, ATP (1 mM) was added for 30 min (D–G). Cells were harvested and lysed with Laemmli's buffer for immunoblotting of pro-IL-1 β (A), NLRP3 (B), ASC (C) or caspase-1 (E). Cells were lysed with caspase-1 lysis buffer for the caspase assay (F). ASC immunoprecipitates were loaded on SDS-PAGE for NLRP-3 and ASC western blot analysis (D). Supernatants were harvested before performing IL-1 β ELISA (G). Results are from one experiment that is representative of at least three others (A–E). Data are the mean \pm SEM of three experiments performed on cells from different donors (F–G). ?, non-specific band (as indicated in the manufacturer's datasheet). * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.005$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g006

p38 and extracellular signal-regulated kinase (ERK1/2) [46]. We and others have previously demonstrated that S100A9 induced phosphorylation of p38, ERK1/2, Syk, and AKT in monocytes and neutrophils [10,11,40]. Interestingly, most of these kinases are regulated by redox-sensitive phosphatases such as tyrosine phosphatases (PTPs) and phosphatase and tensin homologs deleted on chromosome 10 (PTENs) [33,47]. As previously observed in neutrophils, S100A9, and to a lesser extent, S100A8, induce ROS generation in monocytes that could potentially inactivate redox-sensitive phosphatases and thereby sustain cell activation.

ROS act as important second messengers and participate in numerous cellular functions through the regulation of redox-sensitive transcription factors, including NF- κ B and AP-1 [33,35,48]. Our results show that the expression of IL-6 and IL-8 in monocytes primed with S100A8 or S100A9 depends largely on ROS generation, because DPI significantly impaired the expression of both cytokines. Similar effects were observed for IL-1 β , the secretion of which is tightly regulated by caspase-1 activation. Signals triggering formation and activation of the inflammasome involve decreased intracellular K⁺ and the generation of oxidative stress [27,49,50]. These signals are typically provided by the engagement of toll-like receptors (TLRs) or the presence of endogenous stresses that modulate the ionic milieu and redox balance. Various inflammasomes have been identified in monocytes/macrophages such as the NLRP1, NLRP4, and NLRP3, which remains the most studied [51]. All

of them converge toward caspase-1 activation and IL-1 β secretion. The exact mechanisms responsible for the production and secretion of IL-1 β remain unclear, but require two main steps. The first step involves the transcription and expression of proteins and/or precursors including pro-IL-1 β and inflammasome subunits. The second step is inflammasome-mediated caspase-1 activation by stimuli such as ATP that generate robust K⁺ release [24,28]. Tight control and regulation of the inflammasome is required to avoid excessive cytokine secretion. Therefore, regulation of the NLRP3 inflammasome occurs at the transcriptional and post-transcriptional levels. For instance, expression of NLRP3 requires a signal to be highly induced [52]. Interestingly, TLR signaling molecules are potent regulators of NLRP3 expression [37,53]. In fact, efficient priming of cells by TLR signaling molecules is well documented, showing the importance of the priming step for efficient inflammasome activation [26,54]. Our results demonstrate the importance of S100A8 and S100A9 in the regulation of the NLRP3 inflammasome, because these molecules upregulate NLRP3 and pro-IL-1 β expression.

Various microbial-derived products can activate the NLRP3 inflammasome. Moreover, mitochondrial stresses, and host-derived molecules abundantly found in some metabolic and inflammatory diseases, such as excess ATP levels, oxidized low-density lipoproteins (LDLs), excess glucose, cholesterol crystals, monosodium uric acid (MSU), and high-mobility group box 1 (HMGB1) protein are also involved in NLRP3 inflammasome




Figure 7. Role of ASC and NLRP-3 in the production of IL-1 β by S100A8 and S100A9-treated human PBMCs. Cells were transfected with ASC siRNA, NLRP-3 siRNA, Ctrl siRNA or the equivalent volume of buffer (No) for 24 hours. Cells were then harvested and lysed for western blot analysis (A and B) or stimulated for 4 hours with the indicated agonists and another 30 min with ATP 1mM. Supernatants were then harvested before performing ELISA (C). Results are from one experiment representative of three others (A and B). Data are the mean \pm SEM of four experiments performed on cells from different donors (C). * $p \leq 0.05$, ** $p \leq 0.01$, *** $p \leq 0.005$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g007

activation [55–58]. Here, we demonstrate that extracellular S100A8 and S100A9 are potent regulators of the NLRP3 inflammasome, and they act by influencing the redox balance in human monocytes. The mechanisms by which these proteins enhance ROS production are still unknown. However, considering that S100A8 and S100A9 partially mediate their effects through TLR4 activation [8,40], we propose that intracellular signals induced by S100A8 and S100A9 are likely to be similar to LPS, which is known to trigger ROS production crucial for the expression of inflammatory cytokines through the activation of redox-sensitive transcription factors.

Release of mature IL-1 β is greatly potentiated by priming of the NLRP3 inflammasome by NF- κ B-driven protein synthesis stimulated by LPS [26]. However, long-term stimulation of monocytes in the absence of a co-signal, such as ATP, also leads to a slow activation of caspase-1 and liberation of IL-1 β [59–61]. Our results show that S100A8 and S100A9 alone induce IL-1 β secretion after long-term stimulation (24 h). This secretion was greatly enhanced by the addition of ATP. In fact, only 4 h of priming with S100A8 and S100A9 and 30 min of stimulation with ATP were sufficient to observe caspase-1 cleavage and IL-1 β release. Uncontrolled activation of the inflammasome has been previously linked to several immune and non-immune diseases [58,62]. For example, dextran sodium sulfate-mediated colitis and MSU-induced joint inflammation in a murine model of gout are less severe in mice deficient in caspase-1 or NLRP3 [63–65]. Similarly, mice deficient for the NLRP3 inflammasome adaptor protein, ASC, are protected from chronic obesity-induced pancreatic damage [66]. Interestingly, S100A8 and S100A9 concentrations have been correlated to disease activity in inflammatory bowel disease [15,16,67–69], gout [29,70], obesity-mediated type 2 diabetes, and atherosclerosis [reviewed in [71]],

suggesting a possible link between S100A8, S100A9, and exacerbated production of IL-1 β .

Patients with hypercalprotectinemia have blood concentrations of S100A8/A9 that are 1000 to 10,000 times higher than the normal concentrations. These patients suffer from chronic systemic inflammation, presumably due to dysregulation of the S100A8/A9 metabolism [72]. This hereditary syndrome results from a mutation in the gene PSTPIP-1, which increases the binding of S100A8/A9 to PSTPIP-1. Interestingly, high concentrations of cytokines like IL-1 β , IL-8, and TNF α were found in the serum of a patient with hypercalprotectinemia (unpublished observation). These cytokines are also detected in the supernatant of PBMCs stimulated with S100A8 and S100A9, suggesting that S100A8 and S100A9 could be responsible for dysregulation of cytokine production in hypercalprotectinemia patients. The higher concentrations of IL-1 β detected in the serum of the hypercalprotectinemia patient also support the potential effects of S100A8 and S100A9 on NLRP3 inflammasome priming. This effect is likely not restricted to hypercalprotectinemia, because several cytokines actively involved in arthritis [73], including IL-6 and TNF α , were found to be upregulated by S100A8 and S100A9. In addition, blocking S100A9 resulted in diminished TNF α and IL-6 expression in a collagen-induced arthritis model [17].

In conclusion, the results presented in this study suggest a new role for S100A8 and S100A9 in redox-sensitive biological responses, including pro-inflammatory cytokine secretion. These findings also suggest a role of S100A8 and S100A9 in the pathogenesis of various inflammatory diseases.




Figure 8. ROS- and NF- κ B-mediated NLRP3 inflammasome priming by S100A8 and S100A9 in human PBMCs. Cells were pre-treated with various inhibitors (DPI, NAC, BAY-117082 or the equivalent volume of DMSO), then stimulated for 4 h with the indicated agonists. ATP (1 mM) was added for 30 min for caspase-1 immunoblots and for IL-1 β ELISA (B and D). Cells were harvested and lysed in Laemmli's buffer for immunoblotting of NLRP-3, pro-IL-1 β , ASC and caspase-1.(A-C). Supernatants were harvested and IL-1 β secretion was evaluated using ELISA (C). * $p \leq 0.05$, *** $p \leq 0.005$, Dunnett's multiple comparison test. doi:10.1371/journal.pone.0072138.g008

Acknowledgments

We would like to thank Nathalie Pagé and Audrey Plante for the production of recombinant S100A8 and S100A9.

References

- Matzinger P (1994) Tolerance, danger, and the extended family. *Annu Rev Immunol* 12: 991–1045.
- Bianchi ME (2007) DAMPs, PAMPs and alarmins: all we need to know about danger. *J Leukoc Biol* 81: 1–5.
- Kessel C, Holzinger D, Foell D (2012) Phagocyte-derived S100 proteins in autoinflammation: Putative role in pathogenesis and usefulness as biomarkers. *Clin Immunol*.
- Lim SY, Raftery MJ, Goyette J, Hsu K, Geczy CL (2009) Oxidative modifications of S100 proteins: functional regulation by redox. *J Leukoc Biol* 86: 577–587.
- Anceriz N, Vandal K, Tessier PA (2007) S100A9 mediates neutrophil adhesion to fibronectin through activation of beta2 integrins. *Biochem Biophys Res Commun* 354: 84–89.

Author Contributions

Conceived and designed the experiments: JCS AC, JCM MT FA DG PAT. Performed the experiments: JCS AC, JCM MT FA DG PAT. Analyzed the data: JCS AC, JCM MT FA DG PAT. Wrote the paper: JCS AC, JCM MT FA DG PAT.

6. Ryckman C, Vandal K, Rouleau P, Talbot M, Tessier PA (2003) Proinflammatory activities of S100: proteins S100A8, S100A9, and S100A8/A9 induce neutrophil chemotaxis and adhesion. *J Immunol* 170: 3233–3242.
7. Vandal K, Rouleau P, Boivin A, Ryckman C, Talbot M, et al. (2003) Blockade of S100A8 and S100A9 suppresses neutrophil migration in response to lipopolysaccharide. *J Immunol* 171: 2602–2609.
8. Vogl T, Tenbrock K, Ludwig S, Leukert N, Ehrhardt C, et al. (2007) Mrp8 and Mrp14 are endogenous activators of Toll-like receptor 4, promoting lethal, endotoxin-induced shock. *Nat Med* 13: 1042–1049.
9. Hofmann MA, Drury S, Fu C, Qu W, Taguchi A, et al. (1999) RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell* 97: 889–901.
10. Simard JC, Girard D, Tessier PA (2010) Induction of neutrophil degranulation by S100A9 via a MAPK-dependent mechanism. *J Leukoc Biol* 87: 905–914.
11. Simard JC, Simon MM, Tessier PA, Girard D (2011) Damage-associated molecular pattern S100A9 increases bactericidal activity of human neutrophils by enhancing phagocytosis. *J Immunol* 186: 3622–3631.
12. Kane D, Roth J, Frosch M, Vogl T, Bresnahan B, et al. (2003) Increased perivascular synovial membrane expression of myeloid-related proteins in psoriatic arthritis. *Arthritis Rheum* 48: 1676–1685.
13. Bernzten HB, Munthe E, Fagerhol MK (1989) A longitudinal study of the leukocyte protein L1 as an indicator of disease activity in patients with rheumatoid arthritis. *J Rheumatol* 16: 1416–1420.
14. Brun JG, Jonsson R, Haga HJ (1994) Measurement of plasma calprotectin as an indicator of arthritis and disease activity in patients with inflammatory rheumatic diseases. *J Rheumatol* 21: 733–738.
15. Luger N, Stoll R, Kucharzik T, Schmid KW, Rohlmann G, et al. (1995) Immunohistochemical distribution and serum levels of the Ca(2+)-binding proteins MRP8, MRP14 and their heterodimeric form MRP8/14 in Crohn's disease. *Digestion* 56: 406–414.
16. Schmid KW, Luger N, Stoll R, Brinkbaumer P, Winde G, et al. (1995) Immunohistochemical demonstration of the calcium-binding proteins MRP8 and MRP14 and their heterodimer (27E10 antigen) in Crohn's disease. *Hum Pathol* 26: 334–337.
17. Cesaro A, Anceriz N, Plante A, Page N, Tardif MR, et al. (2012) An Inflammation Loop Orchestrated by S100A9 and Calprotectin Is Critical for Development of Arthritis. *PLoS ONE* 7: e45478.
18. van Lent PL, Blom AB, Schelbergen RF, Sloetjes A, Lafèber FP, et al. (2012) Active involvement of alarmins S100A8 and S100A9 in the regulation of synovial activation and joint destruction during mouse and human osteoarthritis. *Arthritis Rheum* 64: 1466–1476.
19. Chang KA, Kim HJ, Suh YH (2012) The role of S100a9 in the pathogenesis of Alzheimer's disease: the therapeutic effects of S100a9 knockdown or knockout. *Neurodegener Dis* 10: 27–29.
20. Loser K, Vogl T, Voskort M, Lueken A, Kupas V, et al. (2010) The Toll-like receptor 4 ligands Mrp8 and Mrp14 are crucial in the development of autoreactive CD8(+) T cells. *Nat Med*.
21. Dinarello CA (2010) IL-1: discoveries, controversies and future directions. *Eur J Immunol* 40: 599–606.
22. Dinarello CA (2009) Immunological and inflammatory functions of the interleukin-1 family. *Annu Rev Immunol* 27: 519–550.
23. Dinarello CA (2011) Interleukin-1 in the pathogenesis and treatment of inflammatory diseases. *Blood* 117: 3720–3732.
24. Franchi L, Eigenbrod T, Munoz-Planillo R, Nunez G (2009) The inflammasome: a caspase-1-activation platform that regulates immune responses and disease pathogenesis. *Nat Immunol* 10: 241–247.
25. Yamamoto M, Yaginuma K, Tsutsui H, Sagara J, Guan X, et al. (2004) ASC is essential for LPS-induced activation of procaspase-1 independently of TLR-associated signal adaptor molecules. *Genes Cells* 9: 1055–1067.
26. Kahlenberg JM, Lundberg KC, Kertesz SB, Qu Y, Dubyak GR (2005) Potentiation of caspase-1 activation by the P2X7 receptor is dependent on TLR signals and requires NF-kappaB-driven protein synthesis. *J Immunol* 175: 7611–7622.
27. Perregraux DG, Gabel CA (1998) Human monocyte stimulus-coupled IL-1beta posttranslational processing: modulation via monovalent cations. *Am J Physiol* 275: C1538–1547.
28. Perregraux D, Gabel CA (1994) Interleukin-1 beta maturation and release in response to ATP and nigericin. Evidence that potassium depletion mediated by these agents is a necessary and common feature of their activity. *J Biol Chem* 269: 15195–15203.
29. Ryckman C, McColl SR, Vandal K, de Medicis R, Lussier A, et al. (2003) Role of S100A8 and S100A9 in neutrophil recruitment in response to monosodium urate monohydrate crystals in the air-pouch model of acute gouty arthritis. *Arthritis Rheum* 48: 2310–2320.
30. Ennaciri J, Girard D (2009) IL-4R(alpha), a new member that associates with Syk kinase: implication in IL-4-induced human neutrophil functions. *J Immunol* 183: 5261–5269.
31. Liu F, Xia Y, Parker AS, Verma IM (2012) IKK biology. *Immunol Rev* 246: 239–253.
32. Rubartelli A (2012) Redox control of NLRP3 inflammasome activation in health and disease. *J Leukoc Biol* 92: 951–958.
33. Miki H, Funato Y (2012) Regulation of intracellular signalling through cysteine oxidation by reactive oxygen species. *J Biochem* 151: 255–261.
34. Cruz CM, Rinna A, Forman HJ, Ventura AL, Persechini PM, et al. (2007) ATP activates a reactive oxygen species-dependent oxidative stress response and secretion of proinflammatory cytokines in macrophages. *J Biol Chem* 282: 2871–2879.
35. Loukili N, Rosenblatt-Velin N, Rolli J, Levrand S, Feihl F, et al. (2010) Oxidants positively or negatively regulate nuclear factor kappaB in a context-dependent manner. *J Biol Chem* 285: 15746–15752.
36. Schreck R, Rieber P, Baeuerle PA (1991) Reactive oxygen intermediates as apparently widely used messengers in the activation of the NF-kappa B transcription factor and HIV-1. *EMBO J* 10: 2247–2258.
37. Bauernfeind FG, Horvath G, Stutz A, Alnemri ES, MacDonald K, et al. (2009) Cutting edge: NF-kappaB activating pattern recognition and cytokine receptors license NLRP3 inflammasome activation by regulating NLRP3 expression. *J Immunol* 183: 787–791.
38. Newton RA, Hogg N (1998) The human S100 protein MRP-14 is a novel activator of the beta 2 integrin Mac-1 on neutrophils. *J Immunol* 160: 1427–1435.
39. Sunahori K, Yamamura M, Yamana J, Takasugi K, Kawashima M, et al. (2006) The S100A8/A9 heterodimer amplifies proinflammatory cytokine production by macrophages via activation of nuclear factor kappa B and p38 mitogen-activated protein kinase in rheumatoid arthritis. *Arthritis Res Ther* 8: R69.
40. Riva M, Kallberg E, Bjork P, Hancz D, Vogl T, et al. (2012) Induction of nuclear factor-kappaB responses by the S100A9 protein is Toll-like receptor-4-dependent. *Immunology* 137: 172–182.
41. Robinson MJ, Tessier P, Poulsom R, Hogg N (2002) The S100 family heterodimer, MRP-8/14, binds with high affinity to heparin and heparan sulfate glycosaminoglycans on endothelial cells. *J Biol Chem* 277: 3658–3665.
42. Kallberg E, Vogl T, Liberg D, Olsson A, Bjork P, et al. (2012) S100A9 interaction with TLR4 promotes tumor growth. *PLoS One* 7: e34207.
43. Dendorfer U, Oettgen P, Libermann TA (1994) Multiple regulatory elements in the interleukin-6 gene mediate induction by prostaglandins, cyclic AMP, and lipopolysaccharide. *Mol Cell Biol* 14: 4443–4454.
44. Libermann TA, Baltimore D (1990) Activation of interleukin-6 gene expression through the NF-kappa B transcription factor. *Mol Cell Biol* 10: 2327–2334.
45. Samuel JM, Kelberman D, Smith AJ, Humphries SE, Woo P (2008) Identification of a novel regulatory region in the interleukin-6 gene promoter. *Cytokine* 42: 256–264.
46. Karin M (1995) The regulation of AP-1 activity by mitogen-activated protein kinases. *J Biol Chem* 270: 16483–16486.
47. Fialkow L, Wang Y, Downey GP (2007) Reactive oxygen and nitrogen species as signaling molecules regulating neutrophil function. *Free Radic Biol Med* 42: 153–164.
48. Ryan KA, Smith MF, Jr., Sanders MK, Ernst PB (2004) Reactive oxygen and nitrogen species differentially regulate Toll-like receptor 4-mediated activation of NF-kappa B and interleukin-8 expression. *Infect Immun* 72: 2123–2130.
49. Bauernfeind F, Bartok E, Rieger A, Franchi L, Nunez G, et al. (2011) Cutting edge: reactive oxygen species inhibitors block priming, but not activation, of the NLRP3 inflammasome. *J Immunol* 187: 613–617.
50. Jiang Y, Wang M, Huang K, Zhang Z, Shao N, et al. (2012) Oxidized low-density lipoprotein induces secretion of interleukin-1beta by macrophages via reactive oxygen species-dependent NLRP3 inflammasome activation. *Biochem Biophys Res Commun* 425: 121–126.
51. Martinon F, Mayor A, Tschopp J (2009) The inflammasomes: guardians of the body. *Annu Rev Immunol* 27: 229–265.
52. Hornung V, Latz E (2010) Critical functions of priming and lysosomal damage for NLRP3 activation. *Eur J Immunol* 40: 620–623.
53. Qiao Y, Wang P, Qi J, Zhang L, Gao C (2012) TLR-induced NF-kappaB activation regulates NLRP3 expression in murine macrophages. *FEBS Lett* 586: 1022–1026.
54. Mariathasan S, Newton K, Monack DM, Vucic D, French DM, et al. (2004) Differential activation of the inflammasome by caspase-1 adaptors ASC and Ipaf. *Nature* 430: 213–218.
55. Martinon F, Petrilli V, Mayor A, Tardivel A, Tschopp J (2006) Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 440: 237–241.
56. Mariathasan S, Weiss DS, Newton K, McBride J, O'Rourke K, et al. (2006) Cryopyrin activates the inflammasome in response to toxins and ATP. *Nature* 440: 228–232.
57. He Q, You H, Li XM, Liu TH, Wang P, et al. (2012) HMGB1 promotes the synthesis of pro-IL-1beta and pro-IL-18 by activation of p38 MAPK and NF-kappaB through receptors for advanced glycation end-products in macrophages. *Asian Pac J Cancer Prev* 13: 1363–1370.
58. Schroder K, Tschopp J (2010) The inflammasomes. *Cell* 140: 821–832.
59. Cheneval D, Ramage P, Kastelic T, Szelestenyi T, Niggl H, et al. (1998) Increased mature interleukin-1beta (IL-1beta) secretion from THP-1 cells induced by nigericin is a result of activation of p45 IL-1beta-converting enzyme processing. *J Biol Chem* 273: 17846–17851.
60. Schumann RR, Belka C, Reuter D, Lamping N, Kirschning CJ, et al. (1998) Lipopolysaccharide activates caspase-1 (interleukin-1-converting enzyme) in cultured monocytic and endothelial cells. *Blood* 91: 577–584.
61. Grahames CB, Michel AD, Chessell IP, Humphrey PP (1999) Pharmacological characterization of ATP- and LPS-induced IL-1beta release in human monocytes. *Br J Pharmacol* 127: 1915–1921.
62. Strowig T, Henao-Mejia J, Elinav E, Flavell R (2012) Inflammasomes in health and disease. *Nature* 481: 278–286.

63. Bauer C, Duewell P, Mayer C, Lehr HA, Fitzgerald KA, et al. (2010) Colitis induced in mice with dextran sulfate sodium (DSS) is mediated by the NLRP3 inflammasome. *Gut* 59: 1192–1199.
64. Siegmund B, Lehr HA, Fantuzzi G, Dinarello CA (2001) IL-1 beta -converting enzyme (caspase-1) in intestinal inflammation. *Proc Natl Acad Sci U S A* 98: 13249–13254.
65. Amaral FA, Costa VV, Tavares LD, Sachs D, Coelho FM, et al. (2012) NLRP3 inflammasome-mediated neutrophil recruitment and hypernociception depend on leukotriene B(4) in a murine model of gout. *Arthritis Rheum* 64: 474–484.
66. Youm YH, Adijiang A, Vandanmagsar B, Burk D, Ravussin A, et al. (2011) Elimination of the NLRP3-ASC inflammasome protects against chronic obesity-induced pancreatic damage. *Endocrinology* 152: 4039–4045.
67. Foell D, Wittkowski H, Ren Z, Turton J, Pang G, et al. (2008) Phagocyte-specific S100 proteins are released from affected mucosa and promote immune responses during inflammatory bowel disease. *J Pathol* 216: 183–192.
68. Roseth AG, Schmidt PN, Fagerhol MK (1999) Correlation between faecal excretion of indium-111-labelled granulocytes and calprotectin, a granulocyte marker protein, in patients with inflammatory bowel disease. *Scand J Gastroenterol* 34: 50–54.
69. Lugerling N, Stoll R, Schmid KW, Kucharzik T, Stein H, et al. (1995) The myeloid related protein MRP8/14 (27E10 antigen)—usefulness as a potential marker for disease activity in ulcerative colitis and putative biological function. *Eur J Clin Invest* 25: 659–664.
70. Ryckman C, Gilbert C, de Medicis R, Lussier A, Vandal K, et al. (2004) Monosodium urate monohydrate crystals induce the release of the proinflammatory protein S100A8/A9 from neutrophils. *J Leukoc Biol* 76: 433–440.
71. Averill MM, Kerkhoff C, Bornfeldt KE (2012) S100A8 and S100A9 in cardiovascular biology and disease. *Arterioscler Thromb Vasc Biol* 32: 223–229.
72. Sampson B, Fagerhol MK, Sunderkotter C, Golden BE, Richmond P, et al. (2002) Hyperzincaemia and hypercalprotectinaemia: a new disorder of zinc metabolism. *Lancet* 360: 1742–1745.
73. McInnes IB, Schett G (2007) Cytokines in the pathogenesis of rheumatoid arthritis. *Nat Rev Immunol* 7: 429–442.