

Genetic diversity of Mycobacterium tuberculosis complex strains isolated from patients with pulmonary tuberculosis in Anambra State, Nigeria.

Gertrude N Uzoewulu, Lovett Lawson, Ibeh S Nnanna, Nalin Rastogi, Madhu Goyal

► **To cite this version:**

Gertrude N Uzoewulu, Lovett Lawson, Ibeh S Nnanna, Nalin Rastogi, Madhu Goyal. Genetic diversity of Mycobacterium tuberculosis complex strains isolated from patients with pulmonary tuberculosis in Anambra State, Nigeria.. International Journal of Mycobacteriology, 2016, 5 (1), pp.74-79. <10.1016/j.ijmyco.2015.06.008>. <pasteur-01304573>

HAL Id: pasteur-01304573

<https://hal-riip.archives-ouvertes.fr/pasteur-01304573>

Submitted on 19 Apr 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HOSTED BY

Available at www.sciencedirect.com

ScienceDirect

journal homepage: www.elsevier.com/locate/IJMYCO

Short Communication

Genetic diversity of *Mycobacterium tuberculosis* complex strains isolated from patients with pulmonary tuberculosis in Anambra State, Nigeria



Gertrude N. Uzoewulu ^a, Lovett Lawson ^b, Ibeh S. Nnanna ^c, Nalin Rastogi ^{d,*}, Madhu Goyal ^{e,*}

^a Nnamdi Azikiwe University Teaching Hospital, Nnewi, Anambra State, Nigeria

^b Zankli Medical Centre, Abuja, Nigeria

^c University of Benin, Edo State, Nigeria

^d World Health Organization TB Supranational Reference Laboratory, Institut Pasteur de Guadeloupe, Les Abymes, Guadeloupe, France

^e University of Hertfordshire, Hatfield, England, UK

ARTICLE INFO

Article history:

Received 8 May 2015

Received in revised form

10 June 2015

Accepted 11 June 2015

Available online 29 July 2015

Keywords:

Exact tandem repeat

Mycobacterium tuberculosis

Nigeria

Spoligotyping

Tuberculosis

Variable number of tandem repeats

ABSTRACT

In this study, we analyzed *Mycobacterium tuberculosis* complex (MTC) genetic diversity in Anambra State, Nigeria based on spoligotyping followed by 5-loci exact tandem repeats (ETRs). Spoligotyping of 180 MTC strains isolated in 2009–2011 from pulmonary tuberculosis (TB) patients led to a total of 31 distinct patterns. A comparison with the SITVIT2 international database showed that all the 31 patterns could be classified as Shared-types (SITs) in this database; briefly, 26/31 SITs ($n = 174$ isolates) matched a preexisting shared-type in the database, whereas 5/31 SITs ($n = 6$ isolates) were newly created due to 2 or more strains belonging to an identical new pattern within this study (SIT3396) or after a match with an orphan in the database (SIT3397, SIT3398, SIT3399 and SIT3400). A total of 18/31 SITs containing 167 or 92.8% isolates were clustered within this study (2–89 isolates per cluster) while 13/31 SITs contained unique strains. Using VNTR typing, a total of 36 distinct patterns were identified; 27 patterns ($n = 157$ isolates) matched a pattern already reported in the SITVIT2 database. Combination of both the methods generated 47 combined patterns for the 180 strains: 17 belonged to clustered isolates ($n = 127$ isolates or 70.5%) while 30 corresponded to as many unique strains (note 23 strains could not be typed using 5-loci ETRs). No correlation was found between the spoligotyping pattern and the HIV status of the patient or drug sensitivity of the strain. This study showed that the LAM10-CAM prototype SIT61 accounted for highest number of isolates ($n = 89$) in Anambra State, showing its relative contribution to the TB burden in the study.

© 2015 Production and hosting by Elsevier Ltd. on behalf of Asian African Society for Mycobacteriology.

* Corresponding authors at: Institut Pasteur de Guadeloupe, BP 484, F97183 Les Abymes, Guadeloupe, France (N. Rastogi). School of Life and Medical Sciences, University of Hertfordshire, Hatfield, England, UK (M. Goyal).

E-mail addresses: nrastogi@pasteur-guadeloupe.fr (N. Rastogi), m.goyal@herts.ac.uk (M. Goyal).

Peer review under responsibility of Asian African Society for Mycobacteriology.

<http://dx.doi.org/10.1016/j.ijmyco.2015.06.008>

2212-5531/© 2015 Production and hosting by Elsevier Ltd. on behalf of Asian African Society for Mycobacteriology.

Introduction

Nigeria, with a population of over 150 million, is among the high-tuberculosis (TB)-burden countries and ranks 13th in the world [1]. Multiple-drug-resistant TB (MDR-TB) is another problem, and in a recent study, it has been found that as much as 8% of all cultured specimens were MDR-TB positive in three states in Nigeria [2]. The information available on the incidence, drug susceptibility, and genotyping of the *Mycobacterium tuberculosis* complex (MTC) in Nigeria is limited [3–6]. Additional data are needed to explore the population structure of strains of MTC to identify specific endemic strains in the study area; monitor transmission dynamics to link outbreak cases in communities, hospitals, or institutions; and for better treatment.

Many molecular-typing techniques have been used to differentiate strains of MTC involved in TB infection, among which the spoligotyping method based on the polymorphism of the direct repeat locus is a widely used first-line typing method [2,6]. However, when used alone, the lower discriminatory power of spoligotyping requires that it is ideally used in association with 12, 15, or 24-loci mycobacterial interspersed repetitive unit-variable number of tandem repeats (MIRU-VNTRs) for *M. tuberculosis* molecular epidemiology, or at minima in association with a more convenient five-loci exact tandem repeats (ETRs, [7]) that have been successfully used to improve the potential of spoligotyping for studying the genetic diversity of TB [7–9]. The present study constitutes a first attempt to describe the genetic population structure of MTC circulating in Anambra State, Nigeria using spoligotyping and five-loci ETRs.

Materials and methods

Setting, clinical isolates, and molecular characterization

The study was conducted among patients between the ages of 10 years and 82 years with pulmonary TB attending Nnamdi Azikiwe University Teaching Hospital and different peripheral DOTS clinics in Anambra State during the period 2009–2011. Data regarding the patients' gender, human-immunodeficiency-virus (HIV) status, and age were collected. MTC strains were isolated and identified from 550 sputum samples of suspected TB patients after smear microscopy by the Ziehl-Neelsen method at Nnamdi Azikiwe University Teaching Hospital, Nnewi, and cultured on Löwenstein-Jensen medium at Zankli TB laboratory, Abuja. DNA was extracted using the classical cetyl-trimethyl-ammonium-bromide method as described previously [8,10], and sent to the University of Hertfordshire, Hatfield, England for molecular typing. Spoligotyping was performed using a commercially available kit (Ocimum Biosolutions, Hyderabad, India), following the manufacturer's instructions, and previously described methodology [11], shown to be useful to study the transmission of *M. tuberculosis* [12]. Five-loci ETR (A, B, C, D, and E) typing was performed, as described by Frothingham and Meeker-O'Connell [7]. The exact number of tandem repeats at each locus was analyzed for each strain using polymerase chain reaction.

Database comparison

The identified spoligotypes and five-loci ETR patterns were analyzed using the BioNumerics software (BioSystematica), and compared with the SITVIT2 proprietary database of the Institut Pasteur de Guadeloupe, which is an updated in-house version of the recently released SITVITWEB database [13], available online at http://www.pasteur-guadeloupe.fr:8081/SITVIT_ONLINE/. In this database, spoligotype international type (SIT) and VNTR international type (VIT) designate spoligotype and five-loci ETR patterns shared by two or more patient isolates, as opposed to "orphan," which designates patterns reported for a single isolate. Major phylogenetic clades were assigned according to the signatures provided in the database defining 62 genetic lineages/sublineages. These include various MTC members, such as *Mycobacterium bovis*, *Mycobacterium caprae*, *Mycobacterium microti*, *Mycobacterium canettii*, *Mycobacterium pinnipedii*, and *Mycobacterium africanum*, as well as rules defining major lineages/sublineages for *M. tuberculosis sensu stricto*. These include the Beijing clade, the Central-Asian clade and two sublineages, the East-African-Indian clade and nine sublineages, the Haarlem clade and three sublineages, the Latin-American-Mediterranean (LAM) clade and 12 sublineages (note that two sublineages, LAM7-TUR and LAM10-CAM, were reclassified as Turkey and Cameroon lineages), the ancestral "Manu" family and three sublineages, the S clade, the IS6110-low-banding X clade and three sublineages, and an ill-defined T clade with five sublineages.

The description of predominant clusters in this study (four or more isolates) and their worldwide distribution was studied in function of their reported numbers in various macrogeographical regions in the SITVIT2 database (reported for regions with more than 3% of a given shared type). The definition of macrogeographical regions and subregions (<http://unstats.un.org/unsd/methods/m49/m49regin.htm>) was according to the United Nations scheme (regions: AFRI [Africa], AMER [Americas], ASIA [Asia], EURO [Europe], and OCE [Oceania], subdivided in E [eastern], M [middle], C [central], N [northern], S [southern], SE [southeastern], and W [western]). Note that, in this scheme, CARIB (Caribbean) belongs to Americas, while Oceania is subdivided in four subregions: AUST (Australasia), MEL (Melanesia), MIC (Micronesia), and POLY (Polynesia). Furthermore, Russia was attributed a new subregion by itself (Northern Asia), instead of including it among the rest of Eastern Europe, reflecting its geographical localization, as well as due to the similarity of specific TB genotypes circulating in Russia (a majority of Beijing genotypes) with those prevalent in Central, Eastern, and Southeastern Asia. Finally, the three-letter country codes were according to http://en.wikipedia.org/wiki/ISO_3166-1_alpha-3.

Ethical considerations

An ethical clearance was granted by the hospital ethical committee, and informed consent was obtained from each patient.

the LAM10–CAM family, as shown previously for the Beijing family [17], remains speculative, but should be studied in future studies.

The result of the comparison of spoligotyping versus VNTRs demonstrated that spoligotyping analysis alone was less discriminative than when used in association with five-loci ETRs; for example, isolates with identical spoligotype pattern SIT61 were all split into different VIT patterns: VIT58, VIT180, VIT299, VIT406, VIT407, and VIT409, and four distinct orphan patterns (Table S1). Thumamo et al. [17] also showed higher discrimination using 12-loci MIRU typing as compared to spoligotyping. Hence, future studies using 15- and/or 24-loci MIRUs might even lead to a much higher discrimination and more conclusive results in our study area.

A significant number of *M. africanum* AFRI_2 sublineage ($n = 19$) were found among pulmonary TB patients attending the Nnamdi Azikiwe University Teaching Hospital ($n = 13$) and Onitsha ($n = 6$), underlining that *M. africanum* also plays an important role in the prevailing TB epidemic in Nigeria. Interestingly, in our study, we had only one isolate belonging to the AFRI_1 sublineage. In a recent study, as high as 33.3% isolates from the Cross River State, Nigeria were shown to belong to the AFRI_2 sublineage [17]. These results are in contrast to the study by Lawson et al. [2], where 13% of the isolates belonged to the AFRI_1 sublineage. Cadmus et al [3] also showed that 13% of TB cases in Nigeria were caused by *M. africanum* (essentially AFRI_1) and *M. bovis*. In another study from Guinea-Bissau, almost all the *M. africanum* strains belonged to the AFRI_1 sublineage [18]. However, Thumamo et al [17] underlined that misidentification of *M. africanum* strains due to substantial heterogeneity leading to their erroneous identification as *M. bovis* and/or *M. tuberculosis* is more common than thought, which requires careful reexamination of the trends of prevalence of the *M. africanum* sublineages in different parts of Africa. Last, but not least, in addition to the LAM10–CAM and AFRI_2 lineages, we also found 4.4% of isolates belonging to the Haarlem sublineage H3 and 4% isolates typed as Manu_ancestor; these results are in agreement with other studies in Nigeria showing a small percentage of isolates belonging to these genotypes [2,6].

Although drug-susceptibility testing was not systematically performed, the partial drug-susceptibility-testing data available ($n = 87$ isolates) showed a relatively high proportion of MDR-TB cases ($n = 14$ or 16%). However, no correlation was found between the spoligotyping pattern and the HIV status of the patient, or drug sensitivity of the strain. However, tracing the route of infection and the risk factors for TB transmission based on patient records, we suggest that TB in clustered patients most probably resulted from recently acquired infection [19,20] versus reactivation among patients infected by unique patterns [20,21]. Last, but not least, out of the total of seven strains belonging to the SIT523/Manu_ancestor lineage (all the 43 spacers present by spoligotyping), only one strain could be typed by VNTRs and belonged to VIT153; the six remaining strains could not be amplified (Table S1). It will be important to look for such strains in future investigations to verify if they did not derive from mixed-strain infections as revealed recently by a novel computational approach [22]. Whether

it is the case or not would require extended MIRU–VNTR typing to exclude multiple bands.

Conflict of interest

None declared.

Acknowledgments

Nalin Rastogi is grateful to David Couvin (Institut Pasteur de Guadeloupe) for helping with the SITVIT2 database query.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.ijmyco.2015.06.008>.

REFERENCES

- [1] World Health Organization (WHO), Global Tuberculosis Report, WHO, Geneva, Switzerland, 2012. Available from http://apps.who.int/iris/bitstream/10665/75938/1/9789241564502_eng.pdf.
- [2] L. Lawson, J. Zhang, M.K. Gomgnimbou, et al, A molecular epidemiological and genetic diversity study of tuberculosis in Ibadan, Nnewi and Abuja, Nigeria, PLoS ONE 7 (2012) e38409.
- [3] S. Cadmus, S. Palmer, M. Okker, et al, Molecular analysis of human and bovine tubercle bacilli from a local setting in Nigeria, J. Clin. Microbiol. 44 (2006) 29–34.
- [4] S.I.B. Cadmus, A.O. Jenkins, J. Godfroid, et al, *Mycobacterium tuberculosis* and *Mycobacterium africanum* in stools from children attending an immunization in Ibadan, Nigeria, Int. J. Infect. Dis. 13 (2009) 740–744.
- [5] A.O. Kehinde, F.A. Obaseki, O.C. Ishola, et al, Multidrug resistance to *Mycobacterium tuberculosis* in a tertiary hospital, J. Natl Med. Assoc. 99 (2007) 1185–1189.
- [6] A. Ani, T. Bruvik, Y. Okoh, et al, Genetic diversity of *Mycobacterium tuberculosis* complex in Jos, Nigeria, BMC Infect. Dis. 10 (2010) 189–193.
- [7] R. Frothingham, W. Meeker-O’Connell, Genetic diversity in *Mycobacterium tuberculosis* complex based on variable numbers of tandem DNA repeats, Microbiology 144 (1998) 1189–1196.
- [8] O.V. Surikova, D.S. Voitech, G. Kuzmicheoi, et al, Efficient differentiation of *Mycobacterium tuberculosis* strains of the W-Beijing family from Russia using highly polymorphic VNTR loci, Eur. J. Epidemiol. 20 (2005) 963–974.
- [9] S. Garbaccio, A. Macias, E. Shimizu, et al, Association between spoligotype-VNTR types and virulence of *Mycobacterium bovis* in cattle, Virulence 5 (2014) 297–302.
- [10] D.L. Williams, T.P. Gills, W.G. Dupree, Ethanol fixation of sputum sediments for DNA-based detection of *Mycobacterium tuberculosis*, J. Clin. Microbiol. 33 (1995) 1558–1561.
- [11] J. Kamerbeek, L. Schouls, A. Kolk, et al, Simultaneous detection and strain differentiation of *Mycobacterium tuberculosis* for diagnosis and epidemiology, J. Clin. Microbiol. 35 (1997) 907–914.
- [12] D. Goguet, Y.O. Salmoniere, H.M. Li, et al, Evaluation of spoligotyping in a study of the transmission of *Mycobacterium tuberculosis*, J. Clin. Microbiol. 35 (1997) 2210–2214.
- [13] C. Demay, B. Liens, T. Burguière, et al, SITVITWEB—a publicly available international multimarker database for studying

- Mycobacterium tuberculosis* genetic diversity and molecular epidemiology, *Infect. Genet. Evol.* 12 (2012) 755–766.
- [14] D. Yeboah-Manu, A. Asante-Poku, T. Bodmer, et al, Genotypic diversity and drug susceptibility patterns among *Mycobacterium tuberculosis* complex isolates from South-Western Ghana, *PLoS ONE* 6 (2011) e21906.
- [15] S. Homolka, E. Post, B. Oberhauser, High genetic diversity among *Mycobacterium tuberculosis* complex strains from Sierra Leone, *BMC Microbiol.* 8 (2008) 103–110.
- [16] S. Godreuil, G. Torrea, D. Terru, First molecular epidemiology study of *Mycobacterium tuberculosis* in Burkina Faso, *J. Clin. Microbiol.* 45 (2007) 921–927.
- [17] B.P. Thumamo, A.E. Asuquo, L.N. Abia-Bassey, et al, Molecular epidemiology and genetic diversity of *Mycobacterium tuberculosis* complex in the Cross River State, Nigeria, *Infect. Genet. Evol.* 12 (2012) 671–677.
- [18] D. Bonard, P. Msellati, L. Rigouts, et al, What is the meaning of repeated isolation of *Mycobacterium africanum*?, *Int J. Tuberc. Lung Dis.* 4 (2000) 1176–1180.
- [19] D. Alland, G.E. Kalkut, A.R. Moss, et al, Transmission of tuberculosis in New York City: an analysis by DNA fingerprinting and conventional epidemiological methods, *N. Engl. J. Med.* 330 (1994) 1710–1716.
- [20] M.W. Borgdorff, N. Nagelkerke, D. van Soolingen, et al, Analysis of tuberculosis transmission between nationalities in the Netherlands in the period 1993–1995 using DNA fingerprinting, *Am. J. Epidemiol.* 147 (1998) 187–195.
- [21] J.R. Glynn, A.C. Crampin, H. Traore, et al, Determinants of cluster size in large, population-based molecular epidemiology study of tuberculosis, Northern Malawi, *Emerg. Infect. Dis.* 13 (1998) 1060–1066.
- [22] L.C. Lazzarini, J. Rosenfeld, R.C. Huard, et al, *Mycobacterium tuberculosis* spoligotypes that may derive from mixed strain infections are revealed by a novel computational approach, *Infect. Genet. Evol.* 12 (2012) 798–806.