



**HAL**  
open science

## Reversible Reprogramming of Circulating Memory T Follicular Helper Cell Function during Chronic HIV Infection.

Rafael Cubas, Julien van Grevenynghe, Saintedym Wills, Lela Kardava, Brian H Santich, Clarisa M Buckner, Roshell Muir, Virginie Tardif, Carmen Nichols, Francesco Procopio, et al.

► **To cite this version:**

Rafael Cubas, Julien van Grevenynghe, Saintedym Wills, Lela Kardava, Brian H Santich, et al.. Reversible Reprogramming of Circulating Memory T Follicular Helper Cell Function during Chronic HIV Infection.. *Journal of Immunology*, 2015, 195 (12), pp.5625-36. 10.4049/jimmunol.1501524 . pasteur-01351053

**HAL Id: pasteur-01351053**

**<https://riip.hal.science/pasteur-01351053>**

Submitted on 2 Aug 2016

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - ShareAlike 4.0 International License

## Immune Checkpoint Proteins & Antibodies in Sino Biological

- ✓ Immune checkpoint targets with hot research issues;
- ✓ Proteins with multiple species, high purity and validated bioactivity;
- ✓ Antibodies generated against full-length proteins with multiple applications.



## Reversible Reprogramming of Circulating Memory T Follicular Helper Cell Function during Chronic HIV Infection

This information is current as of August 9, 2016.

Rafael Cubas, Julien van Grevenynghe, Saintedym Wills, Lela Kardava, Brian H. Santich, Clarisa M. Buckner, Roshell Muir, Virginie Tardif, Carmen Nichols, Francesco Procopio, Zhong He, Talibah Metcalf, Khader Ghneim, Michela Locci, Petronella Ancuta, Jean-Pierre Routy, Lydie Trautmann, Yuxing Li, Adrian B. McDermott, Rick A. Koup, Constantinos Petrovas, Steven A. Migueles, Mark Connors, Georgia D. Tomaras, Susan Moir, Shane Crotty and Elias K. Haddad

*J Immunol* 2015; 195:5625-5636; Prepublished online 6 November 2015;  
doi: 10.4049/jimmunol.1501524  
<http://www.jimmunol.org/content/195/12/5625>

- 
- Supplementary Material** <http://www.jimmunol.org/content/suppl/2015/11/06/jimmunol.1501524.DCSupplemental.html>
- References** This article **cites 42 articles**, 15 of which you can access for free at: <http://www.jimmunol.org/content/195/12/5625.full#ref-list-1>
- Subscriptions** Information about subscribing to *The Journal of Immunology* is online at: <http://jimmunol.org/subscriptions>
- Permissions** Submit copyright permission requests at: <http://www.aai.org/ji/copyright.html>
- Author Choice** Freely available online through *The Journal of Immunology* [Author Choice option](#)
- Email Alerts** Receive free email-alerts when new articles cite this article. Sign up at: <http://jimmunol.org/cgi/alerts/etoc>



# Reversible Reprogramming of Circulating Memory T Follicular Helper Cell Function during Chronic HIV Infection

Rafael Cubas,\* Julien van Grevenynghe,<sup>†</sup> Saintedym Wills,<sup>‡</sup> Lela Kardava,<sup>§</sup> Brian H. Santich,<sup>§</sup> Clarisa M. Buckner,<sup>§</sup> Roshell Muir,\* Virginie Tardif,\* Carmen Nichols,\* Francesco Procopio,<sup>¶</sup> Zhong He,\* Talibah Metcalf,\* Khader Ghneim,\* Michela Locci,<sup>||</sup> Petronella Ancuta,<sup>#,\*\*</sup> Jean-Pierre Routy,<sup>††,‡‡,§§</sup> Lydie Trautmann,\* Yuxing Li,<sup>¶¶,|||</sup> Adrian B. McDermott,<sup>###</sup> Rick A. Koup,<sup>###</sup> Constantinos Petros,<sup>###</sup> Steven A. Migueles,<sup>\*\*\*</sup> Mark Connors,<sup>\*\*\*</sup> Georgia D. Tomaras,<sup>‡</sup> Susan Moir,<sup>§</sup> Shane Crotty,<sup>||,†††,‡‡‡</sup> and Elias K. Haddad\*

Despite the overwhelming benefits of antiretroviral therapy (ART) in curtailing viral load in HIV-infected individuals, ART does not fully restore cellular and humoral immunity. HIV-infected individuals under ART show reduced responses to vaccination and infections and are unable to mount an effective antiviral immune response upon ART cessation. Many factors contribute to these defects, including persistent inflammation, especially in lymphoid tissues, where T follicular helper (T<sub>fh</sub>) cells instruct and help B cells launch an effective humoral immune response. In this study we investigated the phenotype and function of circulating memory T<sub>fh</sub> cells as a surrogate of T<sub>fh</sub> cells in lymph nodes and found significant impairment of this cell population in chronically HIV-infected individuals, leading to reduced B cell responses. We further show that these aberrant memory T<sub>fh</sub> cells exhibit an IL-2-responsive gene signature and are more polarized toward a Th1 phenotype. Treatment of functional memory T<sub>fh</sub> cells with IL-2 was able to recapitulate the detrimental reprogramming. Importantly, this defect was reversible, as interfering with the IL-2 signaling pathway helped reverse the abnormal differentiation and improved Ab responses. Thus, reversible reprogramming of memory T<sub>fh</sub> cells in HIV-infected individuals could be used to enhance Ab responses. Altered microenvironmental conditions in lymphoid tissues leading to altered T<sub>fh</sub> cell differentiation could provide one explanation for the poor responsiveness of HIV-infected individuals to new Ags. This explanation has important implications for the development of therapeutic interventions to enhance HIV- and vaccine-mediated Ab responses in patients under ART. *The Journal of Immunology*, 2015, 195: 5625–5636.

**T** follicular helper (T<sub>fh</sub>) cells are a specialized subset of CD4<sup>+</sup> T cells that provide help to B cells in germinal centers (GCs) and instruct B cell differentiation into

affinity-matured, long-lived memory B cells and plasma cells (1). T<sub>fh</sub> cells deliver critical signals to GC B cells via costimulatory molecules and lymphokine secretion and induce somatic hypermutation

\*Vaccine and Gene Therapy Institute of Florida, Port St. Lucie, FL 34987; <sup>†</sup>Institut National de la Recherche Scientifique, Institut Armand-Frappier, Laval H7V 1B7, Quebec, Canada; <sup>‡</sup>Department of Immunology and the Duke Human Vaccine Institute, Duke University, Durham, NC 27710; <sup>§</sup>Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892; <sup>¶</sup>Service d'Immunologie et Allergie, Centre Hospitalier Universitaire Vaudois, 1011 Lausanne, Switzerland; <sup>||</sup>Division of Vaccine Discovery, La Jolla Institute for Allergy and Immunology, La Jolla, CA 92037; <sup>¶¶</sup>Department of Medicine, University of Montreal, Montreal, Quebec H3C 3J7, Canada; <sup>###</sup>Department of Microbiology, Infectiology and Immunology, Faculty of Medicine, University of Montreal, Montreal, Quebec H3C 3J7, Canada; <sup>††</sup>Chronic Viral Illness Service, McGill University Health Centre, Montreal, Quebec H3H 2R9, Canada; <sup>‡‡</sup>Research Institute, McGill University Health Centre, Montreal, Quebec H3H 2R9, Canada; <sup>§§</sup>Division of Hematology, McGill University Health Centre, Montreal, Quebec H3H 2R9, Canada; <sup>¶¶</sup>International AIDS Vaccine Initiative Neutralizing Antibody Center, The Scripps Research Institute, La Jolla, CA 92037; <sup>|||</sup>Department of Immunology and Microbial Science, The Scripps Research Institute, La Jolla, CA 92037; <sup>###</sup>Immunology Laboratory, Vaccine Research Center, National Institute for Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892; <sup>\*\*\*</sup>HIV-Specific Immunity Section, Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892; <sup>†††</sup>Department of Medicine, School of Immunology, University of California, San Diego, La Jolla, CA 92093; and <sup>‡‡‡</sup>Center for HIV/AIDS Vaccine Immunology and Immunogen Discovery, La Jolla, CA 92037

ORCIDiDs: 0000-0002-2952-4081 (J.v.G.); 0000-0003-2379-8679 (S.W.); 0000-0001-9135-6026 (L.K.); 0000-0001-5869-3661 (B.H.S.); 0000-0002-8178-8076 (C.M.B.); 0000-0003-0364-9625 (C.N.); 0000-0002-7369-9989 (Z.H.); 0000-0003-0548-1942 (K.G.); 0000-0003-1922-5640 (P.A.).

Received for publication July 7, 2015. Accepted for publication October 6, 2015.

This work was supported by National Institutes of Health Grant IRO1AI106482-01A1 (to E.K.H.); the Mathilde Krim Fellowship from the American Foundation for AIDS Research (Grant 10867155-RKVA) (to R.C.); and in part by Canadian Institutes of Health Research Grant 103230, the Canadian Institutes of Health Research Canadian Trials Network/HIV Trial Network (CTN 247), the Fonds de la Recherche en Santé du Québec Maladies Infectieuses, and the Duke Center for AIDS Research Immunology Core (National Institutes of Health Grant AI064518). Y.L. was funded by National Institutes of Health/National Institute of Allergy and Infectious Diseases Grant R01AI102766 and National Institute of Allergy and Infectious Diseases Devel-

opment Grant P30AI36214, the Center for AIDS Research, and the University of California, San Diego. J.-P.R. holds the Louis Lowenstein Chair in Hematology-Oncology, McGill University.

R.C. designed the study, developed and performed experiments, interpreted the data, generated figures, and prepared the manuscript; J.v.G. and Z.H. collaborated on the coculture assay development and planning; S.W. performed and analyzed the HIV-1 multiplex binding assays; L.K. performed and analyzed the evaluation of Ag-specific B cells by flow cytometry; B.H.S. and C.M.B. helped with B cell assays; F.P., C.N., and R.M. helped with the BioMark assays and primer libraries; R.M. and V.T. helped with some of the coculture assays and ELISAs; T.M. assisted in the processing of blood samples; K.G. helped with statistical analysis; M.L. provided intellectual input; P.A. and J.-P.R. provided valuable samples and intellectual input and discussion; L.T. provided valuable discussion; Y.L. provided the HIV YU2 gp140 probe and A.B.M. provided the H1N1 hemagglutinin probe as well as intellectual input for the detection of Ag-specific B cells by flow cytometry; R.A.K. and C.P. provided intellectual input and discussion; S.A.M. and M.C. provided valuable samples and discussion; G.D.T. provided valuable input and discussion and helped to design and analyze the HIV-1 multiplex binding assays; S.M. provided valuable intellectual input and discussion and helped to design and analyze the evaluation of Ag-specific B cells by flow cytometry; S.C. provided valuable intellectual input and discussion and helped design experiments; E.K.H. conceived and directed the research, designed experiments, analyzed data, and helped write the manuscript.

Address correspondence and reprint requests to Dr. Elias K. Haddad, Drexel University, Division of Infectious Diseases and HIV Medicine, 245 North 15th Street, Philadelphia, PA 19102. E-mail address: elias.elhaddad@drexelmed.edu

The online version of this article contains supplemental material.

Abbreviations used in this article: ART, antiretroviral therapy; ASC, Ab-secreting cell; CA, chronic aviremic; cT<sub>fh</sub>, circulating T<sub>fh</sub>; EC, elite controller; GC, germinal center; HC, healthy control; LN, lymph node; MFI, mean fluorescence intensity; NIH, National Institutes of Health; RT, room temperature; SEB, staphylococcal enterotoxin B; T<sub>fh</sub>, T follicular helper.

This article is distributed under The American Association of Immunologists, Inc., [Reuse Terms and Conditions for Author Choice articles](#).

Copyright © 2015 by The American Association of Immunologists, Inc. 0022-1767/15/\$25.00

on Ag-specific B cells (1, 2). The differentiation of Tfh cells involves multiple signals that lead to the activation of various transcription factors, with the core programming being largely mediated by Bcl6, Maf, STAT3, and STAT4 transcription factors (3). IL-2 is a potent inhibitor of Tfh cell differentiation, as it interferes with Bcl6 and CXCR5 expression (4–6) and is dose limiting for Th1 versus Tfh cell differentiation (4). In humans, administration of IL-2 *in vivo* did not yield any clinical benefits to HIV-infected subjects. However, PBMCs obtained from IL-2-treated patients showed increased in CD25 expression and production of inflammatory cytokines (7, 8).

During HIV infection, vast cytokine imbalances (9) in lymphoid tissues can persist despite antiretroviral therapy (ART), affecting both T cell and B cell homeostasis (10). It has recently been shown that HIV continues to undergo low-level replication in lymphoid tissues, maintaining a state of immune activation in individuals undergoing ART. In addition, B cell follicles appear to represent protected niches where viral replication can persist despite the presence of strong antiviral CD8<sup>+</sup> T cell responses (11). It is therefore conceivable that persistent inflammatory signals could skew the differentiation profile of Tfh cells, leading to an altered polarization and adoption of cytokine signaling programs, which could affect their ability to provide adequate B cell help. The lymph node (LN) microenvironment is likely responsible for inducing major changes on Tfh cell function. As evidence for this, we have previously shown that despite an increase in the frequency of Tfh cells, their function is impaired in LNs from HIV-infected individuals (12).

Recently, a population of blood circulating Tfh (cTfh) cells has been described that represents a circulating memory counterpart of LN-resident Tfh cells (13–15). If cTfh cells arise from Tfh cell precursors in LNs, changes in the LN microenvironment could imprint particular effector fates or functional skewing during their

differentiation into memory cells. With the difficulty of obtaining LN samples, investigating the functional status of memory Tfh cells in peripheral blood could provide a glimpse into the microenvironmental changes that took place in LNs during the differentiation of these cells into memory counterparts. With the recent demonstration that HIV continues to undergo low-level replication in lymphoid tissues maintaining a state of immune activation despite ART (16), it is conceivable that Tfh cells encounter environmental signals, which could affect their differentiation and phenotype when transitioning into memory Tfh cells.

In this article, we show that memory Tfh cells from peripheral blood of chronic aviremic (CA) individuals are functionally impaired in their ability to provide adequate B cell help when compared with those from elite controllers (ECs). This impairment stems in part from increased IL-2 signaling, which can affect the differentiation and function of memory Tfh cells. We further demonstrate that memory Tfh cells in CA individuals are more polarized toward a Th1 phenotype with increased levels of Tbet expression; increased secretion of IFN- $\gamma$ ; and reduced expression of Bcl6, CXCR5, and IL-21. Importantly, memory Tfh cells from CA subjects are reprogrammable and still show plasticity, as their function can be restored at least partly by interfering with IL-2 signaling.

## Materials and Methods

### Human samples

All procedures were approved by the Institutional Review Boards at the relevant institutions, and all participants provided signed informed consent. Frozen PBMCs were obtained from CA individuals and ECs who were recruited from the Clinical Research Center, National Institutes of Health (NIH). Patient information can be found in Tables I and II. PBMCs from CA individuals and ECs were also obtained from patients who participated in the blood banking study from the Chronic Viral Illness Service at McGill University Health Centre. Informed consent was signed by each participant, and the study was approved by the Research Ethics Board of

Table I. Patient information on the CA-treated individuals used in various assays

Patient ID	Age	Diagnosis Year	CD4 <sup>+</sup> T Cell Count (cells/mm <sup>3</sup> )	CD8 <sup>+</sup> T Cell Count (cells/mm <sup>3</sup> )	Plasma HIV RNA (copies/ml)	ART Duration (y)
CA1	26	2000	611	509	<50	9
CA2	53	1992	434	583	<50	13
CA3	29	2001	702	586	<50	8
CA4	39	1994	411	368	<50	12
CA5	55	1985	657	1088	<50	16
CA6	35	2003	517	259	<50	6
CA7	47	NA	269	282	<50	NA
CA8	50	2002	569	462	<50	8
CA9	46	1998	391	620	<50	14
CA10	49	2004	309	980	<50	8
CA11	57	1986	670	1006	<50	NA
CA12	59	1996	466	1329	<50	9
CA13	46	1995	1079	324	<50	13
CA14	43	1993	696	768	<50	7
CA15	35	2001	500	625	<50	5
CA16	43	1986	698	769	<50	8
CA17	21	1992	850	648	<50	7
CA18	41	1994	478	669	<50	6
CA19	55	1986	276	414	<50	9
CA20	33	2002	773	634	<50	3
CA21	55	1986	750	1379	<50	9
CA22	53	1985	333	1018	<50	26
CA23	46	2000	530	1280	<50	5
CA24	55	1995	536	676	<50	18
CA25	32	1995	613	302	<50	17
CA26	52	2000	785	1194	<50	14
CA27	45	1999	572	829	<50	15
CA28	24	1992	623	657	<50	6
CA29	53	1997	932	1134	<50	17

HIV-infected CA individuals were assessed for CD4 and CD8 counts as well as virus load. NA, not available.

Table II. Patient information on the ECs used in various assays

Patient ID	Age	Diagnosis Year	CD4 <sup>+</sup> T Cell Count (cells/mm <sup>3</sup> )	CD8 <sup>+</sup> T Cell Count (cells/mm <sup>3</sup> )	Plasma HIV RNA (copies/ml)
EC01	48	NA	788	2498	<50
EC02	36	1999	711	574	<50
EC03	54	1995	711	574	<50
EC05	37	1994	641	1759	121
EC06	47	1991	611	583	<50
EC07	43	NA	613	246	<50
EC08	44	1989	913	519	<50
EC09	40	1992	1116	708	<50
EC10	50	1985	673	833	<50
EC11	47	2004	679	556	<50
EC12	57	1985	1334	805	<50
EC13	61	1988	833	975	<50
EC14	50	1985	902	722	<50
EC15	48	1985	618	588	<50
EC16	44	1989	729	492	<50
EC17	52	2007	1943	716	<50
EC18	60	1994	749	638	<50
EC19	52	1995	552	607	<50
EC20	47	1997	712	260	<50

HIV-infected ECs were assessed for CD4 and CD8 counts as well as virus load. NA, not available.

the Research Institute of McGill University Health Centre. Blood samples from healthy donors were obtained from Martin Memorial Health Systems. PBMCs were isolated from blood by density-gradient centrifugation using Ficoll-Paque PLUS (GE Healthcare). Cells were subsequently frozen and used for the relevant assays. Fresh human tonsils were obtained from tonsillectomies performed at Martin Memorial Health Systems. Tonsils were processed as previously described (10).

#### Abs

The following fluorochrome-conjugated Abs were used: anti-human CD3 (clone UCHT1, 1:200), anti-human CD4 (clone RPA-T4, 1:200), anti-human CD19 (clone HIB19, 1:100), anti-human CD27 (clone O323, 1:100), anti-human CD38 (clone HIT2, 1:100), anti-human CD95 (clone DX2, 1:100), anti-human ICOS (clone C398.4A, 1:100), anti-human CD40L (clone 24-31, 1:10), anti-human PD-1 (clone EH12.2H7, 1:100), anti-human CXCR3 (clone G025H7, 1:200), anti-human CTLA4 (clone L3D10, 1:10), anti-human CD69 (clone FN50, 1:50), and anti-human IL-7R $\alpha$  (clone A019D5, 1:100) were all from BioLegend. Anti-human CXCR5 (clone RF8 B2, 1:200), anti-human Bcl-6 (clone K112-91, 1:20), anti-human IgD (clone IA6-2, 1:50), anti-human Ki-67 (MOPC-21, 1:10), and anti-human CD21 (clone B-ly4, 1:100) were from BD Biosciences, and anti-human CD45RA (clone 2H4LDH11LDB9, 1:200) was from Beckman Coulter. LIVE/DEAD Fixable Dead Cell Stain from Invitrogen was used to gate on live cells as well as Annexin V from BD Biosciences.

#### Coculture assays

The cTfh cells defined as (CD3<sup>+</sup>CD4<sup>+</sup>CD45RA<sup>-</sup>CXCR5<sup>+</sup>CXCR3<sup>-</sup>) or the non-cTfh cells defined as (CD3<sup>+</sup>CD4<sup>+</sup>CD45RA<sup>-</sup>CXCR5<sup>+</sup>CXCR3<sup>+</sup> or CD3<sup>+</sup>CD4<sup>+</sup>CD45RA<sup>-</sup>CXCR5<sup>-</sup>) were sorted and placed in coculture with memory B cells (CD3<sup>-</sup>CD19<sup>+</sup>CD27<sup>+</sup>) at equal numbers ( $1 \times 10^4$  cells each) in the presence of 100 ng/ml staphylococcal enterotoxin B (SEB) (Toxin Technology) in complete RPMI 1640 medium (Life Technologies). Supernatants and cells were harvested at 3, 5, or 7 d for subsequent analysis. SEB was added at the onset of the cocultures to mimic the interaction between T cells and B cells in a polyclonal setting (13). The presence of SEB was required to induce T cell activation and subsequent T/B cell crosstalk (Supplemental Fig. 2C). Of note, SEB did not activate memory B cells when cultured alone (Supplemental Fig. 2C, 2G). The absolute number of cells was calculated using Flow-Count Fluorospheres (Beckman Coulter) and following the manufacturer's instructions. To sort CFSE<sup>lo/hi</sup> cells from the coculture, total PBMCs were first stained with 1.25  $\mu$ M CFSE, followed by thorough washing. Cells were then sorted and placed in coculture, as above. After either 24 h or 5 d, cells were sorted for BioMark analysis. Alternatively, cells were kept for 7 d in coculture followed by flow cytometry analysis.

#### ELISA

ELISAs were performed as previously described (10). In brief, for analysis of total IgG output from the cocultures, 96-well Immulon 2HB plates

(Thermo Scientific) were coated overnight at 4°C with monoclonal anti-human IgG (Mabtech; clone MT91/145) at a concentration of 1  $\mu$ g ml<sup>-1</sup> in PBS. The following day, plates were washed five times with PBS + 0.05% Tween 20. Plates were subsequently blocked for 1 h with PBS + 10% FBS at room temperature (RT). Plates were subsequently washed and the culture supernatants added at different dilutions for 1 h at RT. After washing, the plates were incubated with 1  $\mu$ g ml<sup>-1</sup> of anti-human IgG-biotin (Mabtech; clone MT78/145) for 1 h at RT. The plates were then washed as above and incubated with streptavidin-HRP (Mabtech) for 45 min at RT. Plates were thoroughly washed, and 100  $\mu$ l tetramethylbenzidine substrate (Sigma-Aldrich) was added until the appearance of color. The enzymatic reaction was stopped by the addition of 50  $\mu$ l 1 M H<sub>3</sub>PO<sub>4</sub>. The OD (450 nM) was then measured using a SpectraMax plus 384-plate reader (Molecular Devices).


To detect Env-specific Ab responses, 96-well, high-binding, half-area ELISA plates (Greiner Bio-One) were coated overnight at 4°C with 1  $\mu$ g ml<sup>-1</sup> recombinant HIV-1 envelope protein (ProSpec) in PBS. The following day, plates were washed five times with PBS + 0.05% Tween 20 and blocked for 1 h with PBS + 10% FBS at RT. Plates were subsequently washed and the culture supernatants added at different dilutions for 2 h at RT. For the standard curve, human HIV Ig was used. This reagent was obtained through the NIH AIDS Research and Reference Reagent Program, Division of AIDS, National Institute of Allergy and Infectious Diseases, NIH (cat. no. 3957) from Dr. Luiz Barbosa, National Heart Lung and Blood Institute. The plates were then washed and incubated with 1  $\mu$ g ml<sup>-1</sup> anti-human IgG-biotin (Mabtech; clone MT78/145) for 1 h at RT, followed by washing and incubation with streptavidin-HRP (Mabtech) for 45 min at RT. Plates were then thoroughly washed, and 50  $\mu$ l tetramethylbenzidine substrate (Sigma-Aldrich) was added until the appearance of color. The enzymatic reaction was stopped by the addition of 25  $\mu$ l 1 M H<sub>3</sub>PO<sub>4</sub>. The OD (450 nM) was then measured using a SpectraMax plus 384-plate reader (Molecular Devices).

#### BioMark

Assays (Primers and Probes) were designed using the Roche Universal Probe Library Assay Design Center (www.universalprobelibrary.com) and were designed to detect multiple transcripts, without respect to isoform prevalence. Cells were sorted directly into PCR plates and immediately frozen. RNA extraction and PCR amplifications were done according to manufacturer recommendations and as we have previously published (17).

#### Evaluation of Ag-specific B cells by flow cytometry

Mature B cells were stained with Abs against B cell markers, as well as fluorochrome-conjugated probes derived from HIV YU2 gp140 and influenza H1N1 HA, as previously described (18). HIV-specific memory B cells were identified using a trimeric HIV envelope gp140 probe. This probe contains a site-specific biotinylation motif Avi tag at the C termini that can be fluorescently labeled. The HIV-gp140 probe was conjugated with streptavidin-allophycocyanin (Invitrogen). FACS analyses were



**FIGURE 1.** cTfh cells from CA individuals are functionally impaired in their ability to provide adequate B cell help. **(A)** PBMCs from HC subjects were sorted into cTfh cells ( $CD3^+CD4^+CD45RA^-CXCR5^+CXCR3^-$ ), non-cTfh cells ( $CD3^+CD4^+CD45RA^-CXCR5^+CXCR3^+$  and  $CD3^+CD4^+CD45RA^-CXCR5^-$ ), and  $CD27^+$  memory B cells and placed in coculture for 7 d in the presence of SEB. Shown are the total levels of IgG in the coculture supernatants after 3, 5, and 7 d, as measured by ELISA.  $n = 4-12$ . **(B and C)** PBMCs from HC, EC, and CA subjects were also sorted into cTfh cells and memory B cells and placed in coculture for 7 d in the presence of SEB. **(B)** Total levels of IgG after 7 d. **(C)** Absolute number of live B cells after 7 d.  $n = 16-24$ . **(D-H)** Alternatively, either cTfh cells or memory B cells from a sorted pool of HCs were used to substitute cTfh cells or memory B cells in cocultures from EC or CA subjects. **(D)** Total levels of IgG after replacing cTfh cells from CA subjects with cTfh cells from HCs. **(E)** Total levels of IgG after replacing cTfh cells from ECs with cTfh cells from HCs. **(F)** Comparison of total IgG levels in cocultures from ECs with CA subjects with substituted cTfh cells from HCs. **(G)** Total IgG levels in cocultures after replacing cTfh cells from ECs and CA subjects with cTfh cells from HCs. **(H)** Total IgG levels in cocultures after replacing memory B cells from ECs and CA subjects with memory B cells from HCs. Bars represent mean  $\pm$  SD. **(D-H)**  $n = 8$ . Each dot represents the average of samples done in triplicate.

performed using a FACSCanto II flow cytometer (BD Biosciences). Data were analyzed using FlowJo software (TreeStar).

#### Statistical analysis


All data were presented as means  $\pm$  SD. Comparison between cohorts (healthy control [HC], EC, and CA) were performed using a two-tailed Mann-Whitney  $U$  test. This analysis is optimal for clinical cohorts, as it does not assume normal Gaussian distribution of the data. For statistical analyses within the same group using autologous or allogeneic cocultures (substitution of one cell type), we used paired, nonparametric Wilcoxon matched-pairs signed rank tests. For statistical analyses within the same group (i.e., comparison between Ab levels in cocultures from ECs with or without supplementation of IL-2 or IFN- $\gamma$ ) two-tailed paired  $t$  tests were used. Prism 5.0 (GraphPad) was used to process all the statistical analyses.

## Results

### Memory Tfh cells from CA individuals are functionally impaired

First, we characterized the frequency and phenotype of cTfh cells in peripheral blood from a cohort of HC, EC, and CA subjects (Tables I and II). Memory Tfh cells are Ag experienced ( $CD45RA^-$ ),

have mostly a central-memory phenotype ( $CCR7^+CD27^+$ ), and have stable expression of CXCR5. These cells have also been characterized as lacking the chemokine receptor CXCR3 (13, 14, 19). Unlike their GC counterparts, memory Tfh cells express high levels of IL-7R $\alpha$ , critical for long-term survival, and have a resting phenotype ( $CD69^-$ , Ki-67 $^-$ ). The majority of memory Tfh cells also lack CD38 and ICOS expression (Supplemental Fig. 1A-E), but a proportion of them express PD-1 (Supplemental Fig. 1F), as previously shown (14). When comparing the three patient groups, we did not observe any significant difference in the frequency and absolute number of cTfh cells (Supplemental Fig. 1G). In addition, we did not detect any significant difference in the expression of costimulatory (ICOS, CD40L), inhibitory (PD-1), or activation (CD38) markers following stimulation with anti-CD3/CD28 (Supplemental Fig. 1H). These results suggest that under anti-CD3/CD28 stimulation, at least for the markers studied, there are no apparent differences in the phenotype of memory Tfh cells between HC, EC, and CA subjects. However, the functional aspects of cTfh cells are better investigated in the context of B cell interaction



**FIGURE 2.** Altered IL-2R $\alpha$  and CXCR5 expression in cTfh cells from CA subjects. PBMCs from EC or CA subjects were sorted into cTfh cells and memory B cells and placed in coculture. After 7 d, cTfh cells were resorted for quantitative real-time PCR analysis using the Fluidigm BioMark HD System. Day 0 samples were used for normalization. **(A)** Fold change in the relative expression of genes, as assessed by real-time PCR analysis with respect to measurements at day 0. Results from five EC and four CA subjects are shown. Each dot represents the average of two biological replicates with four to six experimental replicates each. Bars represent mean  $\pm$ SD. **(B–F)** PBMCs were sorted as above and placed in autologous cocultures for 7 d or alternatively placed in coculture with either cTfh cells or memory B cells from a sorted pool of HCs instead of their own autologous counterparts. **(B)** Frequency of CD25 expression on cTfh cells from autologous cocultures or when these cells were replaced in cocultures from CA subjects with cTfh cells from HCs. **(C)** CXCR5 MFI on cTfh. **(D)** Correlation between the frequency of CD25 expression and CXCR5 MFI on cTfh cells. **(E)** Correlation between CD25 expression on cTfh cells and total IgG Ab levels in the coculture. **(F)** Correlation between CXCR5 MFI on cTfh cells and IgG levels.


and crosstalk. To accomplish this, we performed coculture assays with sorted memory Tfh cells (defined as CXCR5<sup>+</sup>CXCR3<sup>-</sup> CD4<sup>+</sup> memory T cells) and sorted memory B cells (defined as CD3<sup>-</sup>CD19<sup>+</sup> CD27<sup>+</sup>). As a control, we also cocultured non-Tfh cells (defined as CXCR5<sup>+</sup>CXCR3<sup>+</sup> or CXCR5<sup>-</sup> CD4<sup>+</sup> memory T cells) with memory B cells.

Similar to their GC counterparts, cTfh cells are specialized in providing B cell help. This characteristic was evidenced by both Ab output (Fig. 1A) and the absolute number of proliferating B cells after 7 d in coculture (Supplemental Fig. 2A). CXCR3<sup>+</sup> cells did not provide adequate B cell help, despite having a similar proliferative capacity to cTfh cells (Fig. 1A, Supplemental Fig. 2B), corroborating previous findings (13). In addition, activation of Tfh cells (using SEB) was required for their adequate function (Supplemental Fig. 2C). The absolute number of live B cells in the coculture was positively correlated with the total level of IgG (data not shown), demonstrating a relationship between cTfh-mediated B cell help and Ab production. From these results, it is evident that cTfh cells defined as CXCR5<sup>+</sup>CXCR3<sup>-</sup> CD4<sup>+</sup> memory T cells provide optimal help to memory B cells.

Using this coculture assay on 17 HC, 16 EC, and 24 CA individuals, we found that cocultures from CA subjects showed

significantly lower levels of total IgG Ab output when compared with cocultures from ECs and HCs (Fig. 1B,  $p < 0.001$ ). The differences in IgG levels were not due to a decrease in the absolute number of live B cells (Fig. 1C) or cTfh cells, as the absolute number of these cells was comparable between EC and CA subjects (Supplemental Fig. 2D). This finding suggests that the blunted IgG Ab levels could be due to defective cTfh/B cell interaction, which could influence memory B cell differentiation into IgG-secreting cells. To investigate this possibility further, we determined if we could restore Ab output by exchanging cTfh cells from CAs with cTfh cells from HC subjects, while maintaining the same memory B cells from CA individuals. Our results showed that replacing cTfh cells from CA subjects with cTfh cells from HCs significantly increased total IgG Ab output in cocultures with memory B cells from CA subjects (Fig. 1D,  $p < 0.01$ ). The restoration in Ab production was not due to an allogeneic effect, as memory B cells cocultured with allogeneic cTfh cells from HCs in the absence of stimulation (SEB) did not secrete detectable levels of IgG and showed poor viability (Supplemental Fig. 2G). Furthermore, in cocultures with memory B cells from ECs, replacing cTfh cells from ECs with cTfh cells from HCs did not enhance Ab production (Fig. 1E). In addition, the mean fluorescence intensity (MFI) of CD27, a

**FIGURE 3.** cTfh cells from CA subjects show increased IL-2 signaling. (A–C) PBMCs from EC or CA subjects were sorted into cTfh cells and memory B cells and placed in autologous cocultures for 7 d or alternatively placed in coculture with either cTfh cells or memory B cells from a sorted pool of HCs instead of their own autologous counterparts. All analyses were performed after 7 d in coculture. (A) Levels of IL-2 in cocultures after 7 d. (B) Correlation between the frequency of CD25 expression on cTfh cells and IL-2 levels in the coculture. (C) Correlation between IL-2 and the total levels of IgG in the cocultures. cTfh (CXCR5<sup>+</sup>CXCR3<sup>-</sup>), non-cTfh (CXCR5<sup>+</sup>CXCR3<sup>+</sup>), and memory B cells were also sorted from HC subjects and cocultured in the presence of SEB. (D) Total levels of IL-2 in the coculture supernatants after 7 d. Bars represent mean  $\pm$ SD.  $n = 8$  for (A),  $n = 16$  for (B) and (C),  $n = 5$  for (D). Each dot represents the average of samples done in triplicate.



marker highly expressed on Ab-secreting cells (ASCs), was significantly reduced in cocultures from CA subjects but restored when these cells were cultured with cTfh cells from HCs (Supplemental Fig. 2E, 2F,  $p < 0.001$ ). This observation suggests that memory B cells from CA subjects have a reduced capacity to differentiate into ASCs in the presence of impaired autologous cTfh cells.

For CA individuals, replacing autologous cTfh cells with those from HCs restored Ab production to the levels observed in ECs (Fig. 1F). This observation indicated that the memory B cell compartment in CA subjects had a similar intrinsic capacity to differentiate into ASCs when compared with ECs. This finding was further corroborated by coculturing memory B cells from CAs and ECs with cTfh cells from HCs, showing that memory B cells from both EC and CA subjects produced similar quantities of IgG (Fig. 1G).


The impairment in cTfh cell function in CA individuals became more apparent when we cocultured memory B cells from HCs with cTfh cells from either ECs or CA subjects. The results indicated that total IgG Ab output with cTfh cells from CA individuals was significantly decreased when compared with cocultures with cTfh cells from ECs (Fig. 1H,  $p < 0.02$ ). Together, these results suggest that the reduced Ab production observed in cocultures from CA subjects was primarily due to dysfunctional cTfh cells rather than intrinsic memory B cell defects.

#### Memory Tfh cells from CA subjects show increased IL-2 signaling

To continue characterizing the cTfh cell defect in CA subjects, we sorted cTfh cells after 7 d in coculture with memory B cells for Fluidigm BioMark analysis and compared their transcriptional gene fold change to sorted cTfh cells on day 0. We observed that some cTfh cells from cocultures of CA subjects showed a tendency toward increased IL-2R $\alpha$  expression and lower levels of BCL6, CXCR5 and IL21, canonical Tfh related genes essential for Tfh cell differentiation, trafficking, and B cell help when compared with cTfh cells from ECs (Fig. 2A). Even though the number of samples used in this assay was limited, the results suggested possible trends in the expression of some of these genes and made us pursue these differences further at the protein levels. Thus, we investigated expression of the high-affinity IL-2 receptor (IL-2R $\alpha$ /CD25) at the

protein level in cocultures from CA individuals and ECs. We observed a significant increase ( $p < 0.02$ ) in the frequency of CD25<sup>+</sup> cTfh cells in cocultures from CA individuals, whereas the expression levels of CXCR5 were significantly reduced ( $p < 0.001$ ) (Fig. 2B, 2C). The increase in CD25<sup>+</sup> cTfh cells was not due to signals coming from the memory B cells, as cocultures of memory B cells from CA subjects with cTfh cells from HCs did not result in an increase of CD25 expression on the cTfh cells (Fig. 2B). This finding suggests that the increase in CD25 expression was due to intrinsic properties in the cTfh cell compartment from CA individuals. In addition, we observed significant negative correlations between CD25 expression on cTfh cells and their expression level of CXCR5 and IgG Ab output in the cocultures (Fig. 2D, 2E,  $p < 0.001$  and  $p < 0.01$ , respectively). On the contrary, we observed a significant positive correlation between CXCR5 MFI on cTfh cells and total IgG Ab levels (Fig. 2F,  $p < 0.0001$ ). Together, these results imply that cTfh cells from CA subjects can display increased IL-2 signaling, which could be partly responsible for reprogramming these cells and altering their polarization and capacity to provide B cell help.

The expression of CD25 is regulated by TCR stimulation but can be sustained by IL-2 in a positive feedback loop involving STAT5 (20, 21). Indeed, we observed significantly higher levels of IL-2 in cocultures from CA subjects (Fig. 3A,  $p < 0.01$ ), and these levels were significantly reduced when exchanging cTfh cells from CA subjects with those from HCs. In addition, exchanging the memory B cells did not affect the levels of IL-2 in cocultures from CA subjects, indicating that the increased levels of IL-2 were coming from the cTfh cells (Fig. 3A). Furthermore, we observed a significant positive correlation between the levels of IL-2 and the expression of CD25 on cTfh cells (Fig. 3B,  $p < 0.03$ ). The increase in IL-2 was also negatively correlated with total IgG Ab output (Fig. 3C,  $p < 0.02$ ). These results suggest that the reprogramming of cTfh cells during chronic HIV infection could be partly the result of increased IL-2 signaling on these cells. A similar increase in the levels of IL-2 was also observed in coculture of non-cTfh cells (CXCR5<sup>+</sup>CXCR3<sup>+</sup>) and memory B cells when compared with cocultures from cTfh cells in HCs (Fig. 3D). This observation suggests that the cTfh cells in CA subjects are skewed toward a non-cTfh phenotype and increased IL-2 production.



**FIGURE 4.** cTfh cells from CA individuals are polarized toward a Th1 phenotype. (A–G) PBMCs from EC or CA subjects were sorted into cTfh cells and memory B cells and placed in autologous cocultures for 7 d or alternatively placed in cocultures with either cTfh cells or memory B cells from a sorted pool of HCs instead of their autologous counterparts. (A) IFN- $\gamma$  levels in the cocultures after 7 d. (B) Correlation between IFN- $\gamma$  and total IgG. (C) TNF- $\alpha$  levels in the cocultures after 7 d. (D) Correlation between TNF- $\alpha$  and total IgG. (E) Correlation between IL-2 and IFN- $\gamma$  levels. (F) Correlation between IL-2 and TNF- $\alpha$  levels. (G) Correlation between IFN- $\gamma$  and TNF- $\alpha$  levels. cTfh (CXCR5<sup>+</sup>CXCR3<sup>-</sup>), non-cTfh (CXCR5<sup>+</sup>CXCR3<sup>+</sup>), and memory B cells were also sorted from HCs and cocultured in the presence of SEB. (H) Total levels of IFN- $\gamma$  in the coculture supernatants after 7 d.  $n = 8$  for (A) and (C),  $n = 16$  for (B) and (D)–(G),  $n = 5$  for (H). Each dot represents the average of samples done in triplicate.


#### Memory Tfh cells from CA individuals are more polarized toward a Th1 phenotype

Triggering of IL-2R could activate downstream signals that could interfere with the STAT3-Bcl6 axis, skewing the polarization of memory Tfh cells. As evidence for this, we observed significantly higher levels ( $p < 0.001$ ) of the characteristic Th1 cytokine IFN- $\gamma$  (22, 23) in cocultures from CA individuals. Similar to what we observed for IL-2, the levels of IFN- $\gamma$  were also significantly reduced when cTfh cells from HCs were used in the cocultures (Fig. 4A,  $p < 0.01$ ). We also observed a significant negative correlation between IFN- $\gamma$  and total IgG levels (Fig. 4B,  $p < 0.001$ ). Moreover, we observed a significant increase in TNF- $\alpha$ , another characteristic Th1 cytokine in cocultures from CA subjects (Fig. 4C,  $p < 0.001$ ). As with IFN- $\gamma$ , the levels of TNF- $\alpha$  were significantly reduced by exchanging with cTfh cells from HCs (Fig. 4C,  $p < 0.01$ ). Furthermore, a significant negative correlation was found between the levels of TNF- $\alpha$  and total IgG (Fig. 4D,  $p < 0.002$ ). IL-2 showed a positive correlation with both IFN- $\gamma$  ( $p < 0.01$ ) and TNF- $\alpha$  ( $p < 0.0001$ ), and TNF- $\alpha$  showed a positive correlation ( $p < 0.0002$ ) with IFN- $\gamma$  (Fig. 4E–G). In addition, we also observed a similar increase in the levels of IFN- $\gamma$  when coculturing non-cTfh cells (CXCR5<sup>+</sup>CXCR3<sup>+</sup>) with mem-

ory B cells in samples from HCs (Fig. 4H), suggesting that cTfh cells in CA subjects appear to be more polarized toward a Th1 phenotype.

#### IL-2 can reprogram memory Tfh cell function

We then supplemented memory Tfh/B cell cocultures from ECs with IL-2 or IFN- $\gamma$  to determine whether we could recapitulate the functional defects observed with cTfh cells from CA subjects. As shown in Fig. 5A and 5B, IL-2 supplementation led to a significant decrease in total IgG Ab levels ( $p < 0.03$ ), whereas IFN- $\gamma$  reduced the levels of total IgG, but not to the same extent as observed with IL-2 supplementation. The decrease in Ab output in the presence of IL-2 was not due to differences in the absolute number of cTfh cells, as their numbers were similar (in the case of IFN- $\gamma$ ) or elevated (in the case of IL-2) when compared with the controls (Fig. 5C). Supplementation with IL-2 also increased the expression of CD25 ( $p < 0.01$ ) on cTfh cells while decreasing CXCR5 expression ( $p < 0.001$ ) (Fig. 5D, 5E). In addition, CD27 MFI was significantly decreased on B cells, suggesting reduced differentiation into ASCs (Fig. 5F,  $p < 0.01$ ). Importantly, IL-2 supplementation significantly increased the levels of the Th1 transcription factor Tbet (Fig. 5G,  $p < 0.01$ ) on cTfh cells as well as the transcription factor



**FIGURE 5.** IL-2 reprograms cT<sub>fh</sub> cells. (A–F) Cocultures of cT<sub>fh</sub> cells and memory B cells ( $1 \times 10^4$  each) from ECs were supplemented with either 10 ng/ml of IL-2 or 10 ng/ml of IFN- $\gamma$ . (A) Total levels of IgG in coculture supernatants from control samples and samples supplemented with exogenous IL-2, as measured by ELISA after 7 d. (B) Total levels of IgG in the presence or absence of IFN- $\gamma$ . (C) Absolute number of live cT<sub>fh</sub> cells after 7 d. Bars represent mean  $\pm$  SD. (D) Frequency of CD25 expression on cT<sub>fh</sub> cells after 7 d in coculture. (E) CXCR5 MFI on cT<sub>fh</sub> cells after 7 d in coculture. (F) CD27 MFI on B cells after 7 d.  $n = 5$ . Each dot represents the average of samples done in triplicate. (G) Fold change in the MFI of Tbet, as measured by intracellular staining (ICS) in control and IL-2-supplemented cocultures after 7 d ( $n = 5$ ). (H) Fold change in the MFI of Blimp1, as measured by ICS after 7 d in coculture ( $n = 5$ ). (I) Correlation between Tbet MFI and Blimp1 MFI, as measured by ICS in control and IL-2-supplemented cocultures after 7 d.


Blimp1, which antagonizes Bcl6 expression and T<sub>fh</sub> cell differentiation (Fig. 5H,  $p < 0.01$ ) (24). We also observed a positive correlation between Tbet and Blimp1 expression on these cells (Fig. 5I,  $p < 0.02$ ). Samples supplemented with IL-2 also showed increased levels of IFN- $\gamma$  and TNF- $\alpha$  production (Supplemental Fig. 3A, 3B). However, samples supplemented with IFN- $\gamma$  did not show an increase in IL-2 production or TNF- $\alpha$  levels (Supplemental Fig. 3C, 3D), suggesting that increased IL-2 signaling might be an important driver or maintainer of memory T<sub>fh</sub> cell polarization toward a Th1-like phenotype in CA subjects.

To determine whether the aberrant Th1 polarization on cT<sub>fh</sub> cells from CA subjects could be reversed, we interfered with IL-2 signaling in the coculture. This was achieved by adding a titrated amount of  $\alpha$ IL-2 Ab to the cocultures to reduce the heightened levels of IL-2 without completely blocking signaling through CD25, which is required for both cT<sub>fh</sub> and B cell survival (data not shown). Reducing the levels of IL-2 in cocultures from CA individuals led to an increase in total IgG Ab output (Fig. 6A,  $p < 0.01$ ) as well as decreased CD25 expression (Fig. 6B,  $p < 0.01$ ) and increased CXCR5 levels on cT<sub>fh</sub> cells (Fig. 6C,  $p < 0.01$ ). In addition, we observed a significant decrease in the expression of the transcription factor Tbet, which plays a major role in driving Th1 differentiation (Fig. 6D,  $p < 0.01$ ) as well as

Blimp1 (Fig. 6E,  $p < 0.02$ ), which can block Bcl6 expression and T<sub>fh</sub> cell differentiation (24). Furthermore, increased Tbet expression was significantly correlated (Fig. 6F,  $p < 0.01$ ) with decreased IgG levels as well as decreased CXCR5 expression on cT<sub>fh</sub> cells (Fig. 6G,  $p < 0.0001$ ). On the contrary, increased expression of Tbet on cT<sub>fh</sub> cells was significantly and positively correlated with increased expression levels of CD25 (Fig. 6H,  $p < 0.0001$ ) and Blimp1 (Fig. 6I,  $p < 0.0001$ ). Thus, the apparent Th1-like programming of cT<sub>fh</sub> cells in CA subjects can be partly reversed by dampening IL-2 signaling, leading to a more functional cT<sub>fh</sub> cell phenotype and an improved capacity to provide B cell help.

#### *HIV-specific Ab responses in CA subjects can be enhanced by adequate cT<sub>fh</sub> cell help*

We next determined whether we could enhance the production of HIV-specific Abs by providing adequate B cell help. We first determined whether HIV-specific memory B cells were present in our cohort of CA subjects, as HIV-infected individuals under ART show a continuous loss of HIV-specific memory B cells (17, 25–28). We used a fluorochrome-labeled gp140 probe to detect Ag-specific memory B cells by flow cytometry. We observed that, compared with ECs, there was a significant reduction in the frequency of




**FIGURE 6.** Interfering with IL-2 signaling enhances cTfh cell differentiation and function in CA individuals. (A–E) Cocultures of sorted cTfh cells and memory B cells ( $2 \times 10^4$  each) from CA individuals were supplemented with an isotype control or a titrated amount of anti-IL-2 Ab (100 ng/ml) on day 0 and day 3 to reduce the levels of IL-2 in the cocultures without completely inhibiting signaling through CD25. After 7 d, the supernatants and cells were collected for further analysis. (A) Total levels of IgG in coculture supernatants from control samples and samples supplemented with an anti-IL-2 Ab, as measured by ELISA after 7 d. (B) Frequency of CD25 expression on cTfh cells after 7 d in coculture. (C) CXCR5 MFI on cTfh cells after 7 d in coculture. (D) Fold change in the MFI of Tbet and (E) Blimp1, as measured by intracellular staining (ICS) in control and anti-IL-2 Ab-treated cocultures after 7 d ( $n = 9$ ). (F) Correlation between Tbet MFI and total IgG Ab output after 7 d. (G) Correlation between Tbet and CXCR5 expression levels, as measured by MFI. (H) Correlation between Tbet and CD25 expression levels, as measured by MFI. (I) Correlation between Tbet and Blimp1 expression levels, as measured by MFI.

gp140-specific memory B cells in CA individuals (Fig. 7A,  $p < 0.001$ ). We also observed a reduction in influenza-specific memory B cells, although the difference did not reach statistical significance (data not shown). We then measured HIV-specific Ab levels in cocultures from EC and CA subjects. As expected, we found significantly lower levels of HIV-specific Abs in cocultures from CA subjects, and the majority of these individuals did not have measurable HIV-specific Abs (Fig. 7B,  $p < 0.00001$ ). This decrease in HIV-specific responses from CA subjects was evident for Abs against gp140 ( $p < 0.001$ ) and gp41 ( $p < 0.001$ ) and the predominant IgG subclasses for anti-HIV responses: IgG1 ( $p < 0.01$ ) and IgG3 ( $p < 0.001$ ) (Supplemental Fig. 4). Reduced production of IL-21 by cTfh cells from CA subjects, as suggested by our BioMark data, could partly explain the reduced levels of HIV-specific IgG1 and IgG3 (29). Importantly, when we supplemented cocultures from ECs with IL-2 or IFN- $\gamma$ , we observed a significant reduction in HIV-specific Ab levels ( $p < 0.001$  and  $p < 0.02$ , respectively) that reached levels similar to those observed in cocultures from CA subjects (Fig. 7C, 7D). However, when cTfh cells from CA individuals were replaced in the cocultures for cTfh cells from ECs, the level of Env-specific IgG was increased (Fig. 7E,  $p < 0.01$ ). We also observed a significant increase in the level of Env-specific IgG when cocultures from CA subjects were supplemented with recombinant IL-21 (Fig. 7F,  $p <$

0.03). Thus, in the presence of functional cTfh cell help, Env-specific IgG levels are enhanced in cocultures from CA subjects.

## Discussion

In this study, we have shown that cTfh cells from CA individuals are functionally impaired in their ability to provide adequate B cell help when compared with memory Tfh cells from ECs and HCs. Memory Tfh cells from CA subjects showed increased IL-2 signaling as well as impaired differentiation and function. In accordance, studies in mice have shown a negative role of IL-2 signaling in mediating Tfh cell differentiation (4–6). Increased autocrine IL-2 levels could be leading to heightened STAT5 signaling, which has been shown to repress Bcl6 expression by upregulating Blimp-1 (4, 6, 24, 30). In fact, we observed that increased IL-2 signaling significantly raised the levels of Blimp1 (Fig. 5H), whereas interfering with IL-2 signaling reduced Blimp1 levels (Fig. 6E). Constitutive STAT5 signaling can selectively block Tfh cell differentiation and GC formation, and this has been shown to be largely mediated by IL-2 signaling (4). However, the relationship between phosphorylated STAT5 and STAT3 levels on these cells still needs to be investigated. Nevertheless, IL-2 could be playing an important role by antagonizing Tfh cell differentiation, and signals through the IL-2R (CD25) could be contributing to shifting



**FIGURE 7.** Decreased HIV-specific Ab responses in CA subjects can be enhanced by adequate cTfh cell help. Mature B cells from EC and CA individuals were stained ex vivo to quantify the frequency of gp140 (APC-gp140) or influenza-specific (PE-H1-CA09) memory B cells (see *Materials and Methods*). **(A)** Frequency of gp140 binding as a percentage of IgG<sup>+</sup> B cells. **(B–E)** PBMCs from HC, EC, or CA subjects were sorted into cTfh cells and memory B cells and placed in coculture for 7 d. **(B)** Total levels of Env-specific IgG after 7 d.  $n = 11, 15,$  and  $23$  for HC, EC, and CA, respectively. **(C)** Fold change in the levels of Env-specific IgG in cocultures from ECs in the presence or absence of 10 ng/ml of IL-2.  $n = 5$ . Also shown is the comparison between the levels of Env-specific IgG in cocultures from ECs with IL-2 supplementation with respect to cocultures from CA subjects. **(D)** Fold change in the levels of Env-specific IgG in cocultures from ECs in the presence or absence of 10 ng/ml of IFN- $\gamma$ .  $n = 5$ . Also shown is the comparison between the levels of Env-specific IgG in cocultures from ECs with IFN- $\gamma$  supplementation with respect to cocultures from CA subjects. Bars represent mean  $\pm$  SD. **(E)** Total levels of Env-specific IgG in cocultures from CA subjects with autologous cTfh cells or with cTfh cells from HCs.  $n = 8$ . Each dot represents the average of samples done in triplicate. **(F)** Total levels of Env-specific IgG, as measured by ELISA in cocultures from CA subjects after 7 d following supplementation with IL-21 (10 ng/ml).

the balance in favor of Th1-effector differentiation (4, 5, 31, 32). Recent evidence also suggests that defects in the IL-2 pathway leading to suboptimal IL-2 signaling could favor increased Tfh cell differentiation in the setting of type 1 diabetes (33). Increased autocrine IL-2 levels could therefore be affecting the differentiation of Tfh cells and could be partly responsible for the altered functional output in memory Tfh cells from CA subjects. In fact, this Tfh cell defect could contribute to the impaired vaccine responses reported in HIV-infected individuals. For example, it has been shown that the immune response to both hepatitis A and B vaccines seems to be reduced in HIV-infected subjects (34). In addition, the levels of Abs to tetanus toxoid and measles are significantly lower in HIV-infected individuals (35), and (36). Furthermore, it has been shown that some HIV-infected subjects have poor responses to influenza vaccination (37).

However, it is important to note that this increase in IL-2 signaling was not observed across all the CA samples studied and additional factors might also be contributing to the skewing of memory Tfh cell differentiation toward a Th1 phenotype. IFN- $\gamma$  and TNF- $\alpha$  could also be playing a role in altering and main-

taining a Th1-like polarization in memory Tfh cells from CA individuals as well as directly affecting B cell responses. We observed increased levels of both of these cytokines in cocultures from CA subjects as well as blunted B cell help in cocultures from ECs following addition of IFN- $\gamma$ . The production of these cytokines is likely the result of memory Tfh cell polarization toward a Th1-like phenotype. IFN- $\gamma$  signaling leads to increased levels of Tbet expression via induction of STAT1 (38). This activity could help tilt the balance in favor of Tbet expression and away from Bcl6. If the levels of Bcl6 go down, this would allow memory Tfh cells to become more polarized toward other effector phenotypes by allowing other transcription factors such as Tbet to outcompete Bcl6 signaling. Indeed, we observed that memory Tfh cells from CA subjects had increased Tbet expression as well as IFN- $\gamma$  and TNF- $\alpha$  production, molecular targets of Tbet.

Our multiplex Fluidigm BioMark analysis also revealed decreased levels of the canonical Tfh cell cytokine IL-21 in cTfh cells from cocultures of CA individuals when compared with ECs. Indeed, blocking IL-21 signaling in cocultures from HC subjects, using an IL-21R Fc chimera, significantly blunted Ab production

and the absolute number of live B cells (Supplemental Fig. 3E). Furthermore, supplementation of cocultures from CA subjects with IL-21 significantly increased total IgG Ab output (Supplemental Fig. 3E). The reduced expression of IL-21 in cTfh cells from CA subjects could be partly the result of increased IL-2 signaling on these cells, which could interfere with the STAT3-Bcl6 axis via STAT5 induction, but the overall functional defect in these cells is likely maintained by a combination of signals and altered transcription factor balance. In this sense, increased IL-2 signaling might be playing an important role in helping to interfere with Bcl6 function, which, together with IFN- $\gamma$  and TNF- $\alpha$ , could further help to drive the altered polarization of memory Tfh cells into a Th1-like phenotype in CA subjects.

How these memory Tfh cells acquire a Th1-like phenotype upon activation is unknown, but it is reasonable to speculate that memory Tfh cells in HIV-infected individuals under ART may continue to experience heightened levels of proinflammatory signals in LNs that, despite ART, can maintain residual continuous local inflammation (16) and viral replication (11). It has been shown that the Tfh cell subset in LNs is a preferential target for HIV infection (39). Because the phenotype of cTfh cells encompasses resting memory T cells, which harbor viral DNA, these cells could also contain viral DNA at extremely low frequencies (one in a million cells) (40, 41). Thus, at this very low frequency, it is possible that the polyclonal function of these cells could not be compromised by viral infection. To confirm this, we performed coculture assays in the presence or absence of ART and found no significant difference in cell viability and Ab production, suggesting that virus infection and replication, at low levels, do not contribute to Tfh dysfunction (data not shown).

Furthermore, it is possible that increased Blimp1 expression and IL-2R signaling could be driving these cells toward a regulatory T cell phenotype (42). However, we did not observe a significant frequency of FOXP3<sup>+</sup> cTfh cells in cocultures from CA subjects (data not shown), and the increased expression levels of IFN- $\gamma$  and TNF- $\alpha$  suggest that these cells are not acquiring a T follicular regulatory T cell/regulatory T cell-like phenotype.

It is also important to point out that in this study we investigated only the phenotype and function of cTfh cells in peripheral blood and not LN-resident Tfh cells, owing to the difficulty of obtaining lymphoid tissues from HIV-infected individuals and HCs. It is still unclear to what extent memory Tfh cells represent their LN counterparts. Memory Tfh cells could generate from pre-Tfh cells before these cells enter the GC or could be derived from GC Tfh cells. However, the exact identity of these cells is still under investigation. For this reason we cannot conclude that what we observe in cTfh cells is exactly what is occurring to Tfh cells in LNs, but by investigating these cells we can infer what general microenvironmental conditions were present in lymphoid tissues when these cells differentiated. This understanding provides a glimpse into the conditions experienced in LNs by both of these cell types, which could affect their overall differentiation and polarization. Our findings therefore provide an insight into the mechanisms responsible for the induction and maintenance of Tfh cell differentiation that could be important in vaccine development. Enhancing memory Tfh cell function could also help overcome various humoral immune defects; enhance Ab effector functions; and improve the affinity maturation of HIV-specific Abs in HIV-infected individuals, which could potentially lead to an improved control of viral replication.

## Acknowledgments

We thank Kim Kusser and Yu Shi from the Flow Cytometry Core at Vaccine and Gene Therapy Institute Florida for expertise and help in sorting all the samples; Will Williams and Nicole L. Yates for expertise with the

Ab subclass assays; Drs. H. Liao and B. Haynes for HIV-1 envelope proteins and mAbs; and all the patients who provided samples, without which this study would not have been possible.

## Disclosures

The authors have no financial conflicts of interest.

## References

- Crotty, S. 2011. Follicular helper CD4 T cells (TFH). *Annu. Rev. Immunol.* 29: 621–663.
- Victoria, G. D., and M. C. Nussenzweig. 2012. Germinal centers. *Annu. Rev. Immunol.* 30: 429–457.
- Crotty, S. 2014. T follicular helper cell differentiation, function, and roles in disease. *Immunity* 41: 529–542.
- Johnston, R. J., Y. S. Choi, J. A. Diamond, J. A. Yang, and S. Crotty. 2012. STAT5 is a potent negative regulator of TFH cell differentiation. *J. Exp. Med.* 209: 243–250.
- Ballesteros-Tato, A., B. León, B. A. Graf, A. Moquin, P. S. Adams, F. E. Lund, and T. D. Randall. 2012. Interleukin-2 inhibits germinal center formation by limiting T follicular helper cell differentiation. *Immunity* 36: 847–856.
- Nurieva, R. I., A. Podd, Y. Chen, A. M. Alekseev, M. Yu, X. Qi, H. Huang, R. Wen, J. Wang, H. S. Li, et al. 2012. STAT5 protein negatively regulates T follicular helper (Tfh) cell generation and function. *J. Biol. Chem.* 287: 11234–11239.
- INSIGHT-ESPRIT Study Group, SILCAAT Scientific Committee, D. Abrams, Y. Levy, M. H. Losso, A. Babiker, G. Collins, D. A. Cooper, J. Darbyshire, S. Emery, et al. 2009. Interleukin-2 therapy in patients with HIV infection. *N. Engl. J. Med.* 361: 1548–1559.
- Sereti, I., H. Imamichi, V. Natarajan, T. Imamichi, M. S. Ramchandani, Y. Badralmaa, S. C. Berg, J. A. Metcalf, B. K. Hahn, J. M. Shen, et al. 2005. In vivo expansion of CD4CD45RO-CD25 T cells expressing foxP3 in IL-2-treated HIV-infected patients. *J. Clin. Invest.* 115: 1839–1847.
- Biancotto, A., J. C. Grivel, S. J. Iglehart, C. Vanpouille, A. Lisco, S. F. Sieg, R. Debernardo, K. Garate, B. Rodriguez, L. B. Margolis, and M. M. Lederman. 2007. Abnormal activation and cytokine spectra in lymph nodes of people chronically infected with HIV-1. *Blood* 109: 4272–4279.
- Shive, C. L., J. C. Mudd, N. T. Funderburg, S. F. Sieg, B. Kyi, D. A. Bazdar, D. Mangioni, A. Gori, J. M. Jacobson, A. D. Brooks, J. Hardacre, J. Ammorio, J. D. Estes, T. W. Schacker, B. Rodriguez, and M. M. Lederman. 2014. Inflammatory cytokines drive CD4 T cell cycling and impaired responsiveness to interleukin-7: implications for immune failure in HIV disease. *J. Infect. Dis.* 210: 619–629.
- Fukazawa, Y., R. Lum, A. A. Okoye, H. Park, K. Matsuda, J. Y. Bae, S. I. Hagen, R. Shoemaker, C. Deleage, C. Lucero, et al. 2015. B cell follicle sanctuary permits persistent productive simian immunodeficiency virus infection in elite controllers. *Nat. Med.* 21: 132–139.
- Cubas, R. A., J. C. Mudd, A. L. Savoye, M. Perreau, J. van Grevenynghe, T. Metcalf, E. Connick, A. Meditz, G. J. Freeman, G. Abesada-Terk, Jr., et al. 2013. Inadequate T follicular cell help impairs B cell immunity during HIV infection. *Nat. Med.* 19: 494–499.
- Morita, R., N. Schmitt, S. E. Bentebibel, R. Ranganathan, L. Bourdery, G. Zurawski, E. Foucat, M. Dullaers, S. Oh, N. Sabzghabaei, et al. 2011. Human blood CXCR5(+)CD4(+) T cells are counterparts of T follicular cells and contain specific subsets that differentially support antibody secretion. *Immunity* 34: 108–121.
- Locci, M., C. Havenar-Daughton, E. Landais, J. Wu, M. A. Kroenke, C. L. Arlehamn, L. F. Su, R. Cubas, M. M. Davis, A. Sette, et al. International AIDS Vaccine Initiative Protocol C Principal Investigators. 2013. Human circulating PD-1+CXCR3-CXCR5+ memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody responses. *Immunity* 39: 758–769.
- Boswell, K. L., R. Paris, E. Boritz, D. Ambrozak, T. Yamamoto, S. Darko, K. Wloka, A. Wheatley, S. Narpara, A. McDermott, et al. 2014. Loss of circulating CD4 T cells with B cell helper function during chronic HIV infection. *PLoS Pathog.* 10: e1003853.
- Fletcher, C. V., K. Staskus, S. W. Wietgreffe, M. Rothenberger, C. Reilly, J. G. Chipman, G. J. Beilman, A. Khoruts, A. Thorkelson, T. E. Schmidt, et al. 2014. Persistent HIV-1 replication is associated with lower antiretroviral drug concentrations in lymphatic tissues. *Proc. Natl. Acad. Sci. USA* 111: 2307–2312.
- Goulet, M. L., D. Olgner, Z. Xu, S. Paz, S. M. Belnaoui, E. I. Lafferty, V. Janelle, M. Arguello, M. Paquet, K. Ghneim, et al. 2013. Systems analysis of a RIG-I agonist inducing broad spectrum inhibition of virus infectivity. *PLoS Pathog.* 9: e1003298.
- Kardava, L., S. Moir, N. Shah, W. Wang, R. Wilson, C. M. Buckner, B. H. Santich, L. J. Kim, E. E. Spurlin, A. K. Nelson, et al. 2014. Abnormal B cell memory subsets dominate HIV-specific responses in infected individuals. *J. Clin. Invest.* 124: 3252–3262.
- Schmitt, N., S. E. Bentebibel, and H. Ueno. 2014. Phenotype and functions of memory Tfh cells in human blood. *Trends Immunol.* 35: 436–442.
- Malek, T. R. 2008. The biology of interleukin-2. *Annu. Rev. Immunol.* 26: 453–479.
- Boyman, O., and J. Sprent. 2012. The role of interleukin-2 during homeostasis and activation of the immune system. *Nat. Rev. Immunol.* 12: 180–190.
- Chen, J., and X. Liu. 2009. The role of interferon gamma in regulation of CD4+ T-cells and its clinical implications. *Cell. Immunol.* 254: 85–90.
- Murphy, K. M., and S. L. Reiner. 2002. The lineage decisions of helper T cells. *Nat. Rev. Immunol.* 2: 933–944.

24. Johnston, R. J., A. C. Poholek, D. DiToro, I. Yusuf, D. Eto, B. Barnett, A. L. Dent, J. Craft, and S. Crotty. 2009. Bcl6 and Blimp-1 are reciprocal and antagonistic regulators of T follicular helper cell differentiation. *Science* 325: 1006–1010.
25. Morris, L., J. M. Binley, B. A. Clas, S. Bonhoeffer, T. P. Astill, R. Kost, A. Hurley, Y. Cao, M. Markowitz, D. D. Ho, and J. P. Moore. 1998. HIV-1 antigen-specific and -nonspecific B cell responses are sensitive to combination antiretroviral therapy. *J. Exp. Med.* 188: 233–245.
26. Bussmann, B. M., S. Reiche, B. Bieniek, I. Krznaric, F. Ackermann, and C. Jassoy. 2010. Loss of HIV-specific memory B-cells as a potential mechanism for the dysfunction of the humoral immune response against HIV. *Virology* 397: 7–13.
27. Moir, S., C. M. Buckner, J. Ho, W. Wang, J. Chen, A. J. Waldner, J. G. Posada, L. Kardava, M. A. O'Shea, S. Kottlilil, et al. 2010. B cells in early and chronic HIV infection: evidence for preservation of immune function associated with early initiation of antiretroviral therapy. *Blood* 116: 5571–5579.
28. Moir, S., and A. S. Fauci. 2014. B-cell exhaustion in HIV infection: the role of immune activation. *Curr. Opin. HIV AIDS* 9: 472–477.
29. Pène, J., J. F. Gauchat, S. Lécart, E. Drouet, P. Guglielmi, V. Boulay, A. Delwail, D. Foster, J. C. Lecron, and H. Yssel. 2004. Cutting edge: IL-21 is a switch factor for the production of IgG1 and IgG3 by human B cells. *J. Immunol.* 172: 5154–5157.
30. Crotty, S., R. J. Johnston, and S. P. Schoenberger. 2010. Effectors and memories: Bcl-6 and Blimp-1 in T and B lymphocyte differentiation. *Nat. Immunol.* 11: 114–120.
31. Choi, Y. S., R. Kageyama, D. Eto, T. C. Escobar, R. J. Johnston, L. Monticelli, C. Lao, and S. Crotty. 2011. ICOS receptor instructs T follicular helper cell versus effector cell differentiation via induction of the transcriptional repressor Bcl6. *Immunity* 34: 932–946.
32. Pepper, M., A. J. Pagán, B. Z. Igyártó, J. J. Taylor, and M. K. Jenkins. 2011. Opposing signals from the Bcl6 transcription factor and the interleukin-2 receptor generate T helper 1 central and effector memory cells. *Immunity* 35: 583–595.
33. Kenefick, R., C. J. Wang, T. Kapadi, L. Wardzinski, K. Attridge, L. E. Clough, F. Heuts, A. Kogimtzis, S. Patel, M. Rosenthal, et al. 2015. Follicular helper T cell signature in type 1 diabetes. *J. Clin. Invest.* 125: 292–303.
34. Sticchi, L., B. Bruzzone, P. Caligiuri, E. Rappazzo, M. Lo Casto, L. De Hoffer, G. Gustinetti, C. Viscoli, and A. Di Biagio. 2015. Seroprevalence and vaccination coverage of vaccine-preventable diseases in perinatally HIV-1-infected patients. *Hum. Vaccin. Immunother.* 11: 263–269.
35. Titanji, K., A. De Milito, A. Cagigi, R. Thorstensson, S. Grützmeier, A. Atlas, B. Hejdeman, F. P. Kroon, L. Lopalco, A. Nilsson, and F. Chiodi. 2006. Loss of memory B cells impairs maintenance of long-term serologic memory during HIV-1 infection. *Blood* 108: 1580–1587.
36. De Milito, A., A. Nilsson, K. Titanji, R. Thorstensson, E. Reizenstein, M. Narita, S. Grützmeier, A. Sönnberg, and F. Chiodi. 2004. Mechanisms of hypergammaglobulinemia and impaired antigen-specific humoral immunity in HIV-1 infection. *Blood* 103: 2180–2186.
37. Pallikkuth, S., A. Parmigiani, S. Y. Silva, V. K. George, M. Fischl, R. Pahwa, and S. Pahwa. 2012. Impaired peripheral blood T-follicular helper cell function in HIV-infected nonresponders to the 2009 H1N1/09 vaccine. *Blood* 120: 985–993.
38. Lighvani, A. A., D. M. Frucht, D. Jankovic, H. Yamane, J. Aliberti, B. D. Hissong, B. V. Nguyen, M. Gadina, A. Sher, W. E. Paul, and J. J. O'Shea. 2001. T-bet is rapidly induced by interferon-gamma in lymphoid and myeloid cells. *Proc. Natl. Acad. Sci. USA* 98: 15137–15142.
39. Perreau, M., A. L. Savoye, E. De Crignis, J. M. Corpataux, R. Cubas, E. K. Haddad, L. De Leval, C. Graziosi, and G. Pantaleo. 2013. Follicular helper T cells serve as the major CD4 T cell compartment for HIV-1 infection, replication, and production. *J. Exp. Med.* 210: 143–156.
40. Chomont, N., M. El-Far, P. Ancuta, L. Trautmann, F. A. Procopio, B. Yassine-Diab, G. Boucher, M. R. Boulassel, G. Ghattas, J. M. Brenchley, et al. 2009. HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. *Nat. Med.* 15: 893–900.
41. Finzi, D., M. Hermankova, T. Pierson, L. M. Carruth, C. Buck, R. E. Chaisson, T. C. Quinn, K. Chadwick, J. Margolick, R. Brookmeyer, et al. 1997. Identification of a reservoir for HIV-1 in patients on highly active antiretroviral therapy. *Science* 278: 1295–1300.
42. Cretney, E., A. Kallies, and S. L. Nutt. 2013. Differentiation and function of Foxp3(+) effector regulatory T cells. *Trends Immunol.* 34: 74–80.