

Family History of Early Infant Death Correlates with Earlier Age at Diagnosis But Not Shorter Time to Diagnosis for Severe Combined Immunodeficiency

Anderson Dik Wai Luk, Pamela P. Lee, Huawei Mao, Koon-Wing Chan, Xiang Chen, Tong-Xin Chen, Jian Xin He, Nadia Kechout, Deepti Suri, Yin Tao, et al.

► **To cite this version:**

Anderson Dik Wai Luk, Pamela P. Lee, Huawei Mao, Koon-Wing Chan, Xiang Chen, et al.. Family History of Early Infant Death Correlates with Earlier Age at Diagnosis But Not Shorter Time to Diagnosis for Severe Combined Immunodeficiency. *Frontiers in Immunology*, Frontiers, 2017, 8, pp.808 - 808. <10.3389/fimmu.2017.00808>. <pasteur-01638829>

HAL Id: pasteur-01638829

<https://hal-riip.archives-ouvertes.fr/pasteur-01638829>

Submitted on 20 Nov 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Family History of Early Infant Death Correlates with Earlier Age at Diagnosis But Not Shorter Time to Diagnosis for Severe Combined Immunodeficiency

OPEN ACCESS

Edited by:

Andrew Gennery,
Newcastle University,
United Kingdom

Reviewed by:

Jennifer Woodard Leiding,
University of South Florida,
United States
Helene Decaluwe,
Centre Hospitalier Universitaire
Sainte-Justine, Canada

*Correspondence:

Yu Lung Lau
lauylung@hku.hk

Specialty section:

This article was submitted to
Primary Immunodeficiencies,
a section of the journal
Frontiers in Immunology

Received: 18 April 2017

Accepted: 26 June 2017

Published: 12 July 2017

Citation:

Luk ADW, Lee PP, Mao H, Chan K-W,
Chen XY, Chen T-X, He JX,
Kechout N, Suri D, Tao YB, Xu YB,
Jiang LP, Liew WK,
Jirapongsananuruk O,
Daengsuwan T, Gupta A, Singh S,
Rawat A, Abdul Latiff AH, Lee ACW,
Shek LP, Nguyen TVA, Chin TJ,
Chien YH, Latiff ZA, Le TMH,
Le NNQ, Lee BW, Li Q, Raj D,
Barbouche M-R, Thong M-K,
Ang MCD, Wang XC, Xu CG, Yu HG,
Yu H-H, Lee TL, Yau FYS, Wong WH-S,
Tu W, Yang W, Chong PCY, Ho MHK
and Lau YL (2017) Family History of
Early Infant Death Correlates with
Earlier Age at Diagnosis But Not
Shorter Time to Diagnosis for Severe
Combined Immunodeficiency.
Front. Immunol. 8:808.
doi: 10.3389/fimmu.2017.00808

Anderson Dik Wai Luk¹, Pamela P. Lee¹, Huawei Mao^{1,2}, Koon-Wing Chan¹, Xiang Yuan Chen³, Tong-Xin Chen⁴, Jian Xin He⁵, Nadia Kechout⁶, Deepti Suri⁷, Yin Bo Tao³, Yong Bin Xu⁸, Li Ping Jiang⁹, Woei Kang Liew¹⁰, Orathai Jirapongsananuruk¹¹, Tassalapa Daengsuwan¹², Anju Gupta⁷, Surjit Singh⁷, Amit Rawat⁷, Amir Hamzah Abdul Latiff¹³, Anselm Chi Wai Lee¹⁴, Lynette P. Shek¹⁵, Thi Van Anh Nguyen¹⁶, Tek Jee Chin¹⁷, Yin Hsiu Chien¹⁸, Zarina Abdul Latiff¹⁹, Thi Minh Huong Le¹⁶, Nguyen Ngoc Quynh Le¹⁶, Bee Wah Lee¹⁵, Qiang Li²⁰, Dinesh Raj²¹, Mohamed-Ridha Barbouche²², Meow-Keong Thong²³, Maria Carmen D. Ang²⁴, Xiao Chuan Wang²⁵, Chen Guang Xu²⁶, Hai Guo Yu²⁷, Hsin-Hui Yu¹⁸, Tsz Leung Lee¹, Felix Yat Sun Yau²⁸, Wilfred Hing-Sang Wong¹, Wenwei Tu^{1,2}, Wangling Yang^{1,2}, Patrick Chun Yin Chong¹, Marco Hok Kung Ho¹ and Yu Lung Lau^{1,2*}

¹LKS Faculty of Medicine, Department of Paediatrics and Adolescent Medicine, The University of Hong Kong, Hong Kong, Hong Kong, ²Shenzhen Primary Immunodeficiency Diagnostic and Therapeutic Laboratory, The University of Hong Kong-Shenzhen Hospital, Shenzhen, China, ³Guangzhou Children's Hospital, Guangzhou, China, ⁴Department of Allergy and Immunology, Shanghai Children's Medical Center, Shanghai Jiao Tong University School of Medicine, Shanghai, China, ⁵Beijing Children's Hospital, Capital Medical University, Beijing, China, ⁶Institut Pasteur d'Algérie, Algeria, North Africa, ⁷Allergy Immunology Unit, Advanced Pediatrics Centre, Postgraduate Institute of Medical Education and Research, Chandigarh, India, ⁸Guang Zhou Women and Children's Medical Center, Guangzhou, China, ⁹Children's Hospital of Chongqing Medical University, Chongqing, China, ¹⁰KK Women's and Children's Hospital, Singapore, Singapore, ¹¹Department of Pediatrics, Siriraj Hospital Mahidol University, Bangkok, Thailand, ¹²Queen Sirikit National Institute of Child Health, Bangkok, Thailand, ¹³Monash University, Selangor, Malaysia, ¹⁴Mount Elizabeth Hospital, Singapore, Singapore, ¹⁵National University of Singapore, Singapore, Singapore, ¹⁶National Children's Hospital, Hanoi, Vietnam, ¹⁷Sarawak General Hospital Malaysia, Kuching, Malaysia, ¹⁸National Taiwan University Children's Hospital, Taipei, Taiwan, ¹⁹University Kebangsaan Malaysia Medical Center, Kuala Lumpur, Malaysia, ²⁰Sichuan Second West China Hospital, Sichuan, China, ²¹Department of Paediatrics, Holy Family Hospital, New Delhi, India, ²²Department of Immunology, Institut Pasteur de Tunis and University Tunis-El Manar, Tunis, Tunisia, ²³Faculty of Medicine, Department of Paediatrics, University of Malaya, Kuala Lumpur, Malaysia, ²⁴San Pedro Hospital, Davao, Philippines, ²⁵Children's Hospital of Fudan University, Shanghai, China, ²⁶The First Affiliated Hospital, Sun Yat-sen University, Guangzhou, China, ²⁷Nanjing Children's Hospital, Nanjing, China, ²⁸Queen Elizabeth Hospital, Hong Kong, Hong Kong

Background: Severe combined immunodeficiency (SCID) is fatal unless treated with hematopoietic stem cell transplant. Delay in diagnosis is common without newborn screening. Family history of infant death due to infection or known SCID (FH) has been associated with earlier diagnosis.

Objective: The aim of this study was to identify the clinical features that affect age at diagnosis (AD) and time to the diagnosis of SCID.

Methods: From 2005 to 2016, 147 SCID patients were referred to the Asian Primary Immunodeficiency Network. Patients with genetic diagnosis, age at presentation (AP), and AD were selected for study.

Results: A total of 88 different SCID gene mutations were identified in 94 patients, including 49 *IL2RG* mutations, 12 *RAG1* mutations, 8 *RAG2* mutations, 7 *JAK3* mutations, 4 *DCLRE1C* mutations, 4 *IL7R* mutations, 2 *RFXANK* mutations, and 2 *ADA* mutations. A total of 29 mutations were previously unreported. Eighty-three of the 94 patients fulfilled the selection criteria. Their median AD was 4 months, and the time to diagnosis was 2 months. The commonest SCID was X-linked ($n = 57$). A total of 29 patients had a positive FH. Candidiasis ($n = 27$) and bacillus Calmette–Guérin (BCG) vaccine infection ($n = 19$) were the commonest infections. The median age for candidiasis and BCG infection documented were 3 months and 4 months, respectively. The median absolute lymphocyte count (ALC) was $1.05 \times 10^9/L$ with over 88% patients below $3 \times 10^9/L$. Positive FH was associated with earlier AP by 1 month ($p = 0.002$) and diagnosis by 2 months ($p = 0.008$), but not shorter time to diagnosis ($p = 0.494$). Candidiasis was associated with later AD by 2 months ($p = 0.008$) and longer time to diagnosis by 0.55 months ($p = 0.003$). BCG infections were not associated with age or time to diagnosis.

Conclusion: FH was useful to aid earlier diagnosis but was overlooked by clinicians and not by parents. Similarly, typical clinical features of SCID were not recognized by clinicians to shorten the time to diagnosis. We suggest that lymphocyte subset should be performed for any infant with one or more of the following four clinical features: FH, candidiasis, BCG infections, and ALC below $3 \times 10^9/L$.

Keywords: severe combined immunodeficiency, family history, candidiasis, absolute lymphocyte count, newborn screening

INTRODUCTION

Severe combined immunodeficiency (SCID) is a group of genetic diseases causing profound developmental and functional impairment of T cells, affecting cellular and humoral immunities. Currently, at least 49 genes are identified to be responsible for SCID and its variants (1–3). Of all the SCID genes, the commonest gene involved is the IL-2 receptor gamma chain gene (*IL2RG*), which accounted for 45 and 19% of SCID cases before and after the T-cell receptor excision circle (TREC) newborn screening, and was introduced in USA (3–5). Patients typically present with recurrent infections from opportunistic pathogens and live-attenuated vaccines, such as bacillus Calmette–Guérin (BCG) (6), chronic diarrhea, and failure to thrive (FTT), eventually die within the first 2 years of life if left untreated (7). Patients typically have low absolute lymphocyte count (ALC). They have been classified by the number of

B lymphocytes as B+ or B– and recently by the causative genetic mutation.

The definitive treatment for SCID is hematopoietic stem cell transplant (HSCT). In addition, gene therapy serves as an alternative for X-linked and adenosine deaminase (ADA)-deficient SCID if suitable HSC donors are not available (8). SCID patients have a 94% survival rate if they undergo HSCT within the first 3.5 months of life (9). To facilitate timely HSCT, an early diagnosis must be made. However, delay in diagnosis is common due to the lack of awareness of the distinctive presenting features of SCID, such as recurrent and persistent opportunistic infections (2). To date, the only feature that is associated with an earlier diagnosis is a positive family history of infant death due to infection or known SCID in USA (10). In addition, family history of SCID is associated with earlier HSCT before 3.5 months (9, 11). Our present study aimed to identify the clinical features that could help clinicians diagnose SCID earlier by comparing the age and time to the diagnosis of patients with or without certain clinical features.

MATERIALS AND METHODS

Patient Source and Selection

The Asian Primary Immunodeficiency Network (APIN) is a primary immunodeficiency (PID) referral network established in 2009 by The University of Hong Kong as a platform for consultation and offering free genetic testing for suspected PID in over 70 centers in Asia and Africa. Its database stores clinical information

Abbreviations: ADA, adenosine deaminase gene; ALC, absolute lymphocyte count; AD, age at diagnosis; AP, age at presentation; APIN, Asian Primary Immunodeficiency Network; BCG, bacillus Calmette–Guérin; CMV, cytomegalovirus; *DCLRE1C*, DNA cross-link repair enzyme 1C (Artemis); FH, family history of early infant death; FN3, fibronectin type-III; FTT, failure to thrive; HSCT, hematopoietic stem cell transplant; ICU, intensive care unit; *IL2RG*, IL-2 receptor gamma chain; *IL7R*, IL-7 receptor alpha chain; *JAK 3*, janus kinase 3; PCP, *Pneumocystis jiroveci*; PID, primary immunodeficiency; *RAG1*, recombinase activating gene 1; *RAG2*, recombinase activating gene 2; *RFXANK*, regulatory factor X associated ankyrin containing protein; SCID, severe combined immunodeficiency; TREC, T-cell receptor excision circle.

provided by the referring doctors, laboratory results, and genetic test reports (12–14). From 2005 to 2016, 147 SCID patients were referred from 23 centers to the APIN for consultation and genetic testing, 42 of whom were reported in our previous study (13). In our study, we included patients with documented age at presentation (AP) and diagnosis. Among them, we selected patients with genetic diagnosis for identifying factors that affected age and time to diagnosis of SCID.

Data Collection

The referring doctors provided the clinical records of patients, together with the laboratory results. Demographic data, clinical presentation, and progress as well as investigation results including ALC and lymphocyte subsets of the patients were recorded. We only considered the clinical features and progression before the diagnosis of SCID was made. We defined AP as the age when the first clinical symptom was documented in the clinical record, age at diagnosis (AD) as the age when clinical diagnosis of SCID was made, and time to diagnosis as the duration between AP and AD. We only considered patients to have certain clinical feature if that feature was stated in the referral summary. We defined recurrent infections as more than one episode of infections affecting similar

systems. We considered the infection to be severe if at least one of the following was present: life-threatening complications (such as acute respiratory distress syndrome and sepsis), intensive care unit (ICU) admission, and life support being used (intubation, ventilation, and resuscitation). We defined opportunistic infection as an infection with at least one of the following pathogens was involved: BCG, *Candida*, *Pseudomonas aeruginosa*, *Acinetobacter baumannii*, *Pneumocystis jiroveci* (PCP), *Aspergillus*, cytomegalovirus (CMV), and herpes zoster virus. We defined opportunistic infections by *Candida* if patients were documented to have invasive candidiasis, candidemia or persistent oral candidiasis (15). We presented the ALC recorded at the time of SCID diagnosis. Patients were said to have lymphopenia if ALC below $3 \times 10^9/L$ as described previously (10). Since there is no universally agreed cutoff of B-cell number to distinguish B+ and B– SCID, we defined the cut-off as 134/ μL based on the CD19⁺ B-cell counts of patients with B+ genotypes (*IL2RG*, *IL7R*, and *JAK3*) and those with B– genotypes (*ADA*, *DCLRE1C*, *RAG1*, and *RAG2*) (Figure 1).

Genetic analysis was performed in the Department of Pediatrics and Adolescent Medicine of the University of Hong Kong using PCR and direct sequencing (Table SE1 in Supplementary Material) (13). Genetic and functional studies




FIGURE 1 | CD19⁺ cell counts of patients with B+ and B– genotypes. B+ genotypes group consisted of CD19⁺ cell counts of patients with mutations found in *IL2RG* ($n = 43$), *IL7R* ($n = 3$), and *JAK3* ($n = 5$). B– genotypes group consisted of CD19⁺ cell counts of patients with mutations found in *ADA* ($n = 1$), *DCLRE1C* ($n = 3$), *RAG1* ($n = 3$), and *RAG2* ($n = 2$). The cutoff for distinguishing B+ and B– patients was 134 CD19⁺ cells/ μL . Three patients with *IL2RG* mutations were classified as having B– SCID.

on PID, data archival in the APIN database, and DNA storage were approved by the Clinical Research Ethics Review Board of the University of Hong Kong and Queen Mary Hospital (Ref. no. UW 08-301) in accordance with the Declaration of Helsinki, with written informed consent obtained from parents of subjects. HGMD Pro version 2016.4 (16) and Immunodeficiency mutation databases (IDbases) (17) were used to identify unreported mutations. The nomenclatures of cDNA mutations were based on coding region. For each unreported mutation, the population frequency was analyzed by Exome Aggregation Consortium Browser (18). Effects of missense mutations on protein functions were predicted by PANTHER (19), PHD-SNP (20), SIFT (21), SNAP (22), Meta-SNP (23), and PolyPhen2 (24). The protein structure predicted to be involved was identified using NCBI Protein database (25) and UniProt Knowledgebase database (26).

Statistical Analysis

For descriptive statistics, all data were expressed in median and range (month). Univariate analysis was performed using Mann–Whitney *U* test; multivariate linear regression was performed for all factors that were significant ($p < 0.05$) in univariate analysis. We defined statistical significance as $p < 0.05$, and 95% confidence interval did not contain 0 in multivariate analysis. We did not include opportunistic infection group in the multivariate linear regression to avoid multicollinearity.

Patients with missing categorical data such as clinical features were considered to be without the features. Patients with missing numerical data such as ALC were not analyzed when analyzing median and range.

RESULTS

Patients Selection

From 147 SCID patients referred to the APIN, 131 of them had documented AP and diagnosis. Among these patients, 83 of them

had genetic diagnosis (Figure 2). Sixteen patients were excluded from the study due to the lack of AP ($n = 4$), the lack of AD ($n = 9$), and being diagnosed by screening ($n = 3$). Among the 16 patients excluded from the study, 11 of them had genetic diagnosis. Altogether, molecular diagnosis of SCID was identified in 94 patients in our cohort.

Genetic Mutations in Patients

The genetic mutations of the 83 SCID patients included in our study and 11 SCID patients excluded from our study are shown in Table 1 and Table SE1 in Supplementary Material, respectively. The commonest gene identified was *IL2RG* ($n = 65$), followed by *RAG1* ($n = 7$), *RAG2* ($n = 7$), *JAK3* ($n = 5$), *DCLRE1C* ($n = 4$), *IL7R* ($n = 3$), *RFXANK* ($n = 2$), and *ADA* ($n = 1$). Eighty-eight different mutations were identified in this study (49 *IL2RG* mutations, 12 *RAG1* mutations, 8 *RAG2* mutations, 7 *JAK3* mutations, 4 *DCLRE1C* mutations, 4 *IL7R* mutations, 2 *RFXANK* mutations, and 2 *ADA* mutations). There was no difference in clinical features between X-linked and autosomal recessive SCID patients (Table SE2 in Supplementary Material).

Genetic mutations in all SCID genes were not evenly distributed, and two mutations were seen three or more times in unrelated patients. c.854G>A mutation was seen in six unrelated patients with *IL2RG* mutation. c.104G>T mutation was observed in three unrelated patients with *RAG2* mutation.

Twenty-two C>T or G>A mutations within CpG dinucleotides were documented (8 *IL2RG* mutations, 5 *RAG1* mutations, 4 *RAG2* mutations, 3 *JAK3* mutations, 1 *IL7R* mutation, and 1 *ADA* mutation). These mutations accounted for 25% of all mutations and were involved in 31 patients (18 in *IL2RG*, 5 in *RAG1*, 3 in *RAG2*, 2 in *JAK3*, 1 in *IL7R*, and 1 in *ADA*).

There were 29 unreported mutations identified in our patients, including 23 *IL2RG* mutations, 3 *RAG1* mutations, 1 *JAK3* mutation, 1 *RAG2* mutation, and 1 *RFXANK* mutations (Table SE3 in Supplementary Material). Effects of these unreported mutations




FIGURE 2 | Patients selection algorithm in this study. From 147 SCID entries in the Asian Primary Immunodeficiency Network (APIN) database, 131 patients were included in our study and 16 patients were excluded from our study. Three patients were excluded as they were diagnosed by screening either antenatally or at birth. Thirteen patients were excluded due to the lack of age at presentation ($n = 4$) or the lack of age at diagnosis ($n = 9$). ¹Cordocentesis was performed due to positive family history of SCID, revealed low CD4⁺ count. ²Complete blood count, lymphocyte subsets, and immunoglobulins measurement were performed in one patient due to positive family history, revealed severe T- and B-cell lymphopenia and low serum IgA and IgM; newborn T-cell receptor excision circle (TREC) screening revealed 0 TREC copy in another patient.

TABLE 1 | Genetic mutations of SCID patients (*n* = 83).

No	Gene	Intron (I)/exon (E)	Nucleotide change	Predicted change
P001	<i>IL2RG</i>	E1	c.3G>T	M1I
P002	<i>IL2RG</i>	E2	c.127delA	T43fsX70
P003	<i>IL2RG</i>	E2	c.202G>T	E68X
P004a	<i>IL2RG</i>	E2	c.202G>A	E68K
P004b	<i>IL2RG</i>	E2	c.202G>A	E68K
P005	<i>IL2RG</i>	E2	c.202G>A	E68K
P006	<i>IL2RG</i>	E2	c.252C>A	N84K
P007	<i>IL2RG</i>	I2	g.IVS2-15A>G	Predicted aberrant splicing
P008	<i>IL2RG</i>	I2	g.IVS2-15A>G	Predicted aberrant splicing
P011 ^e	<i>IL2RG</i>	E3	c.310_311delinsG	H104fsX146
P012	<i>IL2RG</i>	E3	c.340G>T	G114C
P013 ^e	<i>IL2RG</i>	E3	c.359dupA	K120fsX167
P014	<i>IL2RG</i>	E3	c.362delA	E121fsX146
P015	<i>IL2RG</i>	E3	c.365T>C	I122T
P016	<i>IL2RG</i>	E3	c.365T>C	I122T
P017 ^e	<i>IL2RG</i>	E3	c.371T>C	L124P
P018	<i>IL2RG</i>	E3	c.376C>T	Q126X
P019 ^e	<i>IL2RG</i>	E3	c.376C>T	Q126X
P020	<i>IL2RG</i>	E3	c.383T>C	F128S
P021	<i>IL2RG</i>	E3	c.386T>A	V129D
P022	<i>IL2RG</i>	E3	c.406_415del	R136fsX143
P023 ^e	<i>IL2RG</i>	E3	c.421delC	Q141fsX146
P024	<i>IL2RG</i>	I3/E4 junction	g.IVS3-2A>T	Predicted aberrant splicing
P025 ^e	<i>IL2RG</i>	E4	c.507delG	Q169fsX170
P026	<i>IL2RG</i>	E4	c.507delG	Q169fsX170
P027 ^e	<i>IL2RG</i>	E4	c.562C>T	Q188X
P028	<i>IL2RG</i>	E4	c.562C>T	Q188X
P030	<i>IL2RG</i>	E5	c.658_659del	T220fsX227
P031	<i>IL2RG</i>	E5	c.664C>T	R222C
P032 ^e	<i>IL2RG</i>	E5	c.670C>T	R224W
P033	<i>IL2RG</i>	E5	c.670C>T	R224W
P034	<i>IL2RG</i>	E5	c.676C>T	R226C
P036 ^e	<i>IL2RG</i>	E5	c.677G>A	R226H
P037	<i>IL2RG</i>	E5	c.694G>C	G232R
P038	<i>IL2RG</i>	E5	c.709T>C	W237R
P039 ^e	<i>IL2RG</i>	E5	c.711G>A	W237X
P040 ^e	<i>IL2RG</i>	E5	c.722G>T	S241I
P041	<i>IL2RG</i>	E5	c.741delG	G247fsX272
P042	<i>IL2RG</i>	E5	c.741_742insG	E248fsX302
P044	<i>IL2RG</i>	E6	c.811G>T	G271X
P045 ^e	<i>IL2RG</i>	E6	c.835delG	V279fsX293
P046	<i>IL2RG</i>	E6/I6 junction	c.854G>T	R285L
P047 ^e	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P048 ^e	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P049 ^e	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P050 ^e	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P051	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P052	<i>IL2RG</i>	E6/I6 junction	c.854G>A ⁱ	Predicted aberrant splicing R285Q
P053 ^e	<i>IL2RG</i>	I6-I7	g.IVS6-72_IVS7-11del	Predicted exon 7 deletion
P055	<i>IL2RG</i>	I6/E7 junction	g.IVS6-2A>C	Predicted aberrant splicing
P056	<i>IL2RG</i>	I6	g.IVS6+3G>T	Predicted aberrant splicing
P057 ^e	<i>IL2RG</i>	I6	g.IVS6+5G>A	Predicted aberrant splicing
P058 ^e	<i>IL2RG</i>	I6	g.IVS6+5G>A	Predicted aberrant splicing
P059	<i>IL2RG</i>	E7	c.865C>T	R289X
P060	<i>IL2RG</i>	E8	c.929G>A	W310X
P062	<i>IL2RG</i>	E8	c.979_980delinsTT	E327L
P063a	<i>IL2RG</i>	E8	c.982C>T	R328X

(Continued)

TABLE 1 | Continued

No	Gene	Intron (I)/exon (E)	Nucleotide change	Predicted change
P064	<i>ADA</i>	E7 E11	c.646G>A c.1018_1019del	G216R K340fsX348
P065 ^b	<i>DCLRE1C</i>	E1-E3	Gross deletion	Gross deletion
P066 ^b	<i>DCLRE1C</i>	E1-E4	Gross deletion	Gross deletion
P068 ^e	<i>DCLRE1C</i>	I3/E4 junction	c.IVS3-1G>T Exon 4 skipped ^a	I83-G102del
P069 ^e	<i>IL7R</i>	E8 E1 E2/I2 junction	c.632G>T c.65G>T g.IVS2+2T>A	G211V S22I Predicted aberrant splicing
P070 ^{b,e}	<i>IL7R</i>	E5	c.562delC	L188X
P071 ^b	<i>IL7R</i>	E5	c.616C>T	R206X
P072	<i>JAK3</i>	E2	c.115dupC	Q39fsX51
P073	<i>JAK3</i>	E13 E3 E10	c.1744C>T c.307C>T c.1333C>T	R582W R103C R445X
P074 ^e	<i>JAK3</i>	E13	c.1763A>C	H588P
P075 ^{b,e}	<i>JAK3</i>	I14	g.IVS14-11G>A c.1914_1915insCCCCCTTAG ^a	638_639insPPX
P076 ^c	<i>JAK3</i>	E16	c.2062A>T	I688F
P077	<i>RAG1</i>	E2	c.994C>T	R332X
P078 ^e	<i>RAG1</i>	E2	c.3074dupT c.1178delG	L1025fsX1064 G393fsX402
P079	<i>RAG1</i>	E2	c.2095C>T	R699W
P080	<i>RAG1</i>	E2	c.1328G>A c.2486_2490del	R443K R829fsX869
P081 ^d	<i>RAG1</i>	E2	c.1681C>T	R561C
P083	<i>RAG1</i>	E2	c.2561G>A c.2005G>A	G854D E669K
P084 ^e	<i>RAG2</i>	E2 E1/I1 junction E2	c.2324T>A c.2918G>A c.-28G>C c.358delG	L775Q R973H Predicted aberrant splicing V120fsX130
P085 ^b	<i>RAG2</i>	E2	c.104G>T	G35V
P086 ^b	<i>RAG2</i>	E2	c.104G>T	G35V
P087	<i>RAG2</i>	E2	c.104G>T	G35V
P088 ^b	<i>RAG2</i>	E2	c.475C>T	R159C
P089	<i>RAG2</i>	E2	c.218G>A c.442C>T	R73H R148X
P091	<i>RFXANK</i>	E2 E3/I3 junction E5	c.685C>T g.IVS3+1delG c.299_300del	R229W Predicted aberrant splicing Q100fsX113
P092 ^b	<i>RFXANK</i>	E5	c.299_300del	Q100fsX113

ADA, adenosine deaminase; *DCLRE1C*, DNA cross-link repair enzyme 1C; *IL2RG*, interleukin-2 receptor subunit gamma; *IL7R*, interleukin-7 receptor subunit alpha; *JAK3*, Janus kinase 3; *RAG*, recombinase activating genes; *RFXANK*, regulatory factor X-associated ankyrin-containing protein.

^aFrom RT-PCR results.

^bHomozygous mutations.

^cP076 was a B+NK- patient with hypogammaglobulinemia (IgG 1.45 g/L, IgA 0.23 g/L, and IgM 0.26 g/L) whose mother was a heterozygous carrier.

^dOnly one mutation was found.

^ePatients reported in our previous study (13).

^fPrevious study reported 854G>A may cause R285Q or skipping of exon 6 (27). P004a and P004b and P063a and P063b (Table SE1 in Supplementary Material) were from the same kindred.

on protein functions were predicted by multiple tools and are shown in Table SE3 in Supplementary Material.

Characteristics of Patients That Fulfilled Selection Criteria ($n = 83$)

Characteristics of patients included in our study ($n = 131$) are shown in Tables 2–4. For patients that fulfilled selection criteria ($n = 83$), 88.0% were male ($n = 73$) and 75.9% were Chinese ($n = 63$). The median AP was 2 months (0.1–6 months), AD 4 months (0.5–18 months), and time to diagnosis 2 months

(0–14 months). Twenty-nine patients (34.9%) had a family history of early infant death (FH), among them one patient had a family history of SCID and one patient had a family history of PID. Parental consanguinity was present in four kindreds. The median ALC was $1.05 \times 10^9/L$ ($0.134\text{--}52.2 \times 10^9/L$, $n = 70$) with 88.6% below $3 \times 10^9/L$ ($n = 62$). The major immunophenotype was B+ SCID ($n = 51$) (Tables 2 and 3).

50.6% of patients presented with chronic diarrhea ($n = 42$) and 60.2% of patients recurrent infections ($n = 50$). The commonest site of infection was the respiratory system ($n = 61$), followed by

TABLE 2 | Characteristics of patients included in our study ($n = 131$) at SCID diagnosis.

	With genetic diagnosis	Without genetic diagnosis
	$n = 83$	$n = 48$
Gender	Number (%)	Number (%)
Male	73 (88.0)	33 (68.8)
Female	10 (12.0)	15 (31.3)
Ethnicity	Number (%)	Number (%)
Chinese	63 (75.9)	30 (62.5)
Southeast Asian	12 (14.5)	4 (8.3)
Indonesian	1 (1.2)	0 (0)
Malay	3 (3.6)	3 (6.3)
Philippino	1 (1.2)	0 (0)
Thai	5 (6.0)	1 (2.1)
Vietnamese	2 (2.4)	0 (0)
Indian	2 (2.4)	9 (18.8)
Algerian	5 (6.0)	0 (0)
Arabian	1 (1.2)	3 (6.3)
Australian	0 (0)	1 (2.1)
Korean	0 (0)	1 (2.1)
Positive family history	Number (%)	Number (%)
Early infant death	29 (34.9)	13 (27.1)
Consanguinity	4 (4.8)	1 (2.1)
Age in months	Median (range)	Median (range)
Age at presentation	2 (0.1–6)	2 (0–19)
Age at diagnosis	4 (0.5–18)	4 (0.1–27)
Time to diagnosis	2 (0–14)	2 (0–16)
SCID phenotype	Number (%)	Number (%)
B+	51 (61.4)	18 (37.5)
B–	15 (18.1)	24 (50.0)
Others	17 (20.5) ^a	6 (12.5) ^b
	Median (range)	Median (range)
Absolute lymphocyte count ($10^9/L$)	1.05 (0.134–52.2) ^c	0.77 (0.09–13.46) ^d

^aMaternal engraftment ($n = 1$), unknown ($n = 16$).^bUnknown ($n = 6$).^cIn 70 patients.^dIn 44 patients.**TABLE 3** | Lymphocyte subset for patients included in our study ($n = 131$).

Patient	Mutation gene	ALC ($\times 10^9/L$)	CD3 ⁺ cells/ μL (%)	CD19 ⁺ cells/ μL (%)	CD16/56 ⁺ cells/ μL (%)
B+ SCID					
P006	<i>IL2RG</i>	0.4	9.2 (2.3)	385.6 (96.4)	2.4 (0.6)
P008	<i>IL2RG</i>	0.95	0 (0)	931 (98)	9.5 (1)
P011	<i>IL2RG</i>	1.16	3.48 (0.3)	972 (83.8)	2.3 (0.2)
P013	<i>IL2RG</i>	0.31	0 (0)	270 (87)	0 (0)
P014	<i>IL2RG</i>	2.93	468 (16)	2,344 (80)	58.6 (2)
P015	<i>IL2RG</i>	0.51	0 (0)	459 (90)	10.2 (2)
P017	<i>IL2RG</i>	0.7	0.7 (0.1)	663 (94.7)	36.4 (5.2)
P018	<i>IL2RG</i>	1	20 (2)	890 (89)	0 (0)
P019	<i>IL2RG</i>	2.63	26.3 (1)	2,525 (96)	26.3 (1)
P020	<i>IL2RG</i>	0.38	22.8 (6)	345.8 (91)	0 (0)
P021 ^b	<i>IL2RG</i>	1.66	596 (35.9)	1,061 (63.9)	2 (0.12)
P022	<i>IL2RG</i>	0.33	7.59 (2.3)	313.5 (95)	6.6 (2)
P024 ^b	<i>IL2RG</i>	3.43	504 (14.7)	2,867 (83.6)	58.3 (1.7)
P025	<i>IL2RG</i>	1.4	0 (0)	1,302 (93)	42 (3)
P026	<i>IL2RG</i>	0.33	16.5 (5)	290.4 (88)	9.9 (3)
P027	<i>IL2RG</i>	1.1	5.5 (0.5)	1,022 (92.9)	14.3 (1.3)

(Continued)

gastrointestinal system ($n = 42$). 47.0% of infections were severe ($n = 39$).

Fifty patients developed opportunistic infection (60.2%). The commonest opportunistic infection was candidiasis ($n = 27$), followed by BCG infection ($n = 19$) and viral infection ($n = 9$). The median age for candidiasis documented was 3 months, the median age for BCG infection was 4 months, and the median age for CMV infection was 2.25 months.

For patients included in our study, clinical features were compared between those with ($n = 83$) and without genetic diagnosis ($n = 48$). Patients without genetic diagnosis had higher frequency of FTT (33.3 versus 15.7%, $p = 0.0189$) and CMV infections (25.0 versus 9.6%, $p = 0.0185$) (Table 4).

For patients with documented ALC ($n = 114$), 107 of them (93.9%) had at least one of the following four clinical features: FH, candidiasis, BCG infection, and ALC below $3 \times 10^9/L$. 65 of them (57.0%) had at least two of the four clinical features mentioned.

FH and Pneumonia Were Associated with Earlier AP

Factors that were found to significantly affect AP, AD, and time to diagnosis are shown in Tables 5 and 6.

In univariate analysis, FH, pneumonia, and recurrent infections were associated with earlier AP (FH by 1 month, $p = 0.002$; pneumonia by 1 month, $p = 0.003$; recurrent infections by 1 month, $p = 0.008$). Upon multivariate analysis, only FH and pneumonia were associated with earlier AP (FH by 0.884 month, $p = 0.005$; pneumonia by 0.863 month, $p = 0.009$).

FH Was Associated with Earlier AD

In univariate analysis, FH and CMV infections were associated with an earlier AD (FH by 2 months, $p = 0.008$; CMV by 1 month, $p = 0.025$). Upon multivariate analysis, only FH was associated with earlier AD (by 1.86 months, $p = 0.007$).

TABLE 3 | Continued

Patient	Mutation gene	ALC ($\times 10^9/L$)	CD3 ⁺ cells/ μL (%)	CD19 ⁺ cells/ μL (%)	CD16/56 ⁺ cells/ μL (%)
P028	<i>IL2RG</i>	5	4,600 (92)	300 (6)	100 (2)
P030	<i>IL2RG</i>	0.94	16 (1.7)	620.4 (66)	192.7 (20.5)
P031	<i>IL2RG</i>	5.1	948 (18.6)	3,042 (59.7)	928.7 (18.2)
P032	<i>IL2RG</i>	1.1	11 (1)	979 (89)	110 (10)
P033	<i>IL2RG</i>	0.99	5 (0.5)	585.1 (59.1)	17.8 (1.8)
P034	<i>IL2RG</i>	1.34	0 (0)	1,112 (83)	160.8 (12)
P036	<i>IL2RG</i>	1.11	11.1 (1)	455.1 (41)	577 (52)
P038	<i>IL2RG</i>	0.6	240 (40)	324 (54)	33 (5.5)
P039	<i>IL2RG</i>	1.72	0 (0)	1,170 (68)	498.8 (29)
P041	<i>IL2RG</i>	0.62	12.4 (2)	545.6 (88)	37.2 (6)
P044	<i>IL2RG</i>	0.9	0 (0)	846 (94)	18 (2)
P045	<i>IL2RG</i>	1.84	0 (0)	1,748 (95)	73.6 (4)
P047	<i>IL2RG</i>	1.41	155.1 (11)	1,197 (84.9)	18.3 (1.3)
P048	<i>IL2RG</i>	4.94	0 (0)	4,841 (98)	98.8 (2)
P049	<i>IL2RG</i>	2.1	0 (0)	2,079 (99)	0 (0)
P050	<i>IL2RG</i>	0.53	5.3 (1)	424 (80)	21.2 (4)
P051	<i>IL2RG</i>	1.1	0 (0)	1,067 (97)	11 (1)
P052	<i>IL2RG</i>	1.86	223.2 (12)	1,600 (86)	0 (0)
P053	<i>IL2RG</i>	1	52 (5.2)	892 (89.2)	N/A (N/A)
P055	<i>IL2RG</i>	0.9	9 (1)	828 (92)	9 (1)
P056	<i>IL2RG</i>	1.3	1 (0.08)	1,282 (98.6)	9.6 (0.74)
P058	<i>IL2RG</i>	1.3	0 (0)	611 (47)	18.2 (1.4)
P059	<i>IL2RG</i>	1.18	0 (0)	1,133 (96)	35.4 (3)
P060	<i>IL2RG</i>	1.5	45 (3)	1,350 (90)	30 (2)
P063a	<i>IL2RG</i>	0.62	61.4 (9.9)	484.2 (78.1)	46.5 (7.5)
P069	<i>IL7R</i>	1.89	183.3 (9.7)	1,111 (58.8)	565.1 (29.9)
P070	<i>IL7R</i>	1.21	147.6 (12.2)	756.3 (62.5)	410.2 (33.9)
P071	<i>IL7R</i>	0.785	1.6 (0.2)	148.4 (18.9)	433.3 (55.2)
P072	<i>JAK3</i>	2.52	1,738 (69)	730.8 (29)	0 (0)
P073	<i>JAK3</i>	0.47	4.7 (1)	437.1 (93)	N/A (N/A)
P074	<i>JAK3</i>	0.35	91 (26)	196 (56)	N/A (N/A)
P075	<i>JAK3</i>	0.49	6.4 (1.3)	266.1 (54.3)	19.6 (4)
P076	<i>JAK3</i>	1.5	12.6 (0.84)	1,377 (91.77)	54.5 (3.63)
P078	<i>RAG1</i>	7.64	3,965 (51.9)	267.4 (3.5)	3,705 (48.5)
P091	<i>RFXANK</i>	1.59	624.9 (39.3)	936.5 (58.9)	47.7 (3)
P094	N/A	2.06	195.7 (9.5)	1,788 (86.8)	76.2 (3.7)
P098	N/A	1.23	764.2 (62.13)	156.9 (12.76)	263.1 (21.39)
P099	N/A	13.46	9,826 (73)	1,346 (10)	1,750 (13)
P109	N/A	0.88	295.7 (33.6)	460.2 (52.3)	89.5 (10.2)
P110	N/A	2.06	68 (3.3)	1,593 (77.3)	345.7 (16.8)
P111	N/A	1.46	18.3 (1.25)	1,387 (95)	6.6 (0.45)
P112	N/A	2.42	217.8 (9)	1,500 (62)	532.4 (22)
P115	N/A	1.84	18.4 (1)	1,472 (80)	294.4 (16)
P116	N/A	1.3	26 (2)	1,040 (80)	130 (10)
P117	N/A	2.1	396.9 (18.9)	573.3 (27.3)	136.5 (6.5)
P120	N/A	1.09	21.8 (2)	1,030 (94.5)	3.3 (0.3)
P121	N/A	0.5	20 (4)	245 (49)	205 (41)
P122	N/A	0.41	32.8 (8)	278.8 (68)	86.1 (21)
P123 ^b	N/A	1.28	65.3 (5.1)	833.3 (65.1)	381.4 (29.8)
P124	N/A	0.8	15.2 (1.9)	724 (90.5)	43.2 (5.4)
P126	N/A	0.84	342.7 (40.8)	207.5 (24.7)	197.4 (23.5)
P128	N/A	3.5	2,485 (71)	455 (13)	455 (13)
P137	N/A	0.9	9.9 (1.1)	136.8 (15.2)	419.4 (46.6)
B- SCID					
P001	<i>IL2RG</i>	0.14	0 (0)	70 (50)	1.4 (1)
P002	<i>IL2RG</i>	0.67	636.5 (95)	13.4 (2)	0 (0)
P003	<i>IL2RG</i>	3.6	3,456 (96)	0 (0)	N/A (N/A)
P005	<i>IL2RG</i>	0.18	7.2 (4)	3.6 (2)	145.8 (81)
P064	<i>ADA</i>	0.21	4 (1.9)	1 (0.48)	16.8 (8)
P065	<i>DCLRE1C</i>	0.65	110.5 (17)	26 (4)	78 (12)
P066	<i>DCLRE1C</i>	1.2	36 (3)	24 (2)	1,080 (90)
P068	<i>DCLRE1C</i>	0.72	7.2 (1)	0.72 (0.1)	672.5 (93.4)

(Continued)

TABLE 3 | Continued

Patient	Mutation gene	ALC ($\times 10^9/L$)	CD3 ⁺ cells/ μL (%)	CD19 ⁺ cells/ μL (%)	CD16/56 ⁺ cells/ μL (%)
P079	<i>RAG1</i>	0.96	144 (15)	1.9 (0.2)	796.8 (83)
P080	<i>RAG1</i>	0.134	132 (98.6)	0.04 (0.03)	1.5 (1.1)
P083	<i>RAG1</i>	0.34	80.6 (23.7)	3.1 (0.9)	190.1 (55.9)
P084	<i>RAG2</i>	0.74	7.4 (1)	7.4 (1)	666 (90)
P087	<i>RAG2</i>	2.55	2.6 (0.1)	132.6 (5.2)	2,020 (79.2)
P088	<i>RAG2</i>	28.36	26,772 (94.4)	0 (0)	623.9 (2.2)
P092	<i>RFXANK</i>	1.019	276.1 (27.1)	19.4 (1.9)	25.5 (2.5)
P093	N/A	0.31	N/A (N/A)	2 (0.65)	120.9 (39)
P095	N/A	3.38	3,191 (94.4)	33.8 (1)	33.8 (1)
P097	N/A	2.49	2,366 (95)	18.9 (0.76)	49.8 (2)
P100	N/A	0.489	477.8 (97.7)	2.9 (0.6)	4.9 (1)
P101	N/A	0.242	15 (6.19)	2.9 (1.2)	206.4 (85.3)
P102	N/A	0.09	41.1 (45.7)	1.4 (1.6)	19.5 (21.7)
P103	N/A	1.8	1,499 (83.3)	3.6 (0.2)	257.4 (14.3)
P104	N/A	0.138	26.2 (19)	1.4 (1)	93.8 (68)
P105	N/A	0.72	144 (20)	0.72 (0.1)	537.1 (74.6)
P106	N/A	0.42	408.2 (97.2)	1.3 (0.3)	4.6 (1.1)
P107	N/A	0.65	76.7 (11.8)	29.3 (4.5)	490.8 (75.5)
P108	N/A	0.8	40 (5)	14.4 (1.8)	656 (82)
P114	N/A	0.59	11.8 (2)	15.9 (2.7)	472 (80)
P118	N/A	0.19	0 (0)	39.9 (21)	0.38 (0.2)
P125	N/A	0.84	579.6 (69)	100.8 (12)	134.4 (16)
P127	N/A	0.53	312.2 (58.9)	73.1 (13.8)	19.6 (3.7)
P130	N/A	0.29	70.8 (24.4)	45.8 (15.8)	150.8 (52)
P131	N/A	0.8	16 (2)	0 (0)	768 (96)
P132	N/A	0.74	583.1 (78.8)	17.8 (2.4)	96.2 (13)
P134	N/A	0.7	539 (77)	0 (0)	1.1 (0.16)
P135	N/A	1.03	20.6 (2)	30.9 (3)	875.5 (85)
P136	N/A	0.28	254.8 (91)	2.8 (1)	19.6 (7)
P138	N/A	0.22	72.6 (33)	2.2 (1)	129.8 (59)
P139	N/A	0.1	42.8 (42.8)	4.1 (4.1)	50 (50)
Others					
Maternal engraftment					
P077	<i>RAG1</i>	52.23	49,619 (95)	0 (0)	2,089 (4)
Unknown					
P004a	<i>IL2RG</i>	0.64	0 (0)	N/A ^a	N/A ^a
P004b	<i>IL2RG</i>	N/A	N/A (2)	N/A (85)	N/A (10)
P007	<i>IL2RG</i>	N/A	N/A (0.2)	N/A (86.7)	N/A (6.7)
P012	<i>IL2RG</i>	N/A	N/A (2)	N/A (95)	N/A (0)
P016	<i>IL2RG</i>	N/A	N/A (N/A)	N/A (N/A)	N/A (N/A)
P023	<i>IL2RG</i>	0.5	N/A (N/A)	N/A (N/A)	N/A (N/A)
P037	<i>IL2RG</i>	1.6	N/A (N/A)	N/A (N/A)	N/A (N/A)
P040	<i>IL2RG</i>	N/A	N/A (16)	N/A (82)	N/A (0)
P042	<i>IL2RG</i>	N/A	N/A (2)	N/A (95)	N/A (1)
P046	<i>IL2RG</i>	N/A	N/A (0)	N/A (93)	N/A (2)
P057	<i>IL2RG</i>	N/A	N/A (0)	N/A (89)	N/A (0)
P062	<i>IL2RG</i>	5.4	N/A (N/A)	N/A (N/A)	N/A (N/A)
P081	<i>RAG1</i>	N/A	N/A (1.5)	N/A (0.52)	N/A (74.1)
P085	<i>RAG2</i>	N/A	N/A (13)	N/A (0.1)	N/A (24)
P086	<i>RAG2</i>	N/A	N/A (0.67)	N/A (0)	N/A (76)
P089	<i>RAG2</i>	N/A	N/A (N/A)	N/A (N/A)	N/A (N/A)
P096	N/A	0.26	N/A (N/A)	N/A (N/A)	N/A (N/A)
P113	N/A	N/A	N/A (1)	N/A (87)	N/A (4)
P119	N/A	N/A	N/A (3)	N/A (67.5)	N/A (25.4)
P129	N/A	0.22	N/A (N/A)	N/A (N/A)	N/A (N/A)
P133	N/A	0.6	N/A (N/A)	N/A (N/A)	N/A (N/A)
P140	N/A	N/A	N/A (2.8)	N/A (0.6)	N/A (90)

B+ SCID was defined as having ≥ 134 CD19⁺ cells/ μL and B- SCID was defined as having < 134 CD19⁺ cells/ μL . ALC, absolute lymphocyte count; N/A, not available.

^aMedical record documented as "raised".

^bALC was not provided, derived by summation of CD3⁺ cells, CD19⁺ cells, and CD16/56⁺ cells, in P024 the ALC was documented as $2.7 \times 10^9/L$ in separate test.

TABLE 4 | Clinical features of patients included in our study ($n = 131$).

	With genetic diagnosis	Without genetic diagnosis
	$n = 83$	$n = 48$
	Number (%)	Number (%)
Classical SCID triad		
Failure to thrive	13 (15.7)	16 (33.3) ^a
Chronic diarrhea	42 (50.6)	27 (56.3)
Recurrent infections	50 (60.2)	28 (58.3)
Infection by systems		
Respiratory infection	61 (73.5) ^c	34 (70.8) ^d
Non-bacillus Calmette–Guérin (BCG) skin and soft tissue infection	7 (8.4)	11 (22.9)
Gastrointestinal infection	42 (50.6)	23 (48.9)
Urogenital infection	2 (2.4)	1 (2.1)
Musculoskeletal infection	3 (3.6)	1 (2.1)
Central nervous system infection	1 (1.2)	1 (2.1)
Sepsis	18 (21.7)	8 (16.7)
Severe infections		
Intensive care unit admission	24 (28.9)	14 (29.2)
Life support	26 (31.3)	14 (29.2)
Intubation and ventilation	21 (25.3)	8 (16.7)
Resuscitation and/or inotrope support	5 (6.0)	9 (18.8)
Life-threatening complication	40 (48.2)	18 (37.5)
Sepsis	18 (21.7)	8 (16.7)
Respiratory distress/failure	26 (31.3)	13 (27.1)
Acute heart failure	2 (2.4)	1 (1.2)
Opportunistic infections		
Bacterial	50 (60.2)	28 (58.3)
<i>Bacterium</i>	8 (9.6)	1 (2.1)
<i>Pseudomonas aeruginosa</i>	6 (7.2)	0 (0)
<i>Acinetobacter baumannii</i>	4 (4.8)	1 (2.1)
Viral	9 (10.8)	12 (25.0)
Cytomegalovirus (CMV)	8 (9.6) ^e	12 (25.0) ^b
Herpes zoster	1 (1.2)	0 (0)
Bacillus Calmette–Guérin (BCG) infection	19 (22.9) ^f	8 (16.7)
Local	8 (9.6)	2 (4.2)
Regional	2 (2.4)	1 (2.1)
Disseminated	9 (10.8)	5 (10.4)
Candidiasis	27 (32.5) ^g	16 (33.3)
Persistent oral thrush	22 (26.5)	10 (20.8)
Gastrointestinal tract infection	3 (3.6)	2 (4.2)
Candidemia	2 (2.4)	4 (8.3)
Fungal	3 (3.6)	2 (4.2)
<i>Pneumocystis jirovecii</i>	2 (2.4)	1 (2.1)
Aspergillosis	1 (1.2)	1 (2.1)
Hepatosplenomegaly	12 (14.5)	9 (18.8)

^a $p = 0.0189$.^b $p = 0.0185$.^cA total of 53 patients with genetic diagnosis had pneumonia (63.9%).^dA total of 32 patients without genetic diagnosis had pneumonia (66.7%).^eThe median age for CMV infection documented was 2.25 months ($n = 8$).^fThe median age for BCG infection documented was 4 months ($n = 14$).^gThe median age for candidiasis documented was 3 months ($n = 25$).

Candidiasis and Opportunistic Infections Were Associated with Later AD

In univariate analysis, candidiasis, FTT, opportunistic infections, and BCG infection were associated with a later AD (candidiasis by 2 months, $p = 0.008$; FTT by 3 months, $p = 0.038$; opportunistic infections by 1 month, $p = 0.018$; BCG by 2 months, $p = 0.005$). Upon multivariate analysis, only candidiasis was associated with later AD (by 2.21 months, $p = 0.002$).

TABLE 5 | Univariate analysis of features that affect age at presentation (AP), age at diagnosis, and time to diagnosis in patients fulfilled selection criteria ($n = 83$).

Features	Median AP (months) when		Difference in months (Group A–Group B)	p -Value
	Feature present (Group A)	Feature absent (Group B)		
FH	1	2	–1	0.002
Candidiasis	2	2	0	0.664
Bacillus Calmette–Guérin (BCG)	2	2	0	0.291
CMV	1	2	–1	0.280
FTT	2	2	0	0.954
Chronic diarrhea	2	2	0	0.778
Recurrent infections	2	3	–1	0.008
Severe infections	2	2	0	0.813
Pneumonia	2	3	–1	0.003
Hepatosplenomegaly	2.25	2	0.25	0.347
X-linked SCID	2	2	0	0.057
Low ALC ^a	2	2.25	–0.25	0.771
Features	Median age at diagnosis (months) when		Difference in months (Group C–Group D)	p -Value
	Feature present (Group C)	Feature absent (Group D)		
FH	3	5	–2	0.008
Candidiasis	6	4	2	0.008
BCG	6	4	2	0.005
CMV	3	4	–1	0.025
FTT	7	4	3	0.038
Chronic diarrhea	4	4.5	–0.5	0.949
Recurrent infections	5	4	1	0.241
Severe infections	4	5	–1	0.476
Pneumonia	4	5	–1	0.111
Hepatosplenomegaly	4	4	0	0.544
X-linked SCID	4	3.5	0.5	0.689
Low ALC ^a	4	6.5	–2.5	0.086
Features	Median time to diagnosis (months) when		Difference in months (Group E–Group F)	p -Value
	Feature present (Group E)	Feature absent (Group F)		
FH	2	2	0	0.494
Candidiasis	2.5	1.95	0.55	0.003
BCG	3	2	1	0.052
CMV	1.25	2	–0.75	0.155
FTT	4	2	2	0.104
Chronic diarrhea	1.15	2	–0.85	0.617
Recurrent infections	2.5	1	1.5	<0.001
Severe infections	2	2	0	0.565
Pneumonia	2	1.75	0.25	0.382
Hepatosplenomegaly	1.5	2	–0.5	0.217
X-linked SCID	2	2	0	0.569
Low ALC ^a	2	4.5	–2.5	0.124

^aDefined as ALC below $3 \times 10^9/L$.

FH, family history of early infant death; CMV, cytomegalovirus infection; FTT, failure to thrive, ALC, absolute lymphocyte count.

TABLE 6 | Multivariate linear regression of features that affect age at presentation (AP), age at diagnosis (AD), and time to diagnosis in patients fulfilled selection criteria ($n = 83$).

Features	Regression coefficient (months)	p-Value	95% CI
AP			
FH	-0.884	0.005	-1.499 to -0.269
Recurrent infections	-0.541	0.086	-1.161 to 0.078
Pneumonia	-0.863	0.009	-1.504 to -0.221
AD			
FH	-1.86	0.007	-3.189 to -0.529
Candidiasis	2.21	0.002	0.858 to 3.555
Bacillus Calmette-Guérin	1.11	0.141	-0.375 to 2.595
CMV	-1.58	0.147	-3.727 to 0.569
FTT	1.15	0.190	-0.584 to 2.886
Time to diagnosis			
Candidiasis	1.511	0.018	0.267 to 2.754
Recurrent infections	1.845	0.003	0.655 to 3.036

FH, family history of early infant death; CMV, cytomegalovirus infection; FTT, failure to thrive; 95% CI, 95% confidence interval.

Candidiasis, Opportunistic Infections, and Recurrent Infections Were Associated with Longer Time to Diagnosis

Candidiasis, opportunistic infections, and recurrent infections were shown to be associated with longer time to diagnosis (candidiasis by 0.55 month, $p = 0.003$; opportunistic infections by 1 month, $p = 0.005$; recurrent infections by 1.5 months, $p < 0.001$). Upon multivariate analysis, both candidiasis and recurrent infections were associated with longer time to diagnosis (candidiasis by 1.51 months, $p = 0.018$; recurrent infections by 1.85 months, $p = 0.003$).

Other Features Were Not Significantly Associated with AD and Time to Diagnosis

Analysis of chronic diarrhea, pneumonia, hepatosplenomegaly, severe infections, and lymphopenia revealed no association with AD and time to diagnosis. There was no difference between X-linked and autosomal recessive forms of SCID in AP, AD, and time to diagnosis.

DISCUSSION

We found family history of early infant death was associated with earlier AP and earlier AD but not shorter time to diagnosis. Therefore, the earlier AD could be due to the heightened alertness of family with such history so that medical attention was sought earlier, rather than prompting clinicians in making quicker SCID diagnosis. The association between positive family histories and earlier AD was reported by studies in USA and France (10, 28), but they did not investigate whether the positive family history shortened the time to diagnosis. Moreover, they did not investigate whether presence of family history of early infant death alone is associated with an earlier AD. Previous studies reported 16–60% of patients with positive family histories compared to that of 32% in our study; however, the definition of family history

differs between studies (Table 7). Our findings suggested that the family history of early infant death was valuable in alerting families but not clinicians who failed to recognize this clue as the time to diagnosis remained the same regardless of the presence of family history of early infant death.

We found candidiasis was associated with later AD and longer time to diagnosis. The median age of candidiasis documented was 3 months, and the median AD of SCID for patients with candidiasis was 6 months. Therefore, clinicians required 3 months to diagnose SCID after candidiasis was first documented. This suggested that candidiasis was an overlooked feature by clinicians in Asia. Other studies reported similar percentage at candidiasis in SCID patients but no report of association between candidiasis and AD (Table 7). Although oral candidiasis is relatively common in infants under 6 months old; however, persistent, recurrent, or invasive candidiasis warrants investigation for underlying immunodeficiencies in particular SCID (15). Our finding suggested that candidiasis may be useful as a clue for earlier diagnosis since the median age of candidiasis documented was 3 months, which was earlier than the optimal time for HSCT at 3.5 months (9, 11).

We were surprised to find that BCG infection was not associated with AD and time to diagnosis. This could be due to the relatively low frequency of patients with BCG infections (21%) identified in our study, which was at a lower frequency when compared to that of 45–57% reported previously (6, 29, 30). The population coverage, immunization schedules, and virulence of BCG in countries and regions included in our study were comparable to that in Brazil and Iran (Table SE5 in Supplementary Material) (29, 34–39); therefore, the above factors of BCG policies could not account for the discrepancy in the frequency of BCG complications between our study and that from Brazil and Iran. In addition, the onset of BCG complication in our study at 4 months old was comparable to that in Brazil at 3.7 months old (29). The median AD of SCID in our study was 4 months, which was earlier than the 8 months in Brazil and 5 months in Iran, suggesting that the lower frequency of BCG infections in our study (21%) than that in Brazil (57%) and Iran (45%) (Table 7) could be due to earlier diagnosis of SCID in our study.

The median age for BCG infection documented in our patients was 4 months, which was beyond the optimal time for HSCT. Our findings were in line with a previous report in which 74% of 349 BCG-vaccinated SCID patients developed BCG infection at or after 4 months of age (6). Therefore, despite BCG infections being useful clinical features of SCID as SCID patients have approximately 400-fold increase in risk of having localized BCG complication and 33,000-fold increase in risk of having disseminated complications (6), noticing BCG infection had little value in alerting clinicians to make a timely diagnosis of SCID for optimal HSCT, which should be before 3.5 months (9, 11).

We found that opportunistic infections were associated with later AD, while recurrent infections and opportunistic infections were associated with longer time to diagnosis. Therefore, such clinical features were likely the consequences of delay in the diagnosis of SCID, reflecting that clinicians in Asia were unable to recognize these as SCID features.

We found that pneumonia was associated with an earlier AP but did not affect AD and time to diagnosis. Therefore, parents

TABLE 7 | Comparison with previous SCID studies.

Cohort	n	Origin	Duration	AP (months)	AD (months)	Genotype (%)			ALC ($\times 10^9/L$)	Present in cases (%)		
						IL2RG	Other	Unknown		FH	Candidiasis	BCG infection
Our study	131	Asia	2005–2016	2 (0–19) ^a	4 (0.1–27) ^b	57 (44)	26 (20)	48 (37)	0.975 (0.09–52.20) ^a	42/131 (32) ⁱ	43/131 (33)	27/131 (21) ^b
Stephan et al. (28)	117	France	1970–1992	3 (0–19) ^c	4.6 (0–27) ^c	0 (0)	0 (0)	117 (100)	1.608 (0–30) ^c	70/117 (60)	33/117 (28)	10/28 (36) ^b
Mazzucchelli et al. (29)	70	Brazil	1996–2011	2 (0–19) ^a	8 (0–22) ^b	0 (0)	0 (0)	70 (100)	N/A	19/70 (27) ^b	29/64 (45)	39/69 (57) ^b
Yeganeh et al. (30)	40	Iran	1999–2007	2.26 (± 0.43) ^b	5 (± 0.67) ^b	0 (0)	0 (0)	40 (100)	1.26 ^a	20/40 (50) ^b	23/40 (58)	18/40 (45) ^b
Saleem et al. (31)	13	Pakistan	2006–2011	N/A	5 (1.23–8.93) ^a	0 (0)	0 (0)	13 (100)	0.41 (0.17–2.28) ^a	7/13 (54) ⁱ	N/A	N/A ^a
McWilliams et al. (10)	172	USA	1982–2013	N/A	4.87 (0–18) ^b	77 (45)	91 (53)	4 (2)	0.43 (± 1.28) ^b	63/172 (37) ^b	74/172 (43)	2/172 (1)
Dvorak et al. (32)	50	North America	2010–2012	N/A	1.13 (0–10.13) ^d	20 (40)	27 (54)	3 (6)	1.22 (0.02–10.72) ^a	12/50 (24)	8/50 (16)	0/0 (0)
Rozmus et al. (33)	40	Canada	2004–2010	N/A	4.2 (0–19.4) ^c	4 (10)	16 (40)	20 (50)	1.13 (0.05–14.04) ^a	20/39 (51)	8/39 (21)	0/0 (0)

AP, age at presentation; AD, age at diagnosis; ALC, absolute lymphocyte count; FH, positive family history.

^aMedian (range).

^bMean (SD).

^cMean (range).

^dTypical SCID ($n = 37$) only.

^eWith universal BCG vaccination at birth.

^fFH of early infant death.

^gSuggestive or confirmed FH of SCID.

^hFH of early infant death due to infection or known SCID.

ⁱFH of immunodeficiency.

^jFH of SCID and/or infant death and consanguinity.

may perceive pneumonia as a severe medical condition and then brought their children to seek medical care earlier. However, pneumonia also commonly affects children without SCID in Asia (40–42), and clinicians are not alerted to the possible diagnosis of SCID.

Chronic diarrhea, severe infections, and ALC below $3 \times 10^9/L$ were not associated with AD and time to diagnosis, likely due to the distributions of the AD and the time to diagnosis that were quite wide in patients with these features (Figure S1 in Supplementary Material). In addition, chronic diarrhea is common in Southeast Asia and Western Pacific region (40); thus, it may not be a useful differentiating feature for patients with SCID as compared to those without. CMV infections did not affect AD and time to diagnosis as they were documented in small number of patients in our study ($n = 8$). The low rate of documented CMV infections may be due to the lack of diagnostic capacity (43, 44).

This study presented the largest collection of SCID patients in China and Southeast Asia with 147 patients, including 94 SCID patients with genetic diagnosis. The median AD was 4 months, which was comparable to other cohorts in the world, given no newborn screening of TREC was performed (Table 7) (10, 28–33); however, it was later than the optimal time for HSCT.

The commonest SCID gene found to be mutated in our patients was *IL2RG* because of the low consanguinity rate in our population (45) as well as near absence of newborn screening in Asia. Mutations in *IL2RG* were unevenly distributed. Exons 3 and 5 of *IL2RG* were common sites for mutation, accounting for 45% of all *IL2RG* mutations (Table 1; Table SE1 in Supplementary Material) and 48% of all unreported *IL2RG* mutations (Table SE3 in Supplementary Material), which was comparable with previous study (46). Five mutation hotspots, namely cDNA 670, 676, 677, 854, and 865, were identified previously and accounted for 29% of all *IL2RG* mutations in one study (46). Mutations in these hotspots collectively accounted for 27% of *IL2RG* mutations in our study. Majority of the mutations in these hotspots involved either C>T or G>A mutations in CpG dinucleotides. The mutation frequency of the C nucleotides in CpGs is 10–50 times higher compared to any other bases (47). This is commonly thought to be due to the methylation and subsequently deamination of cytosine to form thymidine in CpG (48, 49). Apart from the mentioned hotspots, we identified 16 additional point mutations in all SCID genes involving such mechanism, suggesting that cytosine methylation and deamination to thymidine in CpG dinucleotide is a relatively common mechanism causing mutations in SCID genes.

Four patients with mutations in *IL2RG* were classified as having B– SCID with CD19+ B cells ranging from 0 to 70/ μL (Table 3). The four patients had typical SCID presentations (Table SE4 in Supplementary Material). One patient was screened for *DCLRE1C*, *RAG1*, and *RAG2* due to his B– phenotype, but no mutation found. Patients suffering from X-linked SCID but with T-B– phenotype have been described previously (28). Patients with documented *IL2RG* mutations but with T-B+*NK*+ phenotype were also described previously (50–52). Many possible mechanisms can lead to atypical SCID immunophenotypes, including concurrent mutations in other SCID genes, modifier gene(s), and mutations, that lead to sparing or disrupting developments of other lineages of lymphocytes. In addition, one

patient with mutations in *RAG1* was classified as having B+ SCID with 267.4 CD19+ B cell/ μ L (Table 3). He likely had either Omenn syndrome or maternal engraftment due to his T+B+NK+ immunophenotype as well as his clinical presentation of severe eczema and eosinophilia (Table 3; Table SE4 in Supplementary Material). The T+B+NK+ immunophenotype of this patient may be explained by his missense mutation (c.2095C>T; p.R699W). This hypomorphic mutation results in a mutant RAG1 enzyme with 19.3% residual recombinase activity (53), thus allowing the generation of B cells in this patient. These cases in our study as well as previous reports demonstrated the imperfect correlation of genotype–immunophenotype in SCID patients.

We reported two patients (P091 and P092) with *RFXANK* mutations, with one (P092) reported previously in Chinese literature (54). These patients were the only two with confirmed *RFXANK* mutations reported in Asia. *RFXANK* mutation causes bare lymphocyte syndrome type 2B (55), commonly observed in North Africa (56) and sometimes in other places such as France and Spain (57). MHC class II deficiency accounted for 32% of all forms of SCID and their variants in North Africa (58–62), while it accounted for 1.4% of all SCID in Asia. Such discrepancy could be explained by the higher consanguinity rate of 50% in North Africa (63) compared to that of less than 10% in Asia (45), as well as presence of founder mutations in North Africa (56, 64). These patients present with features of typical SCID and frequently with sclerosing cholangitis (56, 65). However, they have normal TREC level and cannot be identified by newborn screening (5, 66).

The care of patients with SCID in Asia is still at an early phase of development, as reflected by delay in diagnosis and suboptimal management with no easy access to HSCT (12, 14). Eighty-one of the 83 patients were the first member of their respective families to be diagnosed genetically with SCID, thus explaining the relative lack of family history of SCID in our study. For genetic counseling, we offered testing for family members of SCID patients including prenatal and newborn screening on siblings of six index patients as well as carrier screening for parents, siblings, and maternal aunts of 56 index patients (Table SE6 in Supplementary Material). Unfortunately, there is still a relative lack of clinical genetic service in Asia.

The median ALC in all studies including ours was below $3 \times 10^9/L$, and reaffirming lymphopenia is a feature commonly seen in SCID patients (Table 7). However, clinicians failed to act on this critical clue as lymphopenia did not affect the AD or time to diagnosis. Since 1994, many reports have emphasized the importance of low ALC in alerting clinicians regarding SCID (67–70), but sadly clinicians to this date still failed to appreciate the value of low ALC for the diagnosis of SCID.

In this study, we identified that FH, candidiasis, and ALC below $3 \times 10^9/L$ were overlooked clinical features prompting the diagnosis of SCID. In addition, BCG infections were useful clinical features as they were the second most common opportunistic infections in our SCID patients. Ninety-four percent of patients in our study had at least one of the following four features: FH, candidiasis, BCG infections, and ALC below $3 \times 10^9/L$. Therefore, we suggest a simple guideline mandating that clinicians should order lymphocyte subset analysis for infants with any one of the following four features: FH, candidiasis, BCG infections, and ALC below $3 \times 10^9/L$.

Failure to diagnose SCID in time will lead to delay in HSCT, leading to economic losses in addition to poor outcome. Study has shown that the mean total hospital charges in patients who had HSCT after 3.5 months were four times greater than those before 3.5 months (71). Since all the clinical features we analyzed failed to help clinicians in making earlier SCID diagnosis, newborn screening is the only solution for making early enough diagnosis of SCID for timely HSCT in Asia.

Our retrospective case-series relied on reports made by referring doctors instead of analyzing original charts and results; therefore, underreport of clinical features was possible. In addition, our handling of missing data tends to underestimate the strength of association of clinical features with AD and time to diagnosis.

In conclusion, clinicians failed to recognize typical clinical features of SCID to shorten the time to diagnosis. There is an urgent need to educate clinicians in Asia on SCID. Ultimately, the only solution for early diagnosis of and timely HSCT for patients with SCID is newborn screening.

ETHICS STATEMENT

Genetic and functional studies on PID, data archival in the APIN database, and DNA storage were approved by the Clinical Research Ethics Review Board of the University of Hong Kong and Queen Mary Hospital (Ref. no. UW 08-301).

AUTHOR CONTRIBUTIONS

ADL, PL, and YL designed the study; ADL and YL wrote the manuscript with extensive appraisal from PL, K-WC, and HM; YL, K-WC, WW, and ADL analyzed the data; K-WC, WY, and WT performed the genetic and immunological studies; PL, HM, YL, XC, T-XC, JH, NK, DS, YT, YX, LJ, WL, OJ, TD, AG, SS, AR, AHL, ACL, LS, TN, TC, YC, ZL, TML, NL, BL, QL, DR, M-RB, M-KT, MA, XW, CX, HY, H-HY, TLL, FY, PC, and MH contributed clinical data and took care of patients in this study. PL and YL established the Asian Primary Immunodeficiency Network (APIN).

FUNDING

This work was supported by the Hong Kong Society for Relief of Disabled Children; the Health and Medical Research Fund (01120846); and grant from Shenzhen Development and Reform Commission ([2015]164).

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://journal.frontiersin.org/article/10.3389/fimmu.2017.00808/full#supplementary-material>.

FIGURE S1 | Distribution of age at diagnosis and time to diagnosis of patients with genetic diagnosis ($n = 83$). Distribution of age at diagnosis of patients with chronic diarrhea (A), severe infections (B), and lymphopenia (C) and distribution of time to diagnosis of patients with chronic diarrhea (D) and lymphopenia (E).

REFERENCES

- Picard C, Al-Herz W, Bousfiha A, Casanova JL, Chatila T, Conley ME, et al. Primary immunodeficiency diseases: an update on the classification from the international union of immunological societies expert committee for primary immunodeficiency 2015. *J Clin Immunol* (2015) 35(8):696–726. doi:10.1007/s10875-015-0201-1
- Fischer A, Notarangelo LD, Neven B, Cavazzana M, Puck JM. Severe combined immunodeficiencies and related disorders. *Nat Rev Dis Primers* (2015) 1:15061. doi:10.1038/nrdp.2015.61
- Kwan A, Puck JM. History and current status of newborn screening for severe combined immunodeficiency. *Semin Perinatol* (2015) 39(3):194–205. doi:10.1053/j.semperi.2015.03.004
- Buckley RH, Schiff RI, Schiff SE, Markert ML, Williams LW, Harville TO, et al. Human severe combined immunodeficiency: genetic, phenotypic, and functional diversity in one hundred eight infants. *J Pediatr* (1997) 130(3):378–87. doi:10.1016/S0022-3476(97)70199-9
- Kwan A, Abraham RS, Currier R, Brower A, Andruszewski K, Abbott JK, et al. Newborn screening for severe combined immunodeficiency in 11 screening programs in the United States. *JAMA* (2014) 312(7):729–38. Erratum in: *JAMA* (2014) 312(20):2169. doi:10.1001/jama.2014.9132
- Marciano BE, Huang CY, Joshi G, Rezaei N, Carvalho BC, Allwood Z, et al. BCG vaccination in patients with severe combined immunodeficiency: complications, risks, and vaccination policies. *J Allergy Clin Immunol* (2014) 133(4):1134–41. Erratum in: *J Allergy Clin Immunol* (2014) 134(1):244. doi:10.1016/j.jaci.2014.02.028
- Sponzilli I, Notarangelo LD. Severe combined immunodeficiency (SCID): from molecular basis to clinical management. *Acta Biomed* (2011) 82(1):5–13.
- Gaspar HB, Qasim W, Davies EG, Rao K, Amrolia PJ, Veys P. How I treat severe combined immunodeficiency. *Blood* (2013) 122(23):3749–58. doi:10.1182/blood-2013-02-380105
- Pai SY, Logan BR, Griffith LM, Buckley RH, Parrott RE, Dvorak CC, et al. Transplantation outcomes for severe combined immunodeficiency, 2000–2009. *N Engl J Med* (2014) 371(5):434–46. doi:10.1056/NEJMoa1401177
- McWilliams LM, Dell Railey M, Buckley RH. Positive family history, infection, low absolute lymphocyte count (ALC), and absent thymic shadow: diagnostic clues for all molecular forms of severe combined immunodeficiency (SCID). *J Allergy Clin Immunol Pract* (2015) 3(4):585–91. doi:10.1016/j.jaip.2015.01.026
- Brown L, Xu-Bayford J, Allwood Z, Slatter M, Cant A, Davies EG, et al. Neonatal diagnosis of severe combined immunodeficiency leads to significantly improved survival outcome: the case for newborn screening. *Blood* (2011) 117(11):3243–6. doi:10.1182/blood-2010-08-300384
- Lee PP. *Primary Immunodeficiency Disorders in Southeast Asia: Needs, Priorities and Opportunities [MD Thesis]*. University of Hong Kong, Pokfulam, Hong Kong SAR (2014).
- Lee PP, Chan KW, Chen TX, Jiang LP, Wang XC, Zeng HS, et al. Molecular diagnosis of severe combined immunodeficiency – identification of IL2RG, JAK3, IL7R, DCLRE1C, RAG1, and RAG2 mutations in a cohort of Chinese and Southeast Asian children. *J Clin Immunol* (2011) 31(2):281–96. doi:10.1007/s10875-010-9489-z
- Lee PP, Lau YL. Improving care, education, and research: the Asian primary immunodeficiency network. *Ann N Y Acad Sci* (2011) 1238:33–41. doi:10.1111/j.1749-6632.2011.06225.x
- Krol DM, Keels MA. Oral conditions. *Pediatr Rev* (2007) 28(1):15–22. doi:10.1542/pir.28-1-15
- Stenson PD, Ball EV, Mort M, Phillips AD, Shiel JA, Thomas NS, et al. Human gene mutation database (HGMD): 2003 update. *Hum Mutat* (2003) 21(6):577–81. doi:10.1002/humu.10212
- Piirilä H, Väliäho J, Vihinen M. Immunodeficiency mutation databases (IDbases). *Hum Mutat* (2006) 27(12):1200–8. doi:10.1002/humu.20405
- Lek M, Karczewski KJ, Minikel EV, Samocha KE, Banks E, Fennell T, et al. Analysis of protein-coding genetic variation in 60,706 humans. *Nature* (2016) 536(7616):285–91. doi:10.1038/nature19057
- Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, et al. PANTHER version 11: expanded annotation data from gene ontology and reactome pathways, and data analysis tool enhancements. *Nucleic Acids Res* (2017) 45(D1):D183–9. doi:10.1093/nar/gkw1138
- Capriotti E, Calabrese R, Casadio R. Predicting the insurgence of human genetic diseases associated to single point protein mutations with support vector machines and evolutionary information. *Bioinformatics* (2006) 22(22):2729–34. doi:10.1093/bioinformatics/btl423
- Kumar P, Henikoff S, Ng PC. Predicting the effects of coding non-synonymous variants on protein function using the SIFT algorithm. *Nat Protoc* (2009) 4(7):1073–81. doi:10.1038/nprot.2009.86
- Johnson AD, Handsaker RE, Pulit SL, Nizzari MM, O'Donnell CJ, de Bakker PI. SNAP: a web-based tool for identification and annotation of proxy SNPs using HapMap. *Bioinformatics* (2008) 24(24):2938–9. doi:10.1093/bioinformatics/btn564
- Capriotti E, Altman RB, Bromberg Y. Collective judgment predicts disease-associated single nucleotide variants. *BMC Genomics* (2013) 14(Suppl 3):S2. doi:10.1186/1471-2164-14-S3-S2
- Adzhubei IA, Schmidt S, Peshkin L, Ramensky VE, Gerasimova A, Bork P, et al. A method and server for predicting damaging missense mutations. *Nat Methods* (2010) 7(4):248–9. doi:10.1038/nmeth0410-248
- NCBI Resource Coordinators. Database resources of the national center for biotechnology information. *Nucleic Acids Res* (2016) 44:D7–D19. doi:10.1093/nar/gkv1290
- UniProt Consortium. UniProt: a hub for protein information. *Nucleic Acids Res* (2015) 43(Database issue):D204–12. doi:10.1093/nar/gku989
- Kanai N, Yanai F, Hirose S, Nibu K, Izuhara K, Tani T, et al. A G to A transition at the last nucleotide of exon 6 of the gamma c gene (868G → A) may result in either a splice or missense mutation in patients with X-linked severe combined immunodeficiency. *Hum Genet* (1999) 104(1):36–42. Erratum in: *Hum Genet* (1999) 104(2):196. doi:10.1007/s004390050907
- Stephan JL, Vlekova V, Le Deist F, Blanche S, Donadieu J, De Saint-Basile G, et al. Severe combined immunodeficiency: a retrospective single-center study of clinical presentation and outcome in 117 patients. *J Pediatr* (1993) 123(4):564–72. doi:10.1016/S0022-3476(05)80951-5
- Mazzucchelli JT, Bonfim C, Castro GG, Condino-Neto AA, Costa NM, Cunha L, et al. Severe combined immunodeficiency in Brazil: management, prognosis, and BCG-associated complications. *J Investig Allergol Clin Immunol* (2014) 24(3):184–91.
- Yeganeh M, Heidarzade M, Pourpak Z, Parvaneh N, Rezaei N, Gharagozlu M, et al. Severe combined immunodeficiency: a cohort of 40 patients. *Pediatr Allergy Immunol* (2008) 19(4):303–6. doi:10.1111/j.1399-3038.2007.00647.x
- Saleem AF, Ali Khawaja RD, Shaikh AS, Ali SA, Mehdi Zaidi AK. Severe combined immune deficiency syndrome. *J Coll Physicians Surg Pak* (2013) 23(8):570–3.
- Dvorak CC, Cowan MJ, Logan BR, Notarangelo LD, Griffith LM, Puck JM, et al. The natural history of children with severe combined immunodeficiency: baseline features of the first fifty patients of the primary immune deficiency treatment consortium prospective study 6901. *J Clin Immunol* (2013) 33(7):1156–64. doi:10.1007/s10875-013-9917-y
- Rozmus J, Junker A, Thibodeau ML, Grenier D, Turvey SE, Yacoub W, et al. Severe combined immunodeficiency (SCID) in Canadian children: a national surveillance study. *J Clin Immunol* (2013) 33(8):1310–6. doi:10.1007/s10875-013-9952-8
- Zwerling A, Behr MA, Verma A, Brewer TF, Menzies D, Pai M. The BCG world atlas: a database of global BCG vaccination policies and practices. *PLoS Med* (2011) 8(3):e1001012. doi:10.1371/journal.pmed.1001012
- WHO. *UNICEF: Immunization Summary, a Statistical Reference Containing Data through 2013*. (2014). Available from: http://www.who.int/immunization/monitoring_surveillance/Immunization_Summary_2013.pdf
- Joung SM, Ryou S. BCG vaccine in Korea. *Clin Exp Vaccine Res* (2013) 2(2):83–91. doi:10.7774/cevr.2013.2.2.83
- Wang JF, Dai FY, Gong XL, Bao L. Commonly administered bacilli Calmette-Guerin strains induce comparable immune response. *Int J Clin Exp Med* (2015) 8(9):15834–9.
- Govindarajan KK, Chai FY. BCG adenitis-need for increased awareness. *Malays J Med Sci* (2011) 18(2):66–9.
- Zhang L, Ru HW, Chen FZ, Jin CY, Sun RF, Fan XY, et al. Variable virulence and efficacy of BCG vaccine strains in mice and correlation with genome polymorphisms. *Mol Ther* (2016) 24(2):398–405. doi:10.1038/mt.2015.216

40. Walker CL, Rudan I, Liu L, Nair H, Theodoratou E, Bhutta ZA, et al. Global burden of childhood pneumonia and diarrhoea. *Lancet* (2013) 381(9875):1405–16. doi:10.1016/S0140-6736(13)60222-6
41. Rudan I, Boschi-Pinto C, Biloglav Z, Mulholland K, Campbell H. Epidemiology and etiology of childhood pneumonia. *Bull World Health Organ* (2008) 86(5):408–16. doi:10.2471/BLT.07.048769
42. Gereige RS, Laufer PM. Pneumonia. *Pediatr Rev* (2013) 34(10):438–56; quiz 455–6. Erratum in: *Pediatr Rev* (2014) 35(1):29. doi:10.1542/pir.34-10-438
43. Plosa EJ, Esbenshade JC, Fuller MP, Weitkamp JH. Cytomegalovirus infection. *Pediatr Rev* (2012) 33(4):156–63; quiz 163. doi:10.1542/pir.33-4-156
44. Eguchi H, Horita N, Ushio R, Kato I, Nakajima Y, Ota E, et al. Diagnostic test accuracy of antigenemia assay for polymerase chain reaction proven cytomegalovirus infection: systematic review and meta-analysis. *Clin Microbiol Infect* (2017). doi:10.1016/j.cmi.2017.05.009
45. Hamamy H. Consanguineous marriages: preconception consultation in primary health care settings. *J Community Genet* (2012) 3(3):185–92. doi:10.1007/s12687-011-0072-y
46. Puck JM, Pepper AE, Henthorn PS, Candotti F, Isakov J, Whitwam T, et al. Mutation analysis of IL2RG in human X-linked severe combined immunodeficiency. *Blood* (1997) 89(6):1968–77.
47. Walser JC, Furano AV. The mutational spectrum of non-CpG DNA varies with CpG content. *Genome Res* (2010) 20(7):875–82. doi:10.1101/gr.103283.109
48. Wang RY, Kuo KC, Gehrke CW, Huang LH, Ehrlich M. Heat- and alkali-induced deamination of 5-methylcytosine and cytosine residues in DNA. *Biochim Biophys Acta* (1982) 697(3):371–7. doi:10.1016/0167-4781(82)90101-4
49. Ehrlich M, Zhang XY, Inamdar NM. Spontaneous deamination of cytosine and 5-methylcytosine residues in DNA and replacement of 5-methylcytosine residues with cytosine residues. *Mutat Res* (1990) 238(3):277–86. doi:10.1016/0165-1110(90)90019-8
50. Ginn SL, Smyth C, Wong M, Bennetts B, Rowe PB, Alexander IE. A novel splice-site mutation in the common gamma chain (gammac) gene IL2RG results in X-linked severe combined immunodeficiency with an atypical NK+ phenotype. *Hum Mutat* (2004) 23(5):522–3. doi:10.1002/humu.9235
51. Mou W, He J, Chen X, Zhang H, Ren X, Wu X, et al. A novel deletion mutation in IL2RG gene results in X-linked severe combined immunodeficiency with an atypical phenotype. *Immunogenetics* (2017) 69(1):29–38. doi:10.1007/s00251-016-0949-3
52. Estévez OA, Ortega C, Fernández S, Aguado R, Rumbao J, Perez-Navero J, et al. A novel IL2RG mutation presenting with atypical T(–)B(+)NK+ phenotype: rapid elucidation of NK cell origin. *Pediatr Blood Cancer* (2014) 61(1):178–9. doi:10.1002/pbc.24717
53. Lee YN, Frugoni F, Dobbs K, Walter JE, Giliani S, Gennery AR, et al. A systematic analysis of recombination activity and genotype-phenotype correlation in human recombination-activating gene 1 deficiency. *J Allergy Clin Immunol* (2014) 133(4):1099–108. doi:10.1016/j.jaci.2013.10.007
54. Wu WC, Wang W, Song HM, Ma MS, Tang XY, Jian S, et al. [A major histocompatibility complex class II deficiency case report and literature review]. *Zhonghua Er Ke Za Zhi* (2016) 54(8):614–8. doi:10.3760/cma.j.issn.0578-1310.2016.08.013
55. Shrestha D, Szöllosi J, Jenéi A. Bare lymphocyte syndrome: an opportunity to discover our immune system. *Immunol Lett* (2012) 141(2):147–57. doi:10.1016/j.imlet.2011.10.007
56. Ouederni M, Vincent QB, Frange P, Touzot F, Scerra S, Bejaoui M, et al. Major histocompatibility complex class II expression deficiency caused by a RFXANK founder mutation: a survey of 35 patients. *Blood* (2011) 118(19):5108–18. doi:10.1182/blood-2011-05-352716
57. Renella R, Picard C, Neven B, Ouachée-Chardin M, Casanova JL, Le Deist F, et al. Human leucocyte antigen-identical haematopoietic stem cell transplantation in major histocompatibility complex class II immunodeficiency: reduced survival correlates with an increased incidence of acute graft-versus-host disease and pre-existing viral infections. *Br J Haematol* (2006) 134(5):510–6. doi:10.1111/j.1365-2141.2006.06213.x
58. Mellouli F, Mustapha IB, Khaled MB, Besbes H, Ouederni M, Mekki N, et al. Report of the Tunisian registry of primary immunodeficiencies: 25-years of experience (1988–2012). *J Clin Immunol* (2015) 35(8):745–53. doi:10.1007/s10875-015-0206-9
59. Bousfiha AA, Jeddane L, El Hafidi N, Benajiba N, Rada N, El Bakkouri J, et al. First report on the Moroccan registry of primary immunodeficiencies: 15 years of experience (1998–2012). *J Clin Immunol* (2014) 34(4):459–68. doi:10.1007/s10875-014-0005-8
60. Reda SM, Afifi HM, Amine MM. Primary immunodeficiency diseases in Egyptian children: a single-center study. *J Clin Immunol* (2009) 29(3):343–51. doi:10.1007/s10875-008-9260-x
61. Kechout N, Attal N, Doudou F, Boukari R, Ardjoun M, Abbadi MC. Study of primary immunodeficiencies in highly consanguineous North African populations. *BMC Proc* (2011) 5(Suppl 1):35. doi:10.1186/1753-6561-5-s1-p35
62. Reda SM, El-Ghoneimy DH, Afifi HM. Clinical predictors of primary immunodeficiency diseases in children. *Allergy Asthma Immunol Res* (2013) 5(2):88–95. doi:10.4168/air.2013.5.2.88
63. Barbouche MR, Galal N, Ben-Mustapha I, Jeddane L, Mellouli F, Ailal F, et al. Primary immunodeficiencies in highly consanguineous North African populations. *Ann N Y Acad Sci* (2011) 1238:42–52. doi:10.1111/j.1749-6632.2011.06260.x
64. Naamane H, El Maataoui O, Ailal F, Barakat A, Bennani S, Najib J, et al. The 752delG26 mutation in the RFXANK gene associated with major histocompatibility complex class II deficiency: evidence for a founder effect in the Moroccan population. *Eur J Pediatr* (2010) 169(9):1069–74. doi:10.1007/s00431-010-1179-6
65. Elhasid R, Etzioni A. Major histocompatibility complex class II deficiency: a clinical review. *Blood Rev* (1996) 10(4):242–8. doi:10.1016/S0268-960X(96)90008-9
66. Kuo CY, Chase J, Garcia Lloret M, Stiehm ER, Moore T, Aguilera MJ, et al. Newborn screening for severe combined immunodeficiency does not identify bare lymphocyte syndrome. *J Allergy Clin Immunol* (2013) 131(6):1693–5. doi:10.1016/j.jaci.2013.01.019
67. Hague RA, Rassam S, Morgan G, Cant AJ. Early diagnosis of severe combined immunodeficiency syndrome. *Arch Dis Child* (1994) 70(4):260–3. doi:10.1136/adc.70.4.260
68. Gennery AR, Cant AJ. Diagnosis of severe combined immunodeficiency. *J Clin Pathol* (2001) 54(3):191–5. doi:10.1136/jcp.54.3.191
69. Buckley RH. Molecular defects in human severe combined immunodeficiency and approaches to immune reconstitution. *Annu Rev Immunol* (2004) 22:625–55. doi:10.1146/annurev.immunol.22.012703.104614
70. Rivers L, Gaspar HB. Severe combined immunodeficiency: recent developments and guidance on clinical management. *Arch Dis Child* (2015) 100(7):667–72. doi:10.1136/archdischild-2014-306425
71. Kubiak C, Jyonouchi S, Kuo C, Garcia-Lloret M, Dorsey MJ, Sleasman J, et al. Fiscal implications of newborn screening in the diagnosis of severe combined immunodeficiency. *J Allergy Clin Immunol Pract* (2014) 2(6):697–702. doi:10.1016/j.jaip.2014.05.013

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Luk, Lee, Mao, Chan, Chen, Chen, He, Kechout, Suri, Tao, Xu, Jiang, Liew, Jirapongsananuruk, Daengsuwan, Gupta, Singh, Rawat, Abdul Latiff, Lee, Shek, Nguyen, Chin, Chien, Latiff, Le, Le, Lee, Li, Raj, Barbouche, Thong, Ang, Wang, Xu, Yu, Yu, Lee, Yau, Wong, Tu, Yang, Chong, Ho and Lau. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.