



**HAL**  
open science

## Kearns-Sayre syndrome is genetically and phenotypically heterogeneous

Josef Finsterer, Sinda Zarrouk-Mahjoub

► **To cite this version:**

Josef Finsterer, Sinda Zarrouk-Mahjoub. Kearns-Sayre syndrome is genetically and phenotypically heterogeneous. *Pediatrics Medica e Chirurgica*, 2018, 40 (1), 10.4081/pmc.2018.193 . pasteur-02009260

**HAL Id: pasteur-02009260**

**<https://hal-riip.archives-ouvertes.fr/pasteur-02009260>**

Submitted on 6 Feb 2019

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial | 4.0 International License

# Kearns-Sayre syndrome is genetically and phenotypically heterogeneous

Josef Finsterer,<sup>1</sup> Sinda Zarrouk-Mahjoub<sup>2</sup>

<sup>1</sup>Krankenanstalt Rudolfstiftung, Messerli Institute, Veterinary University of Vienna, Vienna, Austria; <sup>2</sup>University of Tunis El Manar and Genomics Platform, Pasteur Institute of Tunis, Tunisia

Dear Editor,

With interest we read the article by Berio *et al.* about a 13-year-old male with Kearns-Sayre syndrome (KSS), manifesting with cardinal features (progressive external ophthalmoplegia, onset <20y, pigmentary retinopathy), and less infrequent, additional features (hypoacusis, ataxia, PNS involvement, short stature, growth hormone deficiency, lactic acidosis, facial dysmorphism, hypoparathyroidism, emesis, aortic insufficiency, subaortic septum hypertrophy, right bundle-branch block, double vision, and white matter lesions).<sup>1</sup> We have the following comments and concerns.

We do not agree with the statement that cerebral white matter lesions (WMLs) found on imaging are unique to KSS patients.<sup>1</sup> Patchy or diffuse subcortical WMLs are a frequent finding in mitochondrial disorders (MIDs) with cerebral involvement. Particularly, patients with non-specific mitochondrial syndromes show up with WMLs as described in Berio's case.<sup>2</sup>

The authors hypothesize that WMLs of the described patient were partially due to abnormal representation or migration of neural crest cells. If this is the case, WMLs presented should be present

already at birth in KSS patients. However, we are not aware of KSS patients with WMLs already at birth.

Since the mother presented with hypothyroidism, a frequent feature of MIDs, we should know if the mother was tested positive for the mtDNA deletion. Though KSS is usually sporadic, in about 4% of the cases there is transmission via the maternal line.<sup>3</sup>

If cerebral lesions in KSS also occur bilaterally in the *substantia nigra*, as mentioned in the discussion, we can expect that at least some KSS patients present with Parkinsonism. However, no reports about KSS and Parkinsonism have been published.

With regards to growth retardation, we would be interested to know if there was empty sella, pituitary adenoma, or brainstem lesions.

Overall, this interesting case report requires clarification of some inconsistencies and would profit from providing genetic data of first-degree relatives.

## References

1. Berio A, Piazzini A, Traverso CE. Kearns-Sayre syndrome with facial and white matter extensive involvement: a (mitochondrial and nuclear gene related?) neurocristopathy? *Pediatr Med Chir* 2017;39:107-11.
2. Bindu PS, Sonam K, Chiplunkar S, et al. Mitochondrial leukoencephalopathies: A border zone between acquired and inherited white matter disorders in children? *Mult Scler Relat Disord* 2018;20:84-92.
3. Poulton J, Finsterer J, Yu-Wai-Man P. Genetic counselling for maternally inherited mitochondrial disorders. *Mol Diagn Ther* 2017;21:419-29.

Correspondence: Josef Finsterer, Krankenanstalt Rudolfstiftung, Messerli Institute, Veterinary University of Vienna, Postfach 20, 1180 Vienna, Austria.

Tel.: +43.1.71165 – Fax: +43.1.4781711.

E-mail: ffigs1@yahoo.de

Key words: Kearns-Sayre syndrome; mtDNA; Leukoencephalopathy; Phenotype; Genotype.

Contributions: JF: design, literature search, discussion, first draft; SZM: literature search, discussion, critical comments.

Conflict of interest: the authors declare no potential conflict of interest.

Funding: none.

Note: this Letter to the Editor relates to Berio A, Piazzini A, Traverso CE. Kearns-Sayre syndrome with facial and white matter extensive involvement: a (mitochondrial and nuclear gene related?) neurocristopathy? *Pediatr Med Chir* 2017;39:107-11.

Received for publication: 19 March 2018.

Accepted for publication: 22 May 2018.

This work is licensed under a Creative Commons Attribution NonCommercial 4.0 License (CC BY-NC 4.0).

©Copyright J. Finsterer and S. Zarrouk-Mahjoub, 2018

Licensee PAGEPress, Italy

*La Pediatria Medica e Chirurgica* 2018; 40:193

doi:10.4081/pmc.2018.193