



HAL
open science

Mitochondrial multiorgan disorder syndrome (MIMODS) due to a compound heterozygous mutation in the ACAD9 gene

Josef Finsterer, Sinda Zarrouk-Mahjoub

► **To cite this version:**

Josef Finsterer, Sinda Zarrouk-Mahjoub. Mitochondrial multiorgan disorder syndrome (MIMODS) due to a compound heterozygous mutation in the ACAD9 gene. *Molecular Genetics and Metabolism Reports*, 2017, 13, pp.31-32. 10.1016/j.ymgmr.2017.07.009 . pasteur-02010473

HAL Id: pasteur-02010473

<https://riip.hal.science/pasteur-02010473>

Submitted on 7 Feb 2019

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License



Correspondence

Mitochondrial multiorgan disorder syndrome (MIMODS) due to a compound heterozygous mutation in the *ACAD9* gene

ARTICLE INFO

Keywords:

Mitochondrial
mtDNA
Phenotype
Genotype
Lymphocytes
Abnormal immune response

Letter to the Editor

We read with interest the article by Fragaki et al. about a 1yo Algerian female with hypertrophic cardiomyopathy, developmental delay, growth retardation, hepatomegaly, failure-to-thrive, general hypotonia, and lactic acidosis, being attributed to the biallelic variants c.1204G > T and c.358delT in the *ACAD9*-gene encoding for acyl-CoA-dehydrogenase-9, an essential assembly-factor for respiratory-chain complex-I and component of fatty-acid oxidation [1,2]. We have the following comments and concerns.

Was hypotonia attributed to involvement of the skeletal muscles, the peripheral nerves, or the cerebrum? Did cerebral MRI show any abnormalities that could explain the CNS-abnormalities? Leigh-like features are well appreciated as cerebral manifestation of *ACAD9*-mutations (Table 1) [3].

Patients carrying *ACAD9*-mutations may develop myopathy (Table 1) [4]. Was there clinical, blood-chemical, electromyographic, or bioptic evidence for myopathy in the index case? Were nerve-conduction studies normal?

The parents carried one of the mutations each. Was either of them clinically affected? Did any of the other first degree relatives exhibit phenotypic features of *ACAD9*-mutations (Table 1)? Were other first degree relatives neurologically or genetically investigated?

Though liver failure and hepatopathy have been previously described in association with *ACAD9*-mutations (Table 1), hepatomegaly seems to be a unique feature. Since *ADAC9*-mutations may cause cardiac compromise [5], it is conceivable that hepatomegaly was due to systolic dysfunction. Did the patient ever develop clinical or instrumental evidence of heart failure? Were proBNP-values elevated? Was heart transplantation ever considered? Did the patient suffer from an infectious disease or did she carry a second mutation? Was a liver-biopsy taken? Was there deposition of lipids?

Complex-I deficiency is frequently associated with *ACAD9*-mutations [6] but is complex-II and complex-IV deficiency, not being previously described, truly attributable to a secondary effect? Did the patient carry a second mutation?

Overall, this interesting case could be more meaningful if more clinical and instrumental investigations would have been carried out in the index case and in her first degree family members.

Author contribution

JF: design, literature search, discussion, first draft, SZ-M: literature search, discussion, critical comments.

* There are no conflicts of interest.
No funding was received.

<http://dx.doi.org/10.1016/j.ymgmr.2017.07.009>

Received 20 July 2017; Accepted 21 July 2017

Available online 28 July 2017

2214-4269/ © 2017 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Table 1
Phenotypic manifestations of ACAD9 mutations.

Phenotypic manifestation	Reference
Central nervous system	
Developmental delay	[Fragaki, 2017, Garone, 2013]
Failure to thrive	[Fragaki, 2017]
Microcephaly	[Aintablian, 2016]
Dystonia	[Aintablian, 2016]
Seizures	[Collet, 2016]
Cognitive impairment	[Collet, 2016]
Optic atrophy	[Collet, 2016]
Leigh-like features on MRI	[Aintablian, 2016]
Heart	
Hypertrophic cardiomyopathy	[Aintablian, 2016]
Mild ventricular hypertrophy	[Dewulf, 2016]
Dilated cardiomyopathy	[Dewulf, 2016]
Patent ductus arteriosus	[Dewulf, 2016]
Intestines	
Acute liver failure	[Valencia, 2016]
Vomiting	[Schrank, 2017]
Nausea	[Schrank, 2017]
Hepatomegaly	[Fragaki, 2017]
Hepatopathy	[Robinson, 1998]
Episodes of acute liver dysfunction	[He, 2007]
Kidneys	
Proximal tubulopathy	[Collet, 2016]
Renal failure	[Collet, 2016]
Endocrine organs	
Secondary ovarian failure	[Collet, 2016]
Muscles	
Myopathy	[Collet, 2016]
Muscle weakness	[Nouws, 2014]
Fatigue	[Schrank, 2017]
Exercise intolerance	[Schrank, 2017, Garone, 2013]
Others	
Fatal neonatal multiorgan failure	[Leslie, 2016]
Postnatal growth retardation	[Fragaki, 2017]
Lactic acidosis	[Aintablian, 2016, Fragaki, 2017]
Intrauterine growth retardation	[Lagoutte-Renosi, 2015]
Reye-like episode	[He, 2007]
Elevated serum alanine	[Fragaki, 2017]
Reduced serum ornithine and citrulline	[Fragaki, 2017]

References

- [1] K. Fragaki, A. Chaussonot, A. Boutron, S. Bannwarth, C. Rouzier, B. Chabrol, V. Paquis-Flucklinger, Assembly defects of multiple respiratory chain complexes in a child with cardiac hypertrophy associated with a novel ACAD9 mutation, *Mol. Genet. Metab.* 121 (2017) 224–226.
- [2] J. Lagoutte-Renosi, I. Ségalas-Milazzo, M. Crahes, F. Renosi, L. Menu-Bouaouiche, S. Torre, C. Lardennois, M. Rio, S. Marret, C. Brasse-Lagnel, A. Laquerrière, S. Bekri, Lethal neonatal progression of fetal cardiomegaly associated to ACAD9 deficiency, *JIMD Rep.* 17 (2015 Oct).
- [3] H.K. Aintablian, V. Narayanan, N. Belnap, K. Ramsey, T.A. Grebe, An atypical presentation of ACAD9 deficiency: diagnosis by whole exome sequencing broadens the phenotypic spectrum and alters treatment approach, *Mol. Genet. Metab.* 10 (2016) 38–44.
- [4] J.P. Dewulf, C. Barrea, M.F. Vincent, C. De Laet, R. Van Coster, S. Seneca, S. Marie, M.C. Nassogne, Evidence of a wide spectrum of cardiac involvement due to ACAD9 mutations: report on nine patients, *Mol. Genet. Metab.* 118 (2016) 185–189.
- [5] M. Collet, Z. Assouline, D. Bonnet, M. Rio, F. Iserin, D. Sidi, A. Goldenberg, C. Lardennois, M.D. Metodiev, B. Haberberger, T. Haack, A. Munnich, H. Prokisch, A. Rötig, High incidence and variable clinical outcome of cardiac hypertrophy due to ACAD9 mutations in childhood, *Eur. J. Hum. Genet.* 24 (2016) 1112–1116.
- [6] B. Schrank, B. Schoser, T. Klopstock, P. Schneiderat, R. Horvath, A. Abicht, E. Holinski-Feder, S. Augustis, Lifetime exercise intolerance with lactic acidosis as key manifestation of novel compound heterozygous ACAD9 mutations causing complex I deficiency, *Neuromuscul. Disord.* 27 (2017) 473–476.

Josef Finsterer¹, Sinda Zarrouk-Mahjoub¹
 Krankenanstalt Rudolfstiftung, Vienna, Austria
 University of Tunis El Manar and Genomics Platform, Pasteur Institute of Tunis, Tunisia
 E-mail address: ffigs1@yahoo.de

* Corresponding author at: Postfach 20, 1180 Vienna, Austria.

¹ Both authors contributed equally.