A potential robust antiviral defense state in the common vampire bat: Expression, induction and molecular characterization of the three interferon-stimulated genes -OAS1, ADAR1 and PKR

Sarkis Sarkis, Stéphanie Dabo, Marie-Claude Lise, Christine Neuveut, Eliane Meurs, Vincent Lacoste, Anne Lavergne

► To cite this version:

Sarkis Sarkis, Stéphanie Dabo, Marie-Claude Lise, Christine Neuveut, Eliane Meurs, et al.. A potential robust antiviral defense state in the common vampire bat: Expression, induction and molecular characterization of the three interferon-stimulated genes -OAS1, ADAR1 and PKR. *Developmental and Comparative Immunology*, 2018, 85, pp.95-107. 10.1016/J.DCI.2018.04.006 . pasteur-02197397

HAL Id: pasteur-02197397

<https://hal-riip.archives-ouvertes.fr/pasteur-02197397>

Submitted on 6 Apr 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A potential robust antiviral defense state in the common vampire bat: expression, induction and molecular characterization of the three interferon-stimulated genes - OAS1, ADAR1 and PKR

Sarkis Sarkis^{a,*}, Stéphanie Dabo^b, Marie-Claude Lise^a, Christine Neuveut^b, Eliane F. Meurs^b, Vincent Lacoste^a and Anne Lavergne^{a,*}

^a Laboratoire des Interactions Virus-Hôtes, Institut Pasteur de la Guyane, Cayenne, French Guiana

^b Hepacivirus and Innate Immunity, Institut Pasteur, 75015 Paris, France

* Corresponding authors.

Laboratoire des Interactions Virus-Hôtes, Institut Pasteur de la Guyane, 23 avenue Pasteur, 97300 Cayenne, French Guiana.

E-mail addresses : sarkis.sarkis@hotmail.com (S. Sarkis); alavergne@pasteur-cayenne.fr (A. Lavergne)

Keywords: *Desmodus rotundus*, Innate immunity, ISGs, OAS1a, OAS1b, ADAR1, PKR

Abstract

Bats are known to harbor many zoonotic viruses, some of which are pathogenic to other mammals while they seem to be harmless in bats. As the interferon (IFN) response represents the first line of defense against viral infections in mammals, it is hypothesized that activation of the IFN system is one of the mechanisms enabling bats to co-exist with viruses. We have previously reported induction of type I IFN in a cell line from the common vampire bat, *Desmodus rotundus*, upon polyinosinic:polycytidylic acid (poly(I:C)) stimulation. To deepen our knowledge on *D. rotundus*' IFN-I antiviral response, we molecularly characterized three interferon-stimulated genes (ISGs), *OAS1*, *PKR* and *ADAR1*, closely implicated in the IFN-I antiviral response, and tested their functionality in our cellular model. We first found that *D. rotundus* encoded two *OAS1* paralogs, *OAS1a* and *OAS1b*, and that the functional domains of the four ISGs characterized were highly conserved with those of other mammals. Despite their significant transcription level in the absence of stimulation, the transcription of the four ISGs characterized was enhanced by poly(I:C). In addition, the transcription of *OAS1a* and *OAS1b* appears to be differentially regulated. These findings demonstrate an active ISG antiviral response in *D. rotundus* in which *OAS1b* may play an important role.

Highlights

- Four ISGs implicated in the antiviral response were molecularly characterized.
- Two *OAS1* paralogs were characterized in *D. rotundus*.
- The functional domains of the characterized ISGs are highly conserved relative to their mammalian homologs.
- Significant induction of *Oas1b* was observed after stimulation with poly(I:C).

1. Introduction

The interferon (IFN)-mediated innate immune response provides a robust first line of defense against invading pathogens. Pathogen-associated molecular patterns' (PAMP) recognition by pattern recognition receptors (PRRs) leads to a subsequent production of IFN, especially of types I (IFN α/β) and III (IL28A, IL28B, IL29), considered as the main antiviral IFNs (Borden et al., 2007). Newly synthesized IFN molecules up-regulate the transcription of hundreds of interferon-stimulated genes (ISGs) resulting in the production of various antiviral effectors (Der et al., 1998). This cascade leads to a remarkable antiviral state of infected and neighboring cells and limits the spread of infectious agents. Considering the large number of ISGs activated upon IFN signaling, considerable redundancy in the system is likely. Among the molecules with important biological functions induced by IFN are the 2'5' oligoadenylate synthetase/ribonuclease L (OAS/RNase L) system, the adenosine deaminase RNA-specific (ADAR) as well as the dsRNA-activated protein kinase (PKR), which play a critical role in the host's antiviral defense mechanism and are among the most extensively studied ISGs (Der et al., 1998; George and Samuel, 1999; Levy et al., 2002; Sen, 2001). Even though the specific functions of mammalian OAS1, ADAR and PKR have yet to be clarified, their structural (architectural) domain organizations are well characterized and show a high degree of similarity among species.

The *OAS* gene family comprises three classes of genes in human, *OAS1*, *OAS2* and *OAS3*, in addition to an "*OAS-like*" (*OASL*), which is catalytically inactive. The mouse *OAS* cluster, the second most well known after the human *OAS* cluster, contains one *OAS2* and one *OAS3* gene; two *OASLs*, of which one is a pseudogene; and eight paralogs of *OAS1*, *Oas1a*–*Oas1h* (Kristiansen et al., 2011). Activation of the OAS proteins in presence of single-stranded (ss) or double-stranded (ds) RNA triggers the synthesis of 2'5' oligoadenylates. These in turn bind to and activate the latent RNaseL, which restricts viral propagation through direct and indirect

70 mechanisms, including i) viral genome degradation (RNA and DNA), ii) cellular mRNA
71 degradation, including mitochondrial RNA followed by apoptosis, as well as iii) amplification
72 of IFN signaling through the release of additional PAMPs activating cytoplasmic helicases,
73 which in turn activate type I IFN synthesis and create a positive feedback loop (Justesen et al.,
74 2000; Malathi et al., 2007, 2005; Silverman, 2003; Xiang et al., 2003). Regarding its domain
75 organization, on its N-terminus OAS1 comprises a nucleotidyl transferase domain (NTP-
76 transf_2) constituted of a dsRNA-binding domain flanked by two ATP-binding domains,
77 which plays a critical role in OAS1 activation. On its C-terminus, OAS1 presents the
78 OAS1_C domain including the region of enzymatic activity thought to activate latent
79 RNaseL (Sarkar et al., 2002).

80 ADAR belongs to a family of proteins that modulate nucleic acid integrity and play an
81 important role in the defense against viral RNAs. All ADAR family members present double-
82 stranded RNA-binding domains (dsRBDs) at their N-termini and a catalytic deaminase
83 domain located at their C-termini (Mannion et al., 2015). A unique feature of ADAR1, when
84 compared to other members of the ADAR family, is the presence of Z-DNA-binding domains
85 (ZBDs) at its N-terminus with a Z α and a Z β domain (Kim et al., 1999). In addition to the
86 capacity of the Z α domain to bind to Z-DNA/RNA (Herbert et al., 1997), it has been reported
87 that all proteins containing this domain are implicated in the type I IFN response pathway
88 (Athanasiadis et al., 2005). Furthermore, the Z α domain is essential for the location of
89 ADAR1 to cytoplasmic stress granules following activation. Upon binding to dsRNA,
90 ADAR1 induces editing of adenosine to inosine, one of the most abundant modifications of
91 mammalian RNA (Bazak et al., 2014; Goodman et al., 2012). However, depending on the
92 viral species, the editing event can make ADAR1 act as either a proviral or an antiviral factor.
93 Indeed, whereas editing by ADAR1 of lymphocytic choriomeningitis virus introduces
94 mutations in viral proteins and reduces infectivity, editing of HIV-1 rev and tat proteins leads

to increased infectivity (Doria et al., 2009). In addition, ADAR1 can also exert a proviral function through its Z-DNA and dsRBD domains by interacting with PKR, another dsRBD-containing protein. Such interaction negatively regulates the antiviral function of PKR (Clerzius et al., 2009).

PKR (protein kinase dsRNA-dependent) is one of the four eIF-2 α serine-threonine kinases that controls protein synthesis through the phosphorylation of the alpha subunit of eukaryotic initiation factor 2 (eIF-2 α). PKR is ubiquitously expressed and can be induced upon IFN treatment (Meurs et al., 1990, 1992; Williams, 2001). PKR becomes activated upon binding to dsRNA, as during viral infection, or to cellular activators in response to stress (Dabo and Meurs, 2012). Because of PKR's role in controlling protein synthesis, it can inhibit infection by a number of viruses. Nevertheless, other viruses, such as the hepatitis C virus, benefit from the general shut-down of protein synthesis, which promotes their own eIF-2 α -independent translation (Arnaud et al., 2010; Garaigorta and Chisari, 2009). In addition, PKR modulates signaling pathways involved in inflammation independently of its enzymatic activity (Bonnet et al., 2006; García et al., 2007). PKR consists of two functionally distinct domains: a N-terminal dsRNA binding regulatory domain (dsRBD) and a C-terminal kinase catalytic domain. The dsRNA-binding domain contains two dsRNA-binding motifs (dsRBMs). These motifs are rich in basic residues and confer to PKR the capability of binding to dsRNA independently of the sequence. As a member of the kinase proteins, PKR presents all the conserved catalytic kinase subdomains. In addition, the homology within subdomain VI identifies PKR as a serine/threonine kinase (Meurs et al., 1990). PKR also possesses an insert region C-terminal of Domain IV. This insert region and the dsRBD are key determinants of the PKR function and regulation.

Bats are known to harbor many zoonotic viruses, some of which are pathogenic to other mammals while they seem to be harmless or only rarely associated with clinical signs in bats

(Moratelli and Calisher, 2015). Strikingly, mechanisms that allow bats to coexist with these viruses have not yet been fully researched. It is currently acknowledged that early control of viral replication by their innate immune system might be one of the mechanisms implicated, enabling bats to coexist with viruses (Baker et al., 2013). Following the establishment of a *Desmodus rotundus* cell line, FluDero, and the molecular characterization of the RIG-I-like receptor members and of type I interferons (Sarkis et al., 2018), we studied several ISGs from this bat species given that they are responsible for the antiviral activity of IFNs in mammals (Der et al., 1998; He et al., 2014; Levy et al., 2002; Sen, 2001; Zhou et al., 2013). Here, we describe the molecular characterization of three ISGs, *OAS1*, *ADAR1* and *PKR*, from *Desmodus rotundus* and report differential expression profiles for these genes upon stimulation of the FLuDero fetal lung cell line with the synthetic analog of dsRNA, the polyinosinic:polycytidylic acid, poly(I:C).

2. Materials and Methods

2.1. Cell line, cell culture and poly(I:C) stimulation

Establishment of the FluDero cell line has been described previously (Sarkis et al., 2018). The cell line was cultured in prewarmed DMEM/Ham's Nutrient Mixture F12 (Sigma) supplemented with 10% fetal bovine serum (FBS) (Sigma), 100 units/mL penicillin and 100 µg/mL streptomycin and was maintained in a humidified incubator with 5% CO₂ at 37°C. For poly(I:C) stimulation, FLuDero cells were seeded at a density of 2.5×10^5 cells/well in 12-well plates. Cells were stimulated by transfection with 20 µg/mL of poly(I:C) (Sigma, cat #P9582) by calcium phosphate transfection (Invitrogen), 20 h after seeding. Cells were harvested in TRIzol Reagent at 0, 3, 6, 9, 12, 24, 48 and 72 h post-transfection and stored at -80°C until RNA extraction. The experiment was performed three times.

2.2. RNA extraction and cDNA synthesis

Total RNA was extracted from FLuDero cells transfected with poly(I:C) using TRIzol Reagent (Invitrogen) as recommended by the manufacturer. cDNAs were synthesized using Superscript[®] III reverse transcriptase (Invitrogen) and random hexamers following the manufacturer's instructions.

2.3. Sequence identification and domain characterization of *OAS1*, *ADAR1* and *PKR*

To generate the full-length coding sequences of *OAS1*, *ADAR1* and *PKR*, all amplifications were performed on cDNA obtained from poly(I:C)-transfected FLuDero cells. Internal coding regions were amplified by PCR using the AmpliTaq Gold DNA polymerase kit (Thermo Fischer Scientific). Different combinations of consensus and specific degenerate primers were designed using Primer3 (v.0.4.0) based on alignments of orthologous mammal sequences, which were available in the GenBank database (**Supplementary Table 1**). The 5'- and 3'-terminal sequences were obtained by rapid amplification of cDNA ends (RACE) using the FirstChoice[®] RLM-RACE Kit (Ambion, Inc.). PCR and RACE products were cloned using the pCR[™]4-TOPO[®] TA CLONING[®] KIT (Invitrogen). The sequences obtained were confirmed by similarity analysis using the NCBI BLAST search tool (<http://www.ncbi.nlm.nih.gov/BLAST>). The structural domains of each protein were identified using SMART software (<http://smart.embl-heidelberg.de/>) (Letunic et al., 2015). Multi-alignment was performed with the ClustalW program (https://npsa-prabi.ibcp.fr/cgi-bin/npsa_automat.pl?page=/NPSA/npsa_clustalw.html) using default parameters.

2.4. Phylogenetic analysis

The complete coding sequences for *ADAR1* and *PKR* of 18 and 17 representatives Laurasiatherian and Euarchontoglires, respectively, were downloaded from the NCBI database. Available sequences included those from the bat species *Eptesicus fuscus*, *Miniopterus natalensis*, *Myotis brandtii*, *Myotis davidii*, *Myotis lucifugus*, *Pteropus alecto*,

Pteropus vampyrus, *Rhinolophus sinicus* and *Rousettus aegyptiacus* (**Supplementary Table 2**). The ISG sequences of *D. rotundus* obtained in this study were further added to each data set. After alignment and editing, the MEGA version 6.0 software (Tamura et al., 2013) was used to compute pairwise comparisons in the nucleotide and amino acid using uncorrected p-distances. The best-fit amino acid evolutionary substitution model (Jones-Taylor-Thornton + G + F) was determined using MEGA version 6.0. Bayesian phylogenetic analyses were performed using MrBAYES 3. Markov Chain Monte Carlo (MCMC) simulations were run for 10,000,000 generations, with four simultaneous chains, using a sample frequency of 500 and a burn-in of 25,000. Majority rule consensus trees were obtained from the output. Validation of the inference was assessed based on the standard deviation of split frequencies, which was less than the expected threshold value of 0.01 (calculated value of 0.002). ADAR1 and PKR trees were rooted with the rodent *Mus musculus* sequences. Coding sequences from 33 representative mammalian species retrieved in NCBI were used for OAS1 pairwise comparisons and phylogenetic analysis (**Supplementary Table 2**). Eight corresponded to the bat species *E. fuscus* (two paralogs), *M. natalensis* (two paralogs), *M. brandtii* (two paralogs), *M. davidii*, *M. lucifugus* (two paralogs), *P. alecto*, *P. vampyrus* and *R. aegyptiacus*. An OAS1 tree was constructed based on the alignment of amino acid residues and was rooted with OASL sequences from *M. natalensis*, *P. alecto*, *P. vampyrus* and *R. aegyptiacus*.

2.5. Quantitative PCR

Quantitative PCR (qPCR) was performed on a StepOne™ Real-Time PCR System using the TaqMan® Universal PCR Master Mix (Applied Biosystems). Primers and TaqMan® probes were designed using Primer Express 3.0 (Applied Biosystems) (**Supplementary Table 1**). Each reaction was performed in duplicate. The cycling conditions were those recommended by the manufacturer: 55°C for 2 min, 95°C for 2 min followed by 40 cycles at 95°C for 15 s, 60°C for 1 min. For absolute quantification, the exact number of copies of the

gene of interest was calculated using a plasmid DNA standard curve. Individual expression values were normalized and compared to mRNA encoding β -actin. Values are shown graphically as fold induction compared to the mock sample (0 h transfection) for each experiment.

3. Results

3.1. Sequence characterization of *D. rotundus* *OAS1a*, *OAS1b*, *ADAR1* and *PKR* genes

Primers derived from alignments of the orthologous mammalian *OAS1*, *ADAR1* and *PKR* sequences were used to characterize *OAS1*, *ADAR1* and *PKR* genes from mRNA extracted from the FluDero cell line. The full-length coding sequences for *ADAR1* and *PKR* were identified. Sequence analysis revealed that their coding regions were 3444 and 1611 base pairs (bp) long, respectively, encoding proteins of 1149 and 537 amino acids (aa), respectively. In addition, we characterized two *OAS1* paralog genes in *D. rotundus*, which we named *OAS1a* and *OAS1b*. The partial coding sequence of *OAS1a* was 1056 bp in length (352 aa). On the other hand, the complete coding region of *OAS1b* was 1101 bp long encoding a 367-aa protein. When comparing the *OAS1a* and *OAS1b* with their mammalian counterparts, *OAS1a* seems to lack 17aa at its C-terminal.

Pairwise comparison done on 897 nucleotides corresponding to 299 amino acid residues showed that the two *OAS1* paralogs shared 81.6 and 77.3% of sequence identity at the nucleotide and amino acid levels, respectively. However, when compared to their mammalian homologs, *OAS1a* showed a higher level of sequence identity at the nucleotide and amino acid levels than *OAS1b*. For instance, *OAS1a* exhibited 84% and 80% identity at the nucleotide and amino acid levels, respectively, and *OAS1b* shared 78 and 73% identity with the *OAS1* sequence of the *Pteropus* species. In the same way, *OAS1a* shared 73–80% amino acid identity with its equine, canine, bovine and human homologs, while *OAS1b* presented 69–

75% sequence identity (**Supplementary Table 3**). *D. rotundus*' *ADAR1* and *PKR* were also subjected to pairwise comparison with their mammalian homologs. Among the four ISGs characterized, the complete amino acid sequence of *ADAR1* displayed the highest level of sequence identity with its mammalian counterparts, while *PKR* shared the lowest. For instance, *ADAR1* and *PKR* displayed 85.4–89.6% and 64.7–71% sequence identity at the protein level with their bat homologs, respectively (**Supplementary Table 3**). In addition, when comparing *D. rotundus* *PKR* with other mammalian homologs the identity at the protein level was slightly lower.

3.2. Conserved domain organization in *D. rotundus* ISGs

One of the hallmarks of the ISGs is their domain architecture responsible for their function. Sequence analysis of the two OAS1 paralogs identified in *D. rotundus* using the SMART software showed that these two proteins presented a similar structural domain organization to that described in their mammalian homologs. They possess the two characteristic functional domains of OAS1, *i.e.*, the NTP-transf_2 domain on their N-termini and C-terminal OAS1_C domain. These domains are located between aa 36–121 and 162–345, respectively (all values are given relative to *H. sapiens* sequences) (**Figure 1**). The *D. rotundus* OAS1a and OAS1b putative proteins showed conservation of the hallmark motifs crucial for OAS1 activity, *i.e.*, the LxxxP sequence required for enzymatic activity, a P-Loop motif with a (G[G/S]xx) sequence, which precedes the first triad residue corresponding to the second signature of the nucleotidyl transferase superfamily formed by Asparagine residues (represented with an inverted triangle), a K-R rich region and a CFK motif located in the C-terminal domain (**Figure 1**). Similarly, the *D. rotundus* ADAR1 protein contained all domains necessary for its function, including at its N-terminus the two Z α domains, a central dsRBD with three dsRBM, and a catalytic deaminase domain at its C-terminus. These domains and motifs are located between aa 132–202, 291–360, 504–569, 615–680, 727–793 and 840–

1222, respectively (**Figure 2a**). Finally, the deduced protein sequence of *D. rotundus* PKR showed the conservation of all domains relative to its function including the N-terminus dsRBD with the two dsRBM (dsRBM1 and dsRBM2) between aa 9–81 and 102–169 (yellow and light gray boxes, respectively) and all subdomains (open boxes) in the catalytic region. A variable region was also found in *D. rotundus* PKR between the dsRBM2 and the basic region (in dark gray) between aa 229 and 251. In addition, *D. rotundus* PKR presented a conserved insert region at the C-terminal of subdomain IV (blue) between aa 362 and 370. However, *D. rotundus* PKR did not possess the repeat sequence (green) found in the catalytic region of the human PKR, a common feature observed in other bat species and mice (**Figure 2b**).

3.3. The *D. rotundus*-characterized ISGs are closely related to their mammalian counterparts

To elucidate the evolutionary relationships of OAS1a and OAS1b, ADAR1 and PKR, phylogenetic trees were constructed based on multiple amino acid alignments constructed with other published mammal sequences (**Figure 3**). The OAS1 phylogenetic topology obtained was in agreement with the different mammalian orders, showing that OAS1a and OAS1b from *D. rotundus* belonged to a single clade of chiropters. Within this clade, OAS1a and OAS1b clustered with OAS1-like and OAS1 of microbats, respectively, and were supported with high posterior probability values (**Figure 3a**). On the other hand, the phylogenetic analyses of ADAR1 (**Figure 3b**) and PKR (**Figure 3c**) showed that all bat sequences belonged to a clade distinct from other mammal species, with the microbats (*D. rotundus*, *M. natalensis*, *E. fuscus* and *Myotis* species) diverging from the megabats (*Pteropus* species, *R. aegyptiacus*, and *R. sinicus*).

3.4. *D. rotundus* OAS1, ADAR1 and PKR can be induced in response to poly(I:C) stimulation

To investigate the functionality of these ISGs in FLDero, we studied their expression profile after poly(I:C) transfection (**Figure 4**). The kinetics of the expression of OAS1a,

OAS1b, ADAR1 and PKR mRNA revealed a similar induction profile. The induction was observed 6 h post-transfection, reaching a peak at 24 h followed by a decline at 48 h. Among the ISGs studied, *OAS1b* was the most inducible gene (400-fold up-regulation at 24 h) at all time points of the kinetics. In contrast, *ADAR1* and *PKR* were the least induced genes, showing a peak of induction at 24 h with a 15- and nine-fold induction, respectively. Strikingly, in the absence of stimulation, transcription of the four ISGs characterized was already significant, with a strong expression of PKR mRNA (**Figure 5**, 0 hrs).

4. Discussion

Viral infection in mammals often triggers the IFN-I-mediated frontline host defense mechanism, leading to the transcription of a wide range of ISGs (Randall and Goodbourn 2008). These ISGs confer an antiviral state to the host and neighboring cells by several mechanisms, including inhibition of viral transcription, translation and replication, as well as degradation of viral nucleic acids. We previously demonstrated that FLDero cells were competent to mount high levels of IFN-I after transfection with poly(I:C) (Sarkis et al., 2018). To deepen our understanding of the *D. rotundus*' IFN-I antiviral response, we molecularly characterized three ISGs, *OAS1*, *PKR* and *ADAR1*, closely implicated in the IFN-I antiviral response and looked at their expression upon poly(I:C) stimulation of FLDero cells.

Contrary to other mammals such as in human, which only possesses one *OAS1* gene, we characterized two *OAS1* paralogs (*OAS1a* and *OAS1b*) in *D. rotundus*, a feature shared with some bat species such as *Myotis lucifugus*, *M. natalensis* and *E. fuscus*, and other mammals such as the mouse, which has several *OAS1* paralogs. Nevertheless, even if the mouse expresses eight *OAS1* genes, only two (*OAS1a* and *OAS1g*) conserve their activities (Elkhateeb et al., 2016). The other paralogs are non-functional due to mutations or deletions of at least one of their functional domains. Multiple sequence alignment of *D. rotundus* *OAS1a* and *OAS1b* with their mammalian ortholog sequences demonstrated that they have the same structural organization and are thus likely to possess similar enzymatic and antiviral activities. For instance, the P-Loop motif, which is an ATP-binding site (Saraste et al., 1990; Tag-El-Din-Hassan et al., 2012), the DAD motif in D box, which is a Mg^{2+} -binding site (Sarkar et al., 2002; Yamamoto et al., 2000), the KR-rich region, which is an oligoadenylate and ATP-binding site (Marié et al., 1997), and the CFK motif located in the C-terminal domain, required for the tetramerization of the proteins, are all conserved (Ghosh et al., 1997). In a previous study on *P. alecto*, Zhou and colleagues characterized a single *OAS1* gene.

However, they did not exclude the possible existence of additional *OAS1* paralogs given that their analysis was based on a *P. alecto* genome assembly not fully achieved at the time of the study (Zhou et al., 2013). In the database, many bat *OAS1* genes have been annotated as “OAS1-like” such as in *M. lucifigus*, *P. vampyrus*, *M. natalensis* and *E. fuscus*. However, these genes do not show any mutation on their functional sites, which could result in a loss of function (Guénet et al. 2015). Moreover, the *OAS1* phylogenetic tree reveals that some bat *OAS1* and *OAS1-like* sequences are intermixed within different clades (Fig. 3a). A consensus on “OAS1” naming is therefore needed. In this regard, the generation of more *OAS1* sequences from other bat species should extend our understanding of their evolutionary history.

The molecular characterization and domain prediction of PKR show that the regulatory and kinase subdomains of the *D. rotundus*’ PKR protein are remarkably conserved with respect to those of other mammals. For instance, the N-terminal domain of *D. rotundus*’ PKR includes a repeated motif (dsRBD) that possesses dsRNA-binding activity and the C-terminal half of PKR includes the 11 conserved catalytic subdomains characteristic of kinases. Our multiple sequence alignment of mammals’ PKR shows the conservation of the major residues crucial to the function of the protein such as the highly conserved lysine K⁶⁴ of the RNA-binding domain required for its RNA-binding activity, the K²⁹⁶ in the catalytic subdomain II required for its kinase activity and the T⁴⁴⁶ within the activation loop that confers to PKR its specificity as a serine/threonine kinase (Dabo and Meurs, 2012). In addition, the salt bridge interaction of R²⁶² and D²⁶⁶ at the dimer interface of PKR, which is critical to eIF2 α phosphorylation and PKR activation, is also conserved (Dey et al., 2007).

Similarly, depending on the sequence alignment, ADAR1 is also highly conserved with those of other mammals. For instance, we found that residues H⁹¹⁰ and E⁹¹² of the CHAE motif, considered as a key element of the catalytic core characteristic of deaminases, are

conserved. It has been shown that their mutations completely abolish A-to-I editing activity (Lai et al., 1995; Liu and Samuel, 1996). Further, C⁹⁶⁶ and C¹⁰³⁶, which are believed to play a role in zinc coordination along with H⁹¹⁰, are necessary for ADAR1 deaminase activity and are also conserved in *D. rotundus* (Lai et al., 1995). Moreover, as in other mammals, the lysine residues crucial for both RNA binding and enzymatic activity, present in the core sequence of each of the dsRBD domains (K⁵⁵⁴, K⁶⁶⁵, and K⁷⁷⁷) are conserved (Fierro-Monti and Mathews, 2000; Liu and Samuel, 1996; McCormack et al., 1994).

To test the functionality of the ISG system upon viral infection in our bat model, expression of the characterized ISGs was examined following transfection of FLuDero cells with poly(I:C). The induction profile of the ISGs characterized in our bat model does not match that obtained by Zhou and colleagues (Zhou et al., 2013) on lung cells from *P. alecto* after poly(I:C) transfection. Although we also showed an induction of OAS1 and PKR following poly(I:C) transfection, they found higher and earlier induction of PKR as compared to OAS1. The discrepancy between this previous study and our results is probably due to intrinsic factors specific to the bat species leading to distinct expression profiles. Indeed, it has recently been reported that the up-regulation of ISGs was not only lineage- but also species-specific when comparing the megabats and the microbats (Shaw et al., 2017). In addition, the higher induction of OAS1a and OAS1b, as compared to ADAR1 and PKR, may be due to the presence of multiple IFN-stimulated response elements (ISREs) in the OAS1 promoter, as described in *P. alecto* (Zhou et al., 2013). This can conveniently allow bats to mount a more robust expression of OAS1 rather than of ADAR1 or PKR. In fact, we previously showed an induction of expression of IFN β in poly(I:C)-transfected FLuDero, reaching a peak at 24 h (Sarkis et al., 2018). Interestingly, the expression kinetics of the three ISGs characterized was similar to that of IFN β , suggesting that these proteins are induced in an IFN β -dependent manner, acting on the ISREs present in their promoters. It is conceivable that the similar

expression levels of ADAR and PKR observed at the RNA level are also observed at the protein level. Accordingly, given the negative regulation of PKR by ADAR, as indicated above, an active PKR might not be in sufficient quantity to exert a proper antiviral action. Therefore, the higher induction of OAS observed, as compared to PKR and ADAR, tends to show the particular importance of the OAS proteins in the antiviral action of IFN in bats. Another interesting observation was the differential expression of OAS1a and OAS1b in our cellular model after poly(I:C) transfection. Interestingly, this feature has already been described in mice (Pulit-Penaloza et al., 2012). It has been shown that the ISRE as well as the overlapping STAT sites are required for Oas1a promoter induction by IFN- β while Oas1b expression requires only the ISRE. Furthermore, while OAS1a requires both STAT1 and STAT2 for its upregulation by IFN- β , OAS1b only requires STAT2. Therefore, analyzing the promoter region of these two paralogs should clarify their differential expression in our bat species.

We found that the basal transcription level of the ISGs characterized was elevated in the FLuDero cells. This observation is consistent with previous results showing that ISGs are constitutively expressed in bats and suggests that bat cells are equipped to mount a rapid defense against viral infections (Cruz-Rivera et al., 2018; Shaw et al., 2017; Zhou et al., 2016). In addition to their already significant levels of expression in the absence of treatment, these ISGs can be further enhanced in response to poly(I:C). Together these results tend to prove that these ISGs are functional and must play an important antiviral role in *D. rotundus*. Further work is therefore necessary to elucidate their role during viral infection, explore the mechanisms underlying their unique expression profiles and determine whether or not they contribute to the unique ability of bats to manage viral infection.

Acknowledgments

This study was conducted within the CAROLIA and BATIMMUNE programs. CAROLIA was supported by European funds (ERDF/FEDER) and assistance from the Région Guyane and the Direction Régionale pour la Recherche et la Technologie. BATIMMUNE was funded by the Institut Pasteur through a Transversal Research Program (PTR499). This study also received a European Commission "REGPOT-CT-2011-285837-STRonGer" grant within the FP7 and an "Investissement d'Avenir" grant managed by the Agence Nationale de la Recherche (CEBA, Ref. ANR-10-LABEX-25-01). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests

The authors declare that they have no competing interests.

References

- Arnaud, N., Dabo, S., Maillard, P., Budkowska, A., Kalliampakou, K.I., Mavromara, P., Garcin, D., Hugon, J., Gatignol, A., Akazawa, D., Wakita, T., Meurs, E.F., 2010. Hepatitis C Virus Controls Interferon Production through PKR Activation. *PLoS ONE* 5. <https://doi.org/10.1371/journal.pone.0010575>
- Athanasiadis, A., Placido, D., Maas, S., Brown, B.A., Lowenhaupt, K., Rich, A., 2005. The Crystal Structure of the Z β Domain of the RNA-editing Enzyme ADAR1 Reveals Distinct Conserved Surfaces Among Z-domains. *J. Mol. Biol.* 351, 496–507. <https://doi.org/10.1016/j.jmb.2005.06.028>
- Baker, M.L., Schountz, T., Wang, L.-F., 2013. Antiviral immune responses of bats: a review. *Zoonoses Public Health* 60, 104–116. <https://doi.org/10.1111/j.1863-2378.2012.01528.x>
- Bazak, L., Haviv, A., Barak, M., Jacob-Hirsch, J., Deng, P., Zhang, R., Isaacs, F.J., Rechavi, G., Li, J.B., Eisenberg, E., Levanon, E.Y., 2014. A-to-I RNA editing occurs at over a hundred million genomic sites, located in a majority of human genes. *Genome Res.* 24, 365–376. <https://doi.org/10.1101/gr.164749.113>
- Bonnet, M.C., Daurat, C., Ottone, C., Meurs, E.F., 2006. The N-terminus of PKR is responsible for the activation of the NF- κ B signaling pathway by interacting with the IKK complex. *Cell. Signal.* 18, 1865–1875. <https://doi.org/10.1016/j.cellsig.2006.02.010>
- Borden, E.C., Sen, G.C., Uze, G., Silverman, R.H., Ransohoff, R.M., Foster, G.R., Stark, G.R., 2007. Interferons at age 50: past, current and future impact on biomedicine. *Nat. Rev. Drug Discov.* 6, 975–990. <https://doi.org/10.1038/nrd2422>
- Clerzius, G., Gélinas, J.-F., Daher, A., Bonnet, M., Meurs, E.F., Gatignol, A., 2009. ADAR1 Interacts with PKR during Human Immunodeficiency Virus Infection of Lymphocytes and Contributes to Viral Replication. *J. Virol.* 83, 10119–10128. <https://doi.org/10.1128/JVI.02457-08>
- Cruz-Rivera, P.C.D.L., Kanchwala, M., Liang, H., Kumar, A., Wang, L.-F., Xing, C., Schoggins, J.W., 2018. The IFN Response in Bats Displays Distinctive IFN-Stimulated Gene Expression Kinetics with Atypical RNASEL Induction. *J. Immunol.* 200, 209–217. <https://doi.org/10.4049/jimmunol.1701214>
- Dabo, S., Meurs, E.F., 2012. dsRNA-Dependent Protein Kinase PKR and its Role in Stress, Signaling and HCV Infection. *Viruses* 4, 2598–2635. <https://doi.org/10.3390/v4112598>
- Der, S.D., Zhou, A., Williams, B.R., Silverman, R.H., 1998. Identification of genes differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays. *Proc. Natl. Acad. Sci. U. S. A.* 95, 15623–15628.
- Dey, M., Cao, C., Sicheri, F., Dever, T.E., 2007. Conserved Intermolecular Salt Bridge Required for Activation of Protein Kinases PKR, GCN2, and PERK. *J. Biol. Chem.* 282, 6653–6660. <https://doi.org/10.1074/jbc.M607897200>

- Doria, M., Neri, F., Gallo, A., Farace, M.G., Michienzi, A., 2009. Editing of HIV-1 RNA by the double-stranded RNA deaminase ADAR1 stimulates viral infection. *Nucleic Acids Res.* 37, 5848–5858. <https://doi.org/10.1093/nar/gkp604>
- Elkhateeb, E., Tag-El-Din-Hassan, H.T., Sasaki, N., Torigoe, D., Morimatsu, M., Agui, T., 2016. The role of mouse 2',5'-oligoadenylate synthetase 1 paralogs. *Infect. Genet. Evol.* 45, 393–401. <https://doi.org/10.1016/j.meegid.2016.09.018>
- Fierro-Monti, I., Mathews, M.B., 2000. Proteins binding to duplexed RNA: one motif, multiple functions. *Trends Biochem. Sci.* 25, 241–246. [https://doi.org/10.1016/S0968-0004\(00\)01580-2](https://doi.org/10.1016/S0968-0004(00)01580-2)
- Garaigorta, U., Chisari, F.V., 2009. Hepatitis C Virus Blocks Interferon Effector Function by Inducing PKR Phosphorylation. *Cell Host Microbe* 6, 513–522. <https://doi.org/10.1016/j.chom.2009.11.004>
- García, M.A., Meurs, E.F., Esteban, M., 2007. The dsRNA protein kinase PKR: Virus and cell control. *Biochimie* 89, 799–811. <https://doi.org/10.1016/j.biochi.2007.03.001>
- Guénet, J.L., Benavides, F., Panthier, J., Montagutelli, X., 2015. *Genetics of the Mouse*. Springer Berlin Heidelberg. XVII Chapters, 408 pages. <https://doi.org/10.1007/978-3-662-44287-6>.
- George, C.X., Samuel, C.E., 1999. Characterization of the 5'-flanking region of the human RNA-specific adenosine deaminase ADAR1 gene and identification of an interferon-inducible ADAR1 promoter. *Gene* 229, 203–213.
- Ghosh, A., Sarkar, S.N., Guo, W., Bandyopadhyay, S., Sen, G.C., 1997. Enzymatic activity of 2'-5'-oligoadenylate synthetase is impaired by specific mutations that affect oligomerization of the protein. *J. Biol. Chem.* 272, 33220–33226.
- Goodman, R.A., Macbeth, M.R., Beal, P.A., 2012. ADAR proteins: structure and catalytic mechanism. *Curr. Top. Microbiol. Immunol.* 353, 1–33. https://doi.org/10.1007/82_2011_144
- He, X., Korytář, T., Zhu, Y., Pikula, J., Bandouchova, H., Zukal, J., Köllner, B., 2014. Establishment of Myotis myotis cell lines--model for investigation of host-pathogen interaction in a natural host for emerging viruses. *PloS One* 9, e109795. <https://doi.org/10.1371/journal.pone.0109795>
- Herbert, A., Alfken, J., Kim, Y.-G., Mian, I.S., Nishikura, K., Rich, A., 1997. A Z-DNA binding domain present in the human editing enzyme, double-stranded RNA adenosine deaminase. *Proc. Natl. Acad. Sci. U. S. A.* 94, 8421–8426.
- Justesen, J., Hartmann, R., Kjeldgaard, N.O., 2000. Gene structure and function of the 2'-5'-oligoadenylate synthetase family. *Cell. Mol. Life Sci. CMLS* 57, 1593–1612.
- Kim, Y.-G., Lowenhaupt, K., Schwartz, T., Rich, A., 1999. The Interaction between Z-DNA and the Zab Domain of Double-stranded RNA Adenosine Deaminase Characterized Using Fusion Nucleases. *J. Biol. Chem.* 274, 19081–19086. <https://doi.org/10.1074/jbc.274.27.19081>

- Kristiansen, H., Gad, H.H., Eskildsen-Larsen, S., Despres, P., Hartmann, R., 2011. The oligoadenylate synthetase family: an ancient protein family with multiple antiviral activities. *J. Interferon Cytokine Res. Off. J. Int. Soc. Interferon Cytokine Res.* 31, 41–47. <https://doi.org/10.1089/jir.2010.0107>
- Lai, F., Drakas, R., Nishikura, K., 1995. Mutagenic Analysis of Double-stranded RNA Adenosine Deaminase, a Candidate Enzyme for RNA Editing of Glutamate-gated Ion Channel Transcripts. *J. Biol. Chem.* 270, 17098–17105. <https://doi.org/10.1074/jbc.270.29.17098>
- Letunic, I., Doerks, T., Bork, P., 2015. SMART: recent updates, new developments and status in 2015. *Nucleic Acids Res.* 43, D257–260. <https://doi.org/10.1093/nar/gku949>
- Levy, D.E., Marié, I., Smith, E., Prakash, A., 2002. Enhancement and diversification of IFN induction by IRF-7-mediated positive feedback. *J. Interferon Cytokine Res. Off. J. Int. Soc. Interferon Cytokine Res.* 22, 87–93. <https://doi.org/10.1089/107999002753452692>
- Liu, Y., Samuel, C.E., 1996. Mechanism of interferon action: functionally distinct RNA-binding and catalytic domains in the interferon-inducible, double-stranded RNA-specific adenosine deaminase. *J. Virol.* 70, 1961–1968.
- Malathi, K., Dong, B., Gale, M., Silverman, R.H., 2007. Small self-RNA generated by RNase L amplifies antiviral innate immunity. *Nature* 448, 816–819. <https://doi.org/10.1038/nature06042>
- Malathi, K., Paranjape, J.M., Bulanova, E., Shim, M., Guenther-Johnson, J.M., Faber, P.W., Eling, T.E., Williams, B.R.G., Silverman, R.H., 2005. A transcriptional signaling pathway in the IFN system mediated by 2'-5'-oligoadenylate activation of RNase L. *Proc. Natl. Acad. Sci. U. S. A.* 102, 14533–14538. <https://doi.org/10.1073/pnas.0507551102>
- Mannion, N., Arieti, F., Gallo, A., Keegan, L.P., O'Connell, M.A., 2015. New Insights into the Biological Role of Mammalian ADARs; the RNA Editing Proteins. *Biomolecules* 5, 2338–2362. <https://doi.org/10.3390/biom5042338>
- Marié, I., Blanco, J., Rebouillat, D., Hovanessian, A.G., 1997. 69-kDa and 100-kDa isoforms of interferon-induced (2'-5')oligoadenylate synthetase exhibit differential catalytic parameters. *Eur. J. Biochem.* 248, 558–566.
- McCormack, S.J., Ortega, L.G., Doohan, J.P., Samuel, C.E., 1994. Mechanism of Interferon Action Motif I of the Interferon-Induced, RNA-Dependent Protein Kinase (PKR) Is Sufficient to Mediate RNA-Binding Activity. *Virology* 198, 92–99. <https://doi.org/10.1006/viro.1994.1011>
- Meurs, E., Chong, K., Galabru, J., Thomas, N.S.B., Kerr, I.M., Williams, B.R.G., Hovanessian, A.G., 1990. Molecular cloning and characterization of the human double-stranded RNA-activated protein kinase induced by interferon. *Cell* 62, 379–390. [https://doi.org/10.1016/0092-8674\(90\)90374-N](https://doi.org/10.1016/0092-8674(90)90374-N)

- Meurs, E.F., Watanabe, Y., Kadereit, S., Barber, G.N., Katze, M.G., Chong, K., Williams, B.R., Hovanessian, A.G., 1992. Constitutive expression of human double-stranded RNA-activated p68 kinase in murine cells mediates phosphorylation of eukaryotic initiation factor 2 and partial resistance to encephalomyocarditis virus growth. *J. Virol.* 66, 5805–5814.
- Moratelli, R., Calisher, C.H., 2015. Bats and zoonotic viruses: can we confidently link bats with emerging deadly viruses? *Mem. Inst. Oswaldo Cruz* 110, 1–22. <https://doi.org/10.1590/0074-02760150048>
- Pulit-Penalzoza, J.A., Scherbik, S.V., Brinton, M.A., 2012. Activation of Oas1a gene expression by type I IFN requires both STAT1 and STAT2 while only STAT2 is required for Oas1b activation. *Virology* 425, 71–81. <https://doi.org/10.1016/j.virol.2011.11.025>
- Saraste, M., Sibbald, P.R., Wittinghofer, A., 1990. The P-loop--a common motif in ATP- and GTP-binding proteins. *Trends Biochem. Sci.* 15, 430–434.
- Sarkar, S.N., Miyagi, M., Crabb, J.W., Sen, G.C., 2002. Identification of the substrate-binding sites of 2'-5'-oligoadenylate synthetase. *J. Biol. Chem.* 277, 24321–24330. <https://doi.org/10.1074/jbc.M110202200>
- Sarkis, S., Lise, M.-C., Darcissac, E., Dabo, S., Falk, M., Chaulet, L., Neuveut, C., Meurs, E.F., Lavergne, A., Lacoste, V., 2018. Development of molecular and cellular tools to decipher the type I IFN pathway of the common vampire bat. *Dev. Comp. Immunol.* 81, 1–7. <https://doi.org/10.1016/j.dci.2017.10.023>
- Sen, G.C., 2001. Viruses and Interferons. *Annu. Rev. Microbiol.* 55, 255–281. <https://doi.org/10.1146/annurev.micro.55.1.255>
- Shaw, A.E., Hughes, J., Gu, Q., Behdenna, A., Singer, J.B., Dennis, T., Orton, R.J., Varela, M., Gifford, R.J., Wilson, S.J., Palmarini, M., 2017. Fundamental properties of the mammalian innate immune system revealed by multispecies comparison of type I interferon responses. *PLOS Biol.* 15, e2004086. <https://doi.org/10.1371/journal.pbio.2004086>
- Silverman, R.H., 2003. Implications for RNase L in Prostate Cancer Biology. *Biochemistry (Mosc.)* 42, 1805–1812. <https://doi.org/10.1021/bi027147i>
- Tag-El-Din-Hassan, H.T., Sasaki, N., Moritoh, K., Torigoe, D., Maeda, A., Agui, T., 2012. The chicken 2'-5' oligoadenylate synthetase A inhibits the replication of West Nile virus. *Jpn. J. Vet. Res.* 60, 95–103.
- Tamura, K., Stecher, G., Peterson, D., Filipinski, A., Kumar, S., 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729. <https://doi.org/10.1093/molbev/mst197>
- Williams, B.R.G., 2001. Signal Integration via PKR. *Sci STKE* 2001, re2-re2. <https://doi.org/10.1126/stke.2001.89.re2>
- Xiang, Y., Wang, Z., Murakami, J., Plummer, S., Klein, E.A., Carpten, J.D., Trent, J.M., Isaacs, W.B., Casey, G., Silverman, R.H., 2003. Effects of RNase L mutations associated with prostate cancer on apoptosis induced by 2',5'-oligoadenylates. *Cancer Res.* 63, 6795–6801.

Yamamoto, Y., Sono, D., Sokawa, Y., 2000. Effects of specific mutations in active site motifs of 2',5'-oligoadenylate synthetase on enzymatic activity. *J. Interferon Cytokine Res. Off. J. Int. Soc. Interferon Cytokine Res.* 20, 337–344. <https://doi.org/10.1089/1079990000312496>

Zhou, P., Cowled, C., Wang, L.-F., Baker, M.L., 2013. Bat Mx1 and Oas1, but not Pkr are highly induced by bat interferon and viral infection. *Dev. Comp. Immunol.* 40, 240–247. <https://doi.org/10.1016/j.dci.2013.03.006>

Zhou, P., Tachedjian, M., Wynne, J.W., Boyd, V., Cui, J., Smith, I., Cowled, C., Ng, J.H.J., Mok, L., Michalski, W.P., Mendenhall, I.H., Tachedjian, G., Wang, L.-F., Baker, M.L., 2016. Contraction of the type I IFN locus and unusual constitutive expression of IFN- α in bats. *Proc. Natl. Acad. Sci. U. S. A.* 113, 2696–2701. <https://doi.org/10.1073/pnas.1518240113>

Figure 1. Amino acid alignment of *D. rotundus* OAS1 paralogs and other mammalian species. The two important signature domains, NTP-transf_2 and OAS1_C in *D. rotundus* OAS1a and OAS1b are highlighted with solid arrows. Amino acid sequence motifs important for the enzyme activity are shown in boxes. The orange rectangle represents the Serine residue in the putative helical turn, the adjacent Serine, which is either Serine or Threonine (Rune Hartmann et al., 2003 Molecular cell). Inverted triangle: one of the signature features of the nucleotidyl transferase superfamily is a conserved triad of aspartic acids predicted to coordinate the catalytically active magnesium ions. The multi-alignment was performed with the ClustalW program using default parameters. Dashes indicate gaps in the amino acid alignment. Abbreviations: MIOAS1: *Myotis lucifigus* OAS1 (XM_014464612), MIOAS1I: *Myotis lucifigus* OAS1-like (XM_014462497), HsOAS1: *Homo sapiens* OAS1 (BC000562), SsOAS1: *Sus scrofa* (NM_214303), MmOAS1a-h: *Mus musculus* OAS1a-h (XM_006530330, BC012877, NM_033541, BC052835, BC053721, BC116433, BC018470, NM_145228).

Figure 2. Amino acid alignments of *D. rotundus* ADAR1 and PKR with their homologs from other mammalian species. a) All motifs required for ADAR1 activity are highlighted with solid arrows, including two Z α domains at the N-terminus, followed by three double-stranded RNA-binding motifs (dsRBM), dsRBM1, 2 and 3, and a catalytic deaminase domain at its C-terminus. **b)** PKR contains two dsRBM highlighted in yellow (dsRBM1) and light gray (dsRBM2) followed by a basic region (dark gray) and subdomain sequences in boxes found in the catalytic region highlighted with solid arrows, green represents the repeat sequence in human PKR not present in *D. rotundus*, other bat species and mice PKR. An insert region at the C-terminal of the subdomain IV is highlighted in blue. The multi-alignments were performed with the ClustalW program using default parameters. Dashes indicate gaps in the amino acid alignments. Abbreviations: PaADAR1/PKR: *Pteropus alecto*

ADAR1/PKR (XM_015598363/NM_0012910168), HsADAR1/PKR: *Homo sapiens* ADAR1/PKR (NM_001111/XM_011532987), MmADAR1/PKR: *Mus musculus* ADAR1/PKR (NM_019655/NM_011163).

Figure 3. Phylogenetic trees generated from amino acid sequence alignments of OAS1, ADAR1 and PKR from *D. rotundus* and other mammals. Phylogenetic trees were generated based on (a) the alignment of partial amino acid sequence of the mammalian OAS1, and the full-length protein sequence of (b) ADAR1 and (c) PKR. The trees were inferred using the Bayesian method with the JTT + G model. Sequence identifiers include the NCBI accession number. Posterior probabilities of the Bayesian analysis (>50%) are shown next to each node. The scale bar indicates amino acid substitutions per site.

Figure 4. Poly(I:C) transfection induces expression of ISGs. Time-course fold induction of (a) OAS1a, (b) OAS1b, (c) ADAR1 and (d) PKR in FLuDero cells upon poly(I:C) transfection. ISGs mRNA were assessed by RTqPCR. Data were normalized against β -actin. Cells were stimulated with 20 μ g/mL of poly(I:C) and harvested at the indicated time points. Data represent the mean values of three separate experiments, and the error bars represent standard deviations (SD).

Figure 5. Quantitative mRNA expression of OAS1a, OAS1b, ADAR1 and PKR. Time course mRNA expression levels of (red) OAS1a, (green) OAS1b, (blue) ADAR1 and (purple) PKR were assessed by RTqPCR in FLuDero cells upon poly(I:C) transfection. Cells were stimulated with 20 μ g/mL of poly(I:C) and harvested at the indicated time points. The expression was normalized to the housekeeping gene β -actin. Data represent the mean values of three separate experiments.

Figure 1.

A

DrADAR1	----NSRQCITLKNYHNFLQRYEHSRLRQGL--GRSYVNSFQLOQIEFLKQGLFVP	55
PaADAR1	----MEFRQCFLRYPS----PFGYELGHQK--GLGAYPNRQLOQIEFLKQGLFVP	51
HaADAR1	----MFRQCYSLSYTHFPQGYEHRQYQPP--GGSSFSFSLKQIEFLKQGLFVP	55
MmADAR1	MSQGFPGTCTVPHQTQSYLDPSSHEHSKRWYPPQPGSGPSYFSFQLQOIEFLKGLRFLFVP	60
DrADAR1	LIRKQAPVFLPLPEFWRFP---GPPARGRQPEINGIPRGVPLRSQALQSSTQPPFPQG	112
PaADAR1	LIGQTSLSLFPYFGLMWRFP---GPPARGRQSEMGVPRPVLRSQAQGRGFPFLCPRG	108
HaADAR1	VIGQTSLSLFLGLRWFVLLASSTRGRVDYGVNPGVHGLGQCLQMGQFHSFPG	115
MmADAR1	LIGIQTSLSLFPFLGHWFRFP---GPPAQDRGLEIMWFPSVTLRQGFHIGPPLFPFHS	117
DrADAR1	RIPFWRGVRLSSHFQRLRQDQQRVLELLQELGAGQVTTAQDLAKLQAPKEENRI	172
PaADAR1	KTRFWRGVDTLSARLQELYISQEQORVLELLQELGAGKAITAHLARRLQASKEVNSV	168
HaADAR1	RSLPQRGVCLSSHFQELSYQDQQRILKFLLELGEKATTADHLGKGLGTPKKEINRV	175
MmADAR1	ROTFRWAGDGLCSHFRELISISQGVQVLRNLEELGKATTAVHLARELIPFRDINRI	177
DrADAR1	LYSLAKKGLHQEAKFPLRLAASVPRWNQSQASRSDSCSGAPHSDBLETEDRD--	230
PaADAR1	LYSLAKKGLLREGVTFPLRRAVSAQANDRQQAADQWSRPAVSDGLETEDRN--	226
HaADAR1	LYSLAKKGLQEQACTPFLRLAVSTQAWNHSQGVPRDSGGAHSDGLETEDRST	235
MmADAR1	LYSLKKGKLIHRGKGFPLSLVPLSQAWTOPGVVNPDSCTQEPFRGEGLDSEDDG--	235
DrADAR1	-----PPGLEDPTEAD	243
PaADAR1	-----PTSGVEGPEPFLD	239
HaADAR1	SVSEDLLEFFIAVSAQAMNHSQGVPRPDSHSQSGSPNSDPGLEPEDSNSTSALEDPLEFLD	295
MmADAR1	-----PASLEGPSPFLD	248
DrADAR1	-----Za	
PaADAR1	MAETKEICHYLNVFKSSALNKNIGLTKARDVNAVILDLERQGVYRGQTFPTTWYL	303
HaADAR1	MAETKEICELYHAPSSALNKNIGLTKARDVNAVILDLERQGVYRGQTFPTTWYL	299
MsADAR1	MAETKEICDYLNVDSGALNKNIGLTKARDNAVILDMERQGVYRGQTFPTTWYL	355
MmADAR1	MAETKEICDYLNVDSGALNKNIGLTKARDVTVLILDLERQGVYRGQATPTTWYL	308
DrADAR1	TDKKRERIQLRNNTSTLETTAAIPETQRNVELPCNLASDAS--HNVTGNLENGQEP	362
PaADAR1	TDKKRERIQIKNNVNSVETTTQAAVPETRKSEVPPHNSAVPRAS--DETAAPKANGQGL	358
HaADAR1	TDKKRERIQIKNNVNSVETTPAALPETKNNSELTCTIPMSASNNVTVTEKVENQEP	415
MmADAR1	TDKKRERLQKKRSTHSAFAPPTTAVPEATRSFPPFACHFFPAGASSVAASKRVENQEP	368
DrADAR1	VIKLES--QGVTFEPKFPVRDNGSPKTVYDFNGQWATDDIPDLNLSIAAPGEFR	421
PaADAR1	VILKLESPEVVTTFVTPKFPVRDNGSPKTVYDFNGQWATDDIPDLNLSIAAPGEFR	418
HaADAR1	VIKLENR--QEARPEARLKFPVHYNGSPKAGYDFNGQWATDDIPDLNLSIAAPGEFR	474
MmADAR1	ATKLESR--HEARFGRMLRPHAYHNGSPKAGYVASENGQWATDDIPDLNLSIHTAPGEFR	427
DrADAR1	-----dsBm1	
PaADAR1	AIMEMSPFYSHGLFRCSFYKLTQCOLNPIISGLLEYAQFASQTCFNLIQSGSPHPEFR	481
HaADAR1	AIMEMSPFYSHWRKCSFYKLTQCOLNPIISGLLEYAQFASQTCFNLIQSGSPHPEFR	478
MmADAR1	AIMEMSPFYSHGLFRCSFYKLTQCOLNPIISGLLEYAQFASQTCFNLIQSGSPHPEFR	534
DrADAR1	AIMEMSPFYSFTLFRCSFYKLTQCOLNPIISGLLEYAQFASQTCFNLIQSGSPHPEFR	487
PaADAR1	FKFQVVMINGREFFPFAEAGSKKVAQDAATKNAALLEEAKARDGSPESHSVCSEKAG	541
HaADAR1	FKFQVVISGREFFPFAEAGSKKVAQDAATKANTILLEAKARDGSRPESHYCTERESE	538
MmADAR1	FKFQVVMINGREFFPFAEAGSKKVAQDAAMKANTILLEAKARDGSKSESSEHYCTERESE	594
DrADAR1	FKFQVVMINGREFFPFAEAGSKKVAQDAVNAAILLEAKARDGSPQDLSHCMEDESE	547
PaADAR1	-----dsBm2	
HaADAR1	ETASQPTTFSATSFSGKNPVTTILECVHKLGSCEFRLLSREGPAHDFFQVCVAMGN	601
MmADAR1	ETASQPTTFSATSFSGKNPVTTILECVHKLGSCEFRLLSREGPAHDFFQVCVAMGT	597
DrADAR1	KTASQPTTFSATSFSGKSPTTTILECMHKLGSCEFRLLSREGPAHEFFQVCVAVGA	654
PaADAR1	KTASQPTTFSATSFSGKSPTTTILECMHKLGSCEFRLLSREGPAHDFFQVCVAVGA	607
MmADAR1	KPAEQAPSSSATLSFGKSPVTTILECMHKLGSCEFRLLSREGPAHDFFQVCVAVGA	607
DrADAR1	HTFPTASAPSKKVAQMAEAMKALHGEATSTSPEDQGSNTIESFONLESAMNKVR	659
PaADAR1	HTFPTASAPSKKAAQMAEAMKALHGEATDLSPENQGSASSEFQDLSAVPSKVR	657
HaADAR1	OTTFVSAPSKKVAQMAEAMKALHGEATNSMADQPEGMISSELNLESMMNKVR	714
MmADAR1	OTTFVSAPSKKVAQMAEAMKALGEAASE--DDQSGNANTISL--ESMAFLR	663
DrADAR1	-----dsBm3	
PaADAR1	RIGELVRYLNTNPGVGLLEYARSHGFAAFKFLVDQSGPHEPRFVYQAKVGWRFFAVCA	721
HaADAR1	RIGELVRYLNTNPGVGLLEYARSHGFAAFKFLVDQSGPHEPRFVYQAKVGWRFFAVCA	717
MmADAR1	RIGELVRYLNTNPGVGLLEYARSHGFAAFKFLVDQSGPHEPRFVYQAKVGWRFFAVCA	774
MmADAR1	RIGELVRYLNTNPGVGLLEYARSHGFAAFKFLVDQSGPHEPRFVYQAKVGWRFFAVCA	723

DrADAR1	HSKKQKGQEAADALRVLIGENEKAERMGTTEL-----P	755
PaADAR1	HSKKQKGQEAADALRVLIGESEKAERMGTTEL-----P	751
HaADAR1	HSKKQKGQEAADALRVLIGENEKAERMGTTEVTPVTGASLRRTMLLSRSPEAQPKTLP	834
MmADAR1	HSKKQKGQDAADALRVLIGESEKAELGLFAEL-----P	757
DrADAR1	LTGSTFHDQIAMLSHRCFNALTNSFQPSLLGRKILAAIIMKKDSEDLGVVVS LGTGNRVC	815
PaADAR1	LTGSTFHDQIAMLSHRCFNALTNSFQPSLLGRKILAAIIMKKDSEDLGVVVS LGTGNRVC	811
HaADAR1	LTGSTFHDQIAMLSHRCFNLTNSFQPSLLGRKILAAIIMKKDSEDMGVVVS LGTGNRVC	894
MmADAR1	LSGSTFHDQIAMLSHRCFNALTNSFQPSLLGRKILAAIIMKRDPEDMGVVVS LGTGNRVC	817
DrADAR1	KGDSLSLKGETVNDCHAEIISRGGFIRFLYSELMKYSPTAKDSIFEPAKGGEKLQIKKT	875
PaADAR1	KGDSLSLKGETVNDCHAEIISRGGFIRFLYSELMKYNFPTAKDSIFEPAKGGEKLQIKKT	871
HaADAR1	KGDSLSLKGETVNDCHAEIISRGGFIRFLYSELMKYNSTAKDSIFEPAKGGEKLQIKKT	954
MmADAR1	KGDSLSLKGETVNDCHAEIISRGGFIRFLYSELMKYNHTAKNSIFEPLARGGEKLQIKKT	877
DrADAR1	VSFHLYISTAPCGDGLFDKSCSDRAVESTDSRHPVPFENPKQGLRRTKVENGEGTIPVE	935
PaADAR1	VSFHLYISTAPCGDGLFDKSCSDRAVESTDRHPVPFENPKQGLRRTKVENGEGTIPVE	931
HaADAR1	VSFHLYISTAPCGDGLFDKSCSDRAVESTSRHPVPFENPKQGLRRTKVENGEGTIPVE	1014
MmADAR1	VSFHLYISTAPCGDGLFDKSCSDRAVESTSRHPVPFENPKQGLRRTKVENGEGTIPVE	937
catalytic deaminase domain		
DrADAR1	SSDIVPTWDGIRLGERLRTMSCSDKILRWNVLGLQALLTHFLQPVYLKSVTLGYLFSQG	995
PaADAR1	SSDIVPTWDGIRLGERLRTMSCSDKILCWNVLGLQALLTHFLQPVYLKSVTLGYLFSQG	991
HaADAR1	SSDIVPTWDGIRLGERLRTMSCSDKILRWNVLGLQALLTHFLQPIVYLKSVTLGYLFSQG	1074
MmADAR1	SSDIVPTWDGIRLGERLRTMSCSDKILRWNVLGLQALLTHFLQPVYLKSVTLGYLFSQG	997
DrADAR1	HLTRAICCRVTRDGSAFEDGLRHPPFIVNHPKVGRVSYDSKRQSGTKTETSNNWCLADGY	1055
PaADAR1	HLTRAICCRVTRDGSAFEDGLQPPFTVNHPKVGRVSYDSKRQSGTKTETSNNWCLADGC	1051
HaADAR1	HLTRAICCRVTRDGSAFEDGLRHPPFIVNHPKVGRVSYDSKRQSGTKTETSNNWCLADGY	1134
MmADAR1	HLTRAICCRVTRDGFADFDGLRYPPFIVNHPKVGRVSYDSKRQSGTKTETSNNWCLADGY	1057
DrADAR1	DLEILDGTRGTVDGPRNELSRVSKKNIFLLFKRLCSFRARRDLLKLSYGEAKKAARDYET	1115
PaADAR1	DLEILDGTRGTVDGPRNELSRVSKKNIFLLFKRLCSFRARRDLLKLSYGEAKKAARDYEA	1111
HaADAR1	DLEILDGTRGTVDGPRNELSRVSKKNIFLLFKRLCSFRARRDLLKLSYGEAKKAARDYET	1194
MmADAR1	DLEILDGTRGTVDGPKELSRVSKKNIFLQFKRLCSFRARRDLLQLSYGEAKKAARDYDL	1117
DrADAR1	AKNYFKKSLKDMGYGNWISKPOEEKNFYLCVP	1147
PaADAR1	AKNYFKKGLKDMGYGNWISKPOEEKNFYLCVP	1143
HaADAR1	AKNYFKKGLKDMGYGNWISKPOEEKNFYLCVP	1226
MmADAR1	AKNYFKKSLRDMGYGNWISKPOEEKNFYLCVP	1149

B		dsRBM1	
DrPKR	MADNHIRSFICAIWKYSCKNNVNVYRELERTGPPHDSIKKFOVILDEREYPAEGKSK	60	
PaPKR	MADDLSGFFLEELNKYRKNNVEIEYRELKRGPPHLSRTFQVVIGEREFPAEGKSK	60	
HePKR	MAGDLSAGFPMRELNTYRKQGVVLKYQELFNSGPPHDSRTFQVVIIDGEREFPAGEGRSK	60	
MmPKR	MASDTP-GYMDKLNKYRMHGVAITYKELSTSGPPHDSRTFQVVIDEKEFPAGEGRSK	59	
DrPKR	REAKDAAKLAFAELKE-KEEVSASVLLTDTSDDLMSGNYVGQINRLTQKANLSVNYQQ	119	
PaPKR	RGAKNAAKLAVEIINEEKEEVSSLSTTDTSDLSIGNYIGLVNRTAQKALPVNYQL	119	
HePKR	REAKNAAKLAVEIINKEKAVSPLLLTTNSSEGLMSGNYIGLVNRTAQKRLTVNYEQ	120	
MmPKR	QEARNAAKLAVDILDN---ENKVDCHTSASEQGLFVGNVIGLVNRTAQKRLTVNYEQ	115	
dsRBM2		basic	
DrPKR	HESYTNQPERFSCCKIIGQKEYSSAVGSTQKQAKQLAAKLAYEKILAEISSSTKDDSVLLG	179	
PaPKR	GLG-AGEPGRFYTCIIGQKYNVAVGSTQKQAKQLAAKLAYEQILSAKTS-VDDLASLG	177	
HePKR	CASGVHGFEGFHYCKRMGQKEYSIGTGSTQKQAKQLAAKLAYLQILSEETSVKSDYLSGG	180	
MmPKR	CEPNSLEFQRICKCKIGQTMGTGSGVTKQKQAKQLAAKLAYQELKSEPK----TAGTS	171	
DrPKR	SSPAGPGDYKGSFRT--GGSESPSENGFLENGESNDNSGSFINTPSSSPDLRFSSGR	237	
PaPKR	SPSAASSDYGSNSMTNMCDSESPCENGFLANGSERNDNSDFSNLSSPSVSVRRSLGR	237	
HePKR	S-FATTICESQSNLVTSTLASESSSEDDYSDTSEINNSDS-LNYSLLANGLRNQRM	238	
MmPKR	S-----SVVT-----ST--FSGFSSSSMTSNGVQSAPGSPSSENVVTNGLGE	213	
region			
DrPKR	GR--RSIAAKFDSPLIGRNKYTMNERFIKDFSEITQIGGGYGRVFKAKHRIDGKTVIK	295	
PaPKR	TRIFNLAPHLNVLBAARNEYMDHRFLENLEITSLSGGHGVFKAKHRIDGKTVIK	297	
HePKR	AK--RSLAPFYDLPMMKETKYTVDKRYGMDFKIELSSGGFGVFKAKHRIDGKTVIK	296	
MmPKR	NK--RKSGVKVSPDQVQRNKYTLDAFNSDFEDIEELGGFGVFKAKHRIDGKTVIK	271	
DrPKR	RIKYNTKNKEKVERETIVLSALDHRNIVRYNNDGIDDAVESPRD-----	341	
PaPKR	RIKYNT--EKVVETKALAAISHNIVRYNNDGIDDAVESDNGR-----	342	
HePKR	RIKYNT--EKAEEETKALAKLDHNVIVHYNCWDGFTDPEISDSELSYDEPNKSK	353	
MmPKR	RIKYNT--EKAEEETQALAEIHNIVQYHSCWEGVDYDPEHMSMT-----	316	
catalytic kinase domain		VI	
DrPKR	----SPECLEFQMECEGTLQWINDREKGTMDLSLELFEQIVHGVNYIHSKALIHR	397	
PaPKR	--FESCTKCLFIQMECEGTLQWINKRRDQTKDKHLSLELFEQIAHGVNFIHSKALIHR	401	
HePKR	SSRKTKCLFIQMEPCDGTLEQWIEKRRGEKDKVLALSLFEQITKGVDIHSKALIHR	413	
MmPKR	-SRVTKCLFIQMEPCDGTLEQWMRNRNRQSKVDKALILDLYEQIVHGVYIHSKALIHR	375	
DrPKR	DLKPSNIFLVDTKQIKIGDFGIVTFOEYDEK-RTDRDGTFLYMSPEQISPEYGVNEVIF	456	
PaPKR	DLKPSNIFLVDTKKIKIGDFGIVTFLKNDK-RTNNRGTFLRYMSPEQISPEYGVNEVIF	460	
HePKR	DLKPSNIFLVDTKQVKGIGDFGIVTSLKNDK-RTRSKGTFLRYMSPEQISPDYGVKEVDLY	472	
MmPKR	DLKPSNIFLVDERHIKIGDFGIVTALENDGKSRTRTGTFLQYMSPEQLFKHYGVKEVDIF	435	
DrPKR	ALGLIIAELLYISPTHLEKSKIFQDLRSCVVP-DTYDDEKSLLEKLSAEPKKRINTSE	515	
PaPKR	PLGLIIAELLYISRTLETCKIFEDLKKGVL-DVFDDEKILLQKLSKDPKKRINTCE	519	
HePKR	ALGLIIAELLLHVCDTAFETSKFFDLDGDIIS-DIFDKKETLLQKLSKKPEDRINTSE	531	
MmPKR	ALGLIIAELLLHTCTESEKIKFFESLRKGFSDNDFDKKESLLEKLSKKPKDRINTSE	495	
DrPKR	ILKTLQEWKTVVREKMRHTY	536	
PaPKR	ILTTLEWKNVV-EKKKRNTC	539	
HePKR	ILRTLIVWKKSP-EKKKRNTC	551	
MmPKR	ILKTLAEWRNIS-EKKKRNTC	516	

Figure 2.

Figure 3.

Figure 4.

Figure 5.